



# ***Annona muricata* Linn leaves or *Curcuma longa* Linn Rhizomes Ameliorates Oxidative Stress Associated with Hypertension in Uninephrectomized Wistar Rats Daily Loaded with Sodium Chloride**

**Olayinka A. Oridupa<sup>1\*</sup>, Folusho B. Falade<sup>2</sup>, Ademola A. Oyagbemi<sup>3</sup>,  
Bukola A. Abegunde<sup>1</sup>, Precious C. Ekwem<sup>1</sup>, Adedoyin Badmus<sup>1</sup>  
and Temidayo O. Omobowale<sup>4</sup>**

<sup>1</sup>Department of Veterinary Pharmacology and Toxicology, University of Ibadan, Ibadan, Nigeria.

<sup>2</sup>Department of Veterinary Surgery and Radiology, University of Ibadan, Ibadan, Nigeria.

<sup>3</sup>Department of Veterinary Physiology and Biochemistry, University of Ibadan, Ibadan, Nigeria.

<sup>4</sup>Department of Veterinary Medicine, University of Ibadan, Ibadan, Nigeria.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author OAO designed the study, wrote the protocol, and wrote the final draft of the manuscript. Author FBF performed the nephrectomy and carried out the experiment. Author AAO performed the laboratory analysis, author BAA performed the laboratory analysis and managed some of the literature searches. Author PCE managed the laboratory animals, performed the statistical analysis and wrote the first draft, author AB managed the laboratory animals and some literature searches. Author TOO designed the study, wrote the protocol and managed the analyses of the study. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/EJMP/2018/v26i430100

### Editor(s):

- (1) Dr. Sabyasachi Chatterjee, Department of Biotechnology, Burdwan University, India.  
(2) Dr. Marcello Iriti, Professor, Plant Biology and Pathology, Department of Agricultural and Environmental Sciences, Milan State University, Italy.

### Reviewers:

- (1) Yongchun Zhu, Shenyang Normal University, China.  
(2) Uchendu, Mbah Okuwa Umudike, Abia State, Nigeria.  
(3) Muhammad Shahzad Aslam, Universiti Malaysia Perlis, Malaysia.  
Complete Peer review History: <http://www.sdiarticle3.com/review-history/47551>

**Original Research Article**

**Received 11 November 2018**  
**Accepted 22 February 2019**  
**Published 11 March 2019**

## **ABSTRACT**

**Aims:** Oxidative stress sequel to hypertension exacerbates the clinical condition and accelerates associated organopathies, therefore prevention is important. Traditionally in Nigeria, hypertension is treated with *Annona muricata* L. leaves or *Curcuma longa* L. rhizomes, two medicinal plants with antioxidant properties.

**Study Design:** This study assessed the effect of these plants on hypertension-induced oxidative stress in uninephrectomized Wistar rats daily loaded with 1% sodium chloride.

**Place and Duration of Study:** Department of Veterinary Pharmacology and Toxicology Experimental Animal House, University of Ibadan, Nigeria, between August and November 2017.

**Methodology:** Hypertensive rats were treated with methanol extracts of the plants for 42 days. Two other groups of hypertensive rats were treated with lisinopril or chlorothiazide. Blood pressure was monitored by non-invasive tail plethysmography using an electro-sphygmomanometer. Oxidative stress markers were determined in blood and tissue (heart, kidney and liver); GPX, GST, GSH, SOD, MDA and NO.

**Results:** Treatment of uninephrectomized rats with *A. muricata* or *C. longa* significantly ( $p < 0.0001$ ) decreased blood pressure and MDA, while elevating enzymatic and non-enzymatic antioxidant defense mechanisms of GST, GSH, GPx and SOD, comparable to normotensive rats. NO, the ubiquitous molecule required for basal vascular tone, myocardial contractility regulation and platelet adhesion prevention, was restored in the extract-treated rats. However, hypertensive untreated rats showed evidence of oxidative damages with significant increase in MDA, especially in the heart and liver, with decreases in the antioxidant defense system.

**Conclusion:** Results of this study justified the traditional use of *A. muricata* or *C. longa* for management of hypertension in Nigeria and showed that the extracts ameliorated oxidative damage that accompanied hypertension, thus also preventing complications of hypertension.

**Keywords:** *Annona muricata*; *Curcuma longa*; hypertension; oxidative stress markers.

## 1. INTRODUCTION

Oxidative stress, precipitated by excessive production of reactive oxygen species (ROS) which overwhelmed the antioxidant defense mechanisms, has been implicated in pathophysiological conditions that affect cardiovascular system such as hypercholesterolemia, diabetes and hypertension [1,2,3]. In animal models, oxidative stress has been demonstrated in spontaneous hypertension [4], renovascular hypertension [5], deoxycorticosterone acetate-salt model [6] and obesity-related hypertension [7]. Spontaneous hypertension in rats can be significantly decreased by reducing superoxide radicals which can be achieved by infusion of superoxide dismutase (SOD) [8].

In humans, hypertension is also considered as a state of oxidative stress that can contribute to the development of atherosclerosis [9] and other hypertension-induced organ damages [10]. Evaluation of antioxidant activities and lipid peroxidation byproducts in hypertensive subjects show an excessive amount of ROS and a decrease in the mechanism of antioxidant activity in both blood as well as in several other cellular systems [11,12], including vascular wall cells [13]. The instability of critical non-lipid macromolecules as another consequence of the overproduction of ROS may have important consequences on cellular functions. More recent management strategy for hypertension targets

alleviation of oxidative stress, thus more researches are geared towards antihypertensive drug candidates with capacity to reverse and or prevent development of oxidative stress in hypertensive patients.

A major source been explored are natural antioxidants from plants species which have protective effect against oxygen ion derived from free radicals involved in the development of many diseases such as arthritis, cardiovascular disorders, cancer and neurodegenerative diseases such as Parkinson's and Alzheimer's diseases [14]. Phytochemicals such as flavonoids, polyphenols, vitamin C and E and carotenoids as antioxidants have been reported to protect the body system against reactive oxygen species [15,16]. Various efforts are now concentrated on many herbal plant extracts because of their antioxidant effects [17].

This study is focused on two plants traditionally used for management of hypertension in Nigeria and are proven to have antioxidant properties [18,19,20,21,22]. The plants are widely grown in Nigeria and readily available. These medicinal plants; *Annona muricata* L. and *Curcuma longa* L. have a long history of use in African Traditional medicine for treatment of several ailments including diabetes and cancer [18,23,24,25,26]. These medicinal plants are well reported to be traditionally used in Nigeria for treatment of hypertension [27,28,29,30]. This study therefore seeks to evaluate treatment

outcome on oxidative stress status of hypertensive Wistar rats treated with the extracts of *Annona muricata* leaves or *Curcuma longa* rhizomes.

## 2. MATERIALS AND METHODS

### 2.1 Plant Collection and Extract Preparation

Fresh leaves of *Annona muricata* were harvested from a private orchard in Asaba, Delta State, Nigeria and rhizomes of *Curcuma longa* were purchased from Bodija Market, Ibadan, Nigeria. The plants were identified and voucher specimen deposited at Department of Botany, University of Ibadan with Voucher-Numbers UIH-22593 for *A. muricata* and UIH-22595 for *C. longa*. The rhizomes were chopped and air dried, while the leaves were air dried and pulverized. The plant materials were extracted by cold macerated in methanol (96%) for 72 hours. The extract was decanted with a fine cloth sieve and the filtrate decanted was concentrated using a rotary evaporator (BUCHI R-210, Switzerland) and the extract obtained was stored at 4°C. Fresh extract was reconstituted daily for dosing.

### 2.2 Experimental Animals

Seventy male Wistar rats weighing 140-180g were obtained from and housed at the Experimental Animal unit of the Department of Veterinary Pharmacology and Toxicology, University of Ibadan. The rats were maintained in a 12 hour light:dark environment with about 26°C temperature and 50% humidity. They were allowed free access to commercially available standard rat pellets (Ladokun Feeds Nig.Ltd) and fresh water *ad libitum*. The rats were acclimatized for two weeks before commencement of the experiment. All experiments and protocols described in the study were in accordance with the recommendation for animal care and use by University of Ibadan Animal Care and Use Research Ethics Committee (UI-ACUREC/App/11/2017/054) which follow internationally acceptable best practices for experimental animal care and use as adapted from the European Community and US guidelines.

### 2.3 Experimental Protocol

The rats were randomly divided into ten groups with 7 rats in each group. Group A rats were maintained as normal healthy rats (Normotensive

control), while hypertension was induced in groups B-J by unilateral nephrectomy and daily loading with sodium chloride (1%) for 42 days. Group B rats remained hypertensive and untreated throughout the study. Groups C and D rats were hypertensive rats treated with lisinopril or hydrochlorothiazide (standard antihypertensives) respectively. Groups E, F and G were hypertensive rats treated with *Annona muricata* leaves extract (100mg/kg, 200mg/kg or 400mg/kg), while Group H, I and J rats were hypertensive rats treated with *Curcuma longa* rhizomes extract (100mg/kg, 200mg/kg or 400mg/kg). The experimental hypertension was maintained for 42 days, alongside treatment with the antihypertensive drugs or plant extracts. Blood pressure was monitored by non-invasive method using an electro-sphygmomanometer (CODA, Kent Scientific, USA).

### 2.4 Sample Collections and Homogenate Preparation

Blood sample was collected from the retro-orbital sinus into lithium heparinized bottles on day 43. After blood collection, the rats were humanely sacrificed by cervical dislocation. The heart, liver and kidney of each rat was carefully removed, immediately perfused with normal saline and blotted with filter paper. It was homogenized in cold potassium phosphate buffer (0.1 M, pH 7.4) using a Teflon homogenizer. The homogenate was centrifuged at 10,000 rpm for 10 minutes with a cold centrifuge at 4°C to obtain post-mitochondrial fraction. An estimation of serum total protein and nitric oxide (NO) was carried out. Reduced glutathione (GSH), glutathione peroxidase (GPx), glutathione transferase (GST), superoxide dismutase (SOD) and malonaldehyde (MDA) were determination from the supernatant.

### 2.5 Data Analysis

All values were expressed as mean±S.D. The test of significance between two groups was estimated by student's t-test. One-way analysis of variance (ANOVA) with Tukey's post-hoc test was performed using Graph Pad Prism version 4.00.

## 3. RESULTS AND DISCUSSION

### 3.1 Results

#### 3.1.1 Blood pressure


The systolic, diastolic and mean arterial blood pressures of hypertensive rats were significantly

( $p < 0.05$ ) higher than the normotensive rats. These were significantly ( $p < 0.05$ ) reversed in hypertensive rats treated with the methanol extracts of *A. muricata* leaves or *C. longa* rhizomes, or the standard antihypertensive drugs (lisinopril or Chlorothiazide) (Fig. 1).


### 3.1.2 Antioxidant defense systems

The result shows that nitric oxide (NO) levels were significantly ( $p < 0.01$ ) decreased in untreated hypertensive rats, but was reversed in hypertensive rats treated with the extracts of *A. muricata* or *C. longa* or the antihypertensive drugs. NO levels in rats treated with the extracts were comparable to that in normotensive rats

(Fig. 2). Hypertension produced a significant ( $p < 0.01$ ) reduction in the heart, kidney and liver glutathione peroxidase (GPx) level in the untreated hypertensive group when compared to the normotensive rats. This was reversed in only hypertensive rats treated with *A. muricata* or *C. longa* (100mg/kg), while other hypertensive rats had reduced heart GPx levels. Depression of kidney GPx levels was reversed in all treated groups except in rats that received the higher doses of *C. longa* (200mg/kg or 400mg/kg). A reversal of the depression in GPx was also observed in liver of treated rats, with significant ( $p < 0.05$ ) elevations of liver GPx in rats treated with *A. muricata* (200mg/kg and 400mg/kg) (Fig. 3). Reduced glutathione (GSH) levels were



**Fig. 1. Blood pressure of uni-nephrectomized rats loaded daily with NaCl (1%) and treated with methanol extract of *Annona muricata* leaves or *Curcuma longa* rhizomes**  
 ### Significantly ( $p < 0.001$ ) different from normotensive control; \*Significantly different from hypertensive untreated ( $*p < 0.05$ ,  $***p < 0.0001$ )




**Fig. 2. Effects of *Annona muricata* leaves, *Curcuma longa* rhizomes, lisinopril and chlorothiazide on serum NO**  
 Significantly different from the normotensive rats (#### $p < 0.01$ ); Significantly different from hypertensive untreated rats ( $*p < 0.05$ ,  $****p < 0.0001$ )


also depressed in the heart, kidney and liver of untreated rats compared to normotensive rats, but a reversal was observed in all the treated hypertensive rats. Significant ( $p < 0.01$ ) elevations of heart and kidney GSH levels were also observed in hypertensive rats treated with methanol extract of *A. muricata* (400mg/kg) and *C. longa* (200mg/kg) (Fig. 4).

Glutathione s-transferase (GST) levels significantly ( $p < 0.05$ ) declined in hearts, kidney and liver of untreated hypertensive rats compared to normotensive rats but was also reversed in all treated rats with significant elevations in the heart of hypertensive rats treated with *A. muricata* (400mg/kg) and the liver of rats treated with *C. longa* (100mg/kg and

200mg/kg) (Fig. 5). The same pattern of decline in untreated hypertensive rats compared to normotensive rats and reversal in all treated rats was observed for superoxide dismutase (SOD) levels in the heart, kidney and liver of these rats. In addition, significant elevations in SOD levels were observed in kidneys of rats treated with *C. longa* (400mg/kg) and liver of rats treated with *A. muricata* (200mg/kg), while SOD were statistically unchanged in liver of rats treated with *A. muricata* (100mg/kg) and *C. longa* (100 and 400mg/kg) (Fig. 6). Malondialdehyde (MDA) levels were significantly ( $p < 0.0001$ ) elevated in untreated hypertensive rats compared to normotensive rats, but were remarkably reversed to normal levels in treated rats with the extracts or antihypertensives (Fig. 7).



**Fig. 3. Effects of *Annona muricata* leaves, *Curcuma longa* rhizomes, lisinopril and chlorothiazide on heart, kidney and liver glutathione peroxidase**  
 ## Significantly different from the normal control group ( $p < 0.01$ ); \*Significantly different from hypertensive untreated control ( $*p < 0.05$ ,  $***p < 0.001$ )



**Fig. 4. Effects of *Annona muricata* leaves, *Curcuma longa* rhizomes, lisinopril and chlorothiazide on heart, kidney and liver GSH**  
 ## Significantly different from the normal control group ( $p < 0.01$ ); \*Significantly different from hypertensive untreated control ( $*p < 0.05$ ,  $***p < 0.0001$ )

### 3.2 Discussion

Medicinal plants are now considered as the basis for health preservation and care worldwide [31]. Chronic degenerative diseases (hypertension, diabetes, cardiovascular and cancer) have reached epidemic proportions and are considered a serious health problem; therefore, the treatments of these diseases are of clinical importance [32]. In this study, hypertension was induced by unilateral nephrectomy and daily loading with sodium chloride (1%) which resulted


in renal hypertension [33,34]. *Annona muricata* L. (Family: Annonaceae) and *Curcuma longa* (Family: Curcubitaceae) demonstrated potent antihypertensive properties evidenced by the reversal of the elevated blood pressure, restoration of antioxidants and reduction of oxidants generated in the induced hypertensive state.

Previous studies have shown evidences of the relation between hypertension and oxidative stress. In the review by Baradaran *et al.* [35],

hypertensive patients were reported to have reduced superoxide dismutase and glutathione peroxidase levels with an inverse relationship with the blood pressure. The reports by Hirata and Satonaka [36] and Moon and Won [37] further explained the correlation between oxidative stress and hypertension as the resultant stimuli sequel to mechanical stretch of endothelial walls and activation of the renin-angiotensin system (RAS). Activation of RAS

leads to generation of reactive oxygen species which forms the pathogenesis of oxidative stress and also further exacerbates hypertension [38].

The methanol extract of *A. muricata* leaves and *C. longa* rhizomes inhibited development of hypertension shown by normal systolic blood pressure, diastolic blood pressure and mean arterial pressure of these treated hypertensive rats. This is in agreement with an earlier report in




**Fig. 5. Effects of *Annona muricata* leaves, *Curcuma longa* rhizomes, lisinopril and chlorothiazide on heart, kidney and liver GST**  
 #### Significantly different from the normal control group ( $p < 0.01$ ); \*\*\*\* Significantly different from hypertensive untreated control ( $p < 0.0001$ )





free radicals. Antioxidant defense systems of cells contain a variety of enzymatic and non-enzymatic scavengers. The enzymatic antioxidants of cells, including glutathione peroxidase (GPx), glutathione reductase (GR), glutathione-s-transferase (GST) and superoxide dismutase (SOD) play a critical role in the

attenuation of oxidative stress induced by reactive oxygen species [41]. Reduced glutathione substrate augments the activity of GPx and GST in catalyzing the hydrogen peroxide into oxygen and water. The reduced glutathione has the ability to reduce the oxidized glutathione, catalyzed by GR [42].



**Fig. 7. Effects of *Annona muricata* leaves, *Curcuma longa* rhizomes, lisinopril and chlorothiazide on heart, kidney and liver MDA**

#### Significantly different from the normal control group ( $p < 0.0001$ );  
 \*Significantly different from hypertensive untreated control ( $****p < 0.0001$ )

The first defensive mechanism against reactive oxygen species is provided by SOD, which attenuates oxidative stress through dismutation of  $O_2^-$ . Catalase enzyme has an important role in converting the endogenous  $H_2O_2$  to water and oxygen [43]. The accumulation of  $H_2O_2$  in cells results in the generation of highly reactive free hydroxyl radical ( $OH^\cdot$ ) through Fenton reaction, which has an important devastating role in oxidative damages [44]. GPx degrades lipid peroxides to hydroxyl lipids and waters through conversion of glutathione to glutathione disulfide [45,46].

A major marker of lipid peroxidation is malondialdehyde (MDA) which increases during oxidative damage to cell membranes, inhibition of several important enzymes, reduced cellular function, and cell death [47,48]. The degree of lipid peroxidation can be determined by tissue MDA levels, which is a highly reliable marker of oxidative stress [49]. MDA is a highly reactive aldehyde which can cause toxic stress in cells and result in formation of covalent protein adducts known as advanced lipoxidation end-products, an analogy of advanced glycation end-products [50]. The result of this study shows that induction of hypertension produced a significant ( $p < 0.0001$ ) elevation of MDA in the heart, kidney and liver of untreated hypertensive group when compared to the normotensive rats. Treatment with methanol extract of *A. muricata* or *C. longa*, lisinopril and chlorothiazide produced a significant ( $p < 0.0001$ ) reduction in the heart, kidney and liver MDA. This indicates remarkable inhibition of lipid peroxidation which usually accompanies and further exacerbates oxidative stress and hypertension [51].

Depletion of nitric oxide (NO) was reversed in rats treated with methanol extract of *A. muricata* and *C. longa* in this study. Nitric oxide is generated from its precursor L-arginine by nitric oxide synthase (NOS). There are three isoforms of the enzyme; the two constitutive forms, endothelial and neuronal NOS (eNOS and nNOS) and the inducible isoform originally described in immune cells (iNOS). Nitric oxide effects its principle biological actions, including that of vascular smooth muscle relaxation, via soluble guanylate cyclase and production of the second messenger c-GMP [52]. Interestingly, *A. muricata* or *C. longa* treatment exhibited a good therapeutic profile with a marked increase of serum NO level thereby enhancing the vasodilatory effects of NO with resultant lowering of blood pressure.

#### 4. CONCLUSION

In conclusion, methanol extract of *Annona muricata* and *Curcuma longa* ameliorated the oxidative stress which accompanies and exacerbates hypertension in uni-nephrectomized rats loaded with 1% sodium chloride. This study corroborated previous findings on the correlation between hypertension and oxidative stress [51,53], and improvement of renovascular hypertension following antioxidant treatment [54]. Further studies are warranted to establish the pharmacological principle responsible for the antihypertensive activity of these medicinal plants which can be progressed as antihypertensive drug candidates.

#### CONSENT

It is not applicable.

#### ETHICAL APPROVAL

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as regulations set by the Animal Care and Use in Research Ethics Committee of the University of Ibadan with approval number UI-ACUREC/App/11/2017/054.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Steinberg D, Witztum JL. Is the Oxidative Modification Hypothesis Relevant to Human Atherosclerosis? *Circul.* 2002;105: 2107–2111.
2. Chilsom GM, Steimberg D. The Oxidative Modification Hypothesis of Atherogenesis. An Overview. *Free Rad Biol Med.* 2000;28: 1815–1826.
3. Cai H, Harrison DG. Endothelial dysfunction in cardiovascular diseases: The role of oxidant stress. *Circul. Res.* 2000;87(10):840–44.
4. Wu L, Jourlink BH. Increased methylglyoxal and oxidative stress in hypertensive rat vascular smooth muscle cells. *Hypert.* 2002;39:809–814.
5. Lerman LO, Nath KA, Rodriguez-Porcel M, Krier JD, Schwartz RS, Napoli C, Romero JC. Increased oxidative stress in

- experimental renovascular hypertension. *Hypert.* 2001;37:541–546.
6. Rolliet MR, Rudd MA, Loscalzo J. Oxidative stress and renal dysfunction in salt-sensitive hypertension. *Kidney Blood Press Res.* 2001;24:116–123.
  7. Dobrian AD, Davies MJ, Schriver SD, Lauterio TJ, Prewitt RL. Oxidative stress in a rat model of obesity-induced hypertension. *Hypert.* 2001;37:554–560.
  8. Nakazono K, Watanabe N, Matsuno K, Sasaki J, Sato T, Inoue M. Does superoxide underlie the pathogenesis of hypertension? *Proc Natl Acad Sci USA.* 1991;88:10045–10048.
  9. Romero JC, Reckelhoff JF. Role of angiotensin and oxidative stress in essential hypertension. *Hypert.* 1999;34: 943–949.
  10. Raj L. Nitric Oxide in Hypertension. Relationship with Renal Injury and Left Ventricular Hypertrophy. *Hypert.* 1998;31: 189–193.
  11. McIntyre M, Bohr DF, Dominiczak AF. Endothelial function in hypertension: The Role of superoxide anion. *Hypert.* 1999; 34:539–545.
  12. Yasunari K, Maeda K, Nakamura M, Yoshikawa J. Oxidative stress in leukocytes is a possible link between blood pressure, Blood Glucose and C-Reactive Protein. *Hypert.* 2002;39:777–780.
  13. Orié NN, Zidek, W, Tepel M, Reactive Oxygen species in essential hypertension and non-insulin-dependent diabetes mellitus. *Amer J Hypert.* 1999;12:1169–1174.
  14. Almeida MMB, de Lemos LG. Bioactive compound and antioxidant activity of fresh exotic fruits from Northern eastern Brazil. *Food Res Inter.* 2011;44:2155-2159.
  15. Gutteridge JM, Halliwell B. Antioxidants: Molecules, medicines and myths. *Biochem Biophys Res Comm.* 2010;393:561-564.
  16. Mohamed DA, Hamed TE, AL-Okbi SY. Reduction in hypercholesterolemia and risk of cardiovascular diseases by mixtures of plant food. *Grasas Aceites.* 2010;61(4): 378-389.  
DOI:https://doi.org/10.3989/gya.021210
  17. Akinloye DI, Osatuyi OA, Musibau OG, Yusuf AA, Adewuyi S. *In vitro* antioxidant activities, elemental analysis and some bioactive constituents of *Acalypha wilkesiana* Muellarg and *Acalypha wilkesiana* Java white leaf extracts. *J Chem Soc Nigeria.* 2016;41(2):150-157.
  18. Saba AB, Oridupa OA. Search for a novel antioxidant, anti-inflammatory/analgesic or anti-proliferative drug: cucurbitacins hold the ace. *J Med Plants Res.* 2010;4(25): 2821-2826.
  19. George VC, Kumar DRN, Suresh PK, Kumar RA. Antioxidant, DNA protective efficacy and HPLC analysis of *Annona muricata* (soursop) extracts. *J Food Sci Technol.* 2014;52(4):2328–2335.  
DOI:https://doi.org/10.1007/s13197-014-1289-7
  20. Muthu S, Durairaj B. Evaluation of antioxidant and free radical scavenging activity of *Annona muricata*. *Eur J Exper Biol.* 2015;5(3):39-45.
  21. Agu KC, Okolie PN. Proximate composition, phytochemical analysis, and *in vitro* antioxidant potentials of extracts of *Annona muricata* (Soursop). *Food Sci Nutr.* 2017;5(5):1029–1036.  
DOI:https://doi.org/10.1002/fsn3.498
  22. Tanvir EM, Hossen MS, Hossain MF, Afroz R, Gan SH, Khalil MI, Karim N. Antioxidant Properties of Popular Turmeric (*Curcuma longa*) Varieties from Bangladesh. *Journal of Food Quality Volume 2017*, Article ID 8471785, 8 pages.  
DOI:https://doi.org/10.1155/2017/8471785.
  23. Adewole SO, Caxton-Martins EA. Morphological changes and hypoglycemic effects of *Annona muricata* Linn. (*Annonaceae*) leaf aqueous extract on pancreatic B-cells of streptozotocin-treated diabetic rats. *Afr J Biomed Res.* 2006;9: 173–187.  
DOI:https://doi.org/10.4314/ajbr.v9i3.48903
  24. Adewole S, Ojewole J. Protective effects of *Annona muricata* linn. (*annonaceae*) leaf aqueous extract on serum lipid profiles and oxidative stress in hepatocytes of streptozotocin-treated diabetic rats. *Afr J Tradit Complement Altern Med.* 2009;6: 30–41.  
DOI:https://doi.org/10.4314/ajtcam.v6i1.57071
  25. Mishra S, Ahmad S, Kumar N, Sharma BK. *Annona muricata* (the cancer killer): A review. *Glob. J Pharm Res.* 2013;2:1613–1618.
  26. Priyadarsini KI. Chemical and structural features influencing the biological activity of curcumin. *Curr Pharm Des.* 2013;19: 2093–2100.
  27. Nwokocha CR, Owu DU, Gordon A, Thaxter K, McCalla G, Ozolua RI, Young L. Possible mechanisms of action of the

- hypotensive effect of *Annona muricata* (soursop) in normotensive Sprague-Dawley rats. *Pharm Biol.* 2012;50(11): 1436–1441.
28. Adefegha SA, Oyeleye SI, Oboh G. Distribution of Phenolic Contents, Antidiabetic Potentials, Antihypertensive Properties, and Antioxidative Effects of Soursop (*Annona muricata* L.) Fruit Parts In vitro. *Biochem Res Int.* 2015: 347673. DOI:10.1155/2015/347673.
  29. Akinyemi AJ, Adedara IA, Thome GR, Morsch VM, Rovani MT, Mujica LKS, Duarte T, Duarte M, Oboh G, Schetinger MRC. Dietary supplementation of ginger and turmeric improves reproductive function in hypertensive male rats. *Toxicol Rep.* 2015;2:1357-1366. DOI:10.1016/j.toxrep.2015.10.001.
  30. Oyemitan IA, Elusiyana CA, Onifade AO, Akanmu MA, Oyedele AO, McDonald AG. Neuropharmacological profile and chemical analysis of fresh rhizome essential oil of *Curcuma longa* (turmeric) cultivated in Southwest Nigeria. *Toxicol Rep.* 2017;4:391–398. DOI:10.1016/j.toxrep.2017.07.001
  31. Pan SY, Zhou SF, Gao SH, Yu ZL, Zhang SF, Tang MK, Sun JN, Ma DL, Han YF, Fong WF, Ko KM. New Perspectives on How to Discover Drugs from Herbal Medicines: CAM's Outstanding Contribution to Modern Therapeutics. *Evid. Based Complement Alternat Med.* 2013; 627375. DOI:<https://doi.org/10.1155/2013/627375>
  32. World Health Organization (WHO). Preventing chronic diseases a vital investment; 2005. Available:[http://www.who.int/chp/chronic\\_disease\\_report/full\\_report.pdf](http://www.who.int/chp/chronic_disease_report/full_report.pdf) (accessed on 25.08.15)
  33. Mozaffari MS, Wyss JM. Dietary NaCl-induced hypertension in uninephrectomized wistar-kyoto rats: role of kidney function. *J Cardiovasc Pharmacol.* 2005;33(5):814-821.
  34. Lin HY, Lee YT, Chan YW, Tse G. Animal models for the study of primary and secondary hypertension in humans. *Biomed Rep.* 2016;5(6):653–659. DOI:<https://doi.org/10.3892/br.2016.784>
  35. Baradaran A, Nasri H, Rafieian-Kopaei M. Oxidative stress and hypertension: Possibility of hypertension therapy with antioxidants. *J Res Med Sci.* 2014;19(4): 358–367.
  36. Hirata Y, Satonaka H. Hypertension and oxidative stress. *JMAJ.* 2001;44(12):540-545.
  37. Moon JS, Won KC. Oxidative stress: link between hypertension and diabetes. *Korean J Intern Med.* 2017;32(3):439–441. DOI:10.3904/kjim.2017.153
  38. Wu J, Harrison DG. Oxidative stress and hypertension. In: Blood pressure and arterial wall mechanics in cardiovascular diseases. Michael E. Safar, Michael F. O'Rourke and Edward D. Frohlich (Editors). Publisher Springer. 2015;175-193.
  39. Nakmareong S, Kukongviriyapan U, Pakdeechote P, Donpunha W, Kukongviriyapan V, Kongyingoes B, Sompamit K, Phisalaphong C. Antioxidant and vascular protective effects of curcumin and tetrahydrocurcumin in rats with L-NAME-induced hypertension. *Naunyn Schmiedebergs Arch Pharmacol.* 2011; 383:519–529.
  40. Yao Y, Wang W, Li M, Ren H, Chen C, Wang J, Wang WE, Yang J, Zeng C. Curcumin exerts its antihypertensive effect by down regulating the AT1 receptor in vascular smooth muscle cells. *Sci Reports.* 2016;6:25579. DOI:<https://doi.org/10.1038/srep25579>
  41. Singh S, Vrishni S, Singh BK, Rahman I, Kakkar P. Nrf2-ARE stress response mechanism: A control point in oxidative stress-mediated dysfunctions and chronic inflammatory diseases. *Free Rad Res.* 2010;44:1267-1288.
  42. Bazzichi L, Ciompi ML, Betti L, Rossi A, Melchiorre D, Fiorini M, Giannaccini G, Lucacchini A. Impaired glutathione reductase activity and levels of collagenase and elastase in synovial fluid in rheumatoid arthritis. *Clin Exp Rheumat.* 2002;20:761-766.
  43. Fahmy HM, Mohammed FF, Abdelrahman RT, Abu Elfetoh MM, Mohammed YA. Effect of Radiofrequency Waves Emitted from Conventional WIFI Devices on Some Oxidative Stress Parameters in Rat Kidney. *J. Drug Metab. Toxicol.* 2015;6: 195. DOI:<https://doi.org/10.4172/2157-7609.1000195>
  44. Park S, Imlay JA. High levels of intracellular cysteine promote oxidative DNA damage by driving the fenton reaction. *J Bacteriol.* 2003;185:1942-1950. DOI:<https://doi.org/10.1128/JB.185.6.1942>

45. Naziroglu M, Kokcam I. Antioxidants and lipid peroxidation status in the blood of patients with alopecia. *Cell Biochem Funct.* 2008;18:169–173.
46. Blokhina O, Virolainen E, Fagerstedt KV. Antioxidants, oxidative damage and oxygen deprivation stress: A review. *Ann Bot.* 2003;91(2):179–194. DOI:<https://doi.org/10.1093/aob/mcf118>
47. Abdel Fattah NS, Ebrahim AA, El Okda ES. Lipid peroxidation/antioxidant activity in patients with *Alopecia areata*. *J Eur Acad Dermatol Venereol.* 2011;25(4):403-8. DOI:<https://doi.org/10.1111/j.1468-3083.2010.03799.x>.
48. Prie BE, Voiculescu VM, Ionescu-Bozdog OB, Petrutescu B, Iosif L, Gaman LE, Clatici VG, Stoian I, Giurcaneanu C. Oxidative stress and *Alopecia areata*. *J Med Life.* 2015;8(Spec Issue):43–46.
49. Davey MW, Stals E, Panis B, Keulemans J, Swennen RL. High-throughput determination of malondialdehyde in plant tissues. *Analyt Biochem.* 2005;347(2): 201–207. DOI:10.1016/j.ab.2005.09.041.
50. Farmer EE, Davoine C. Reactive electrophile species. *Curr Opin Plant Biol.* 2007;10(4):380–6. DOI:<https://doi.org/10.1016/j.pbi.2007.04.019>.
51. Touyz RM, Briones AM. Reactive oxygen species and vascular biology: Implications in human hypertension. *Hypert Res.* 2011;34:5–14.
52. Francis SH, Busch JL, Corbin JD. cGMP-Dependent protein Kinases and cGMP phosphodiesterases in nitric oxide and cGMP action. *Pharmacol Rev.* 2010; 62(3):525–563. DOI:<https://doi.org/10.1124/pr.110.002907>.
53. Wang H, Li H, Hou Z, Pan L, Shen X, Li G. Role of oxidative stress in elevated blood pressure induced by high free fatty acids. *Hypert Res.* 2009;32:152–158.
54. Nishi EE, Oliveira-Sales EB, Bergamaschi CT, Oliveira, TGC, Boim MA, Campos RR. Chronic antioxidant treatment improves arterial renovascular hypertension and oxidative stress markers in the kidney in Wistar rats. *Am J Hypertens.* 2010;23: 473-480. DOI:<https://doi.org/10.1038/ajh.2010.1>.

© 2018 Oridupa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/47551>