



Assessment of Different Germplasm of *Aglaonema* for Pot and Removal of Particulate Matter in Air

Yukthamukhi R ^{a*}, Alka Singh ^b, Nikhitha ^c, Anusha S R ^d,
HP Shah ^b and A. P. Chaudhary ^e

^a ASPEE College of Horticulture, NAU, Navsari, (Gujarat)-396450, India.

^b Department of Floriculture and Landscape Architecture, NAU, Navsari, (Gujarat)-396450, India.

^c Department of Post Harvest Management, College of Horticulture, Bengaluru, Karnataka-560065, India.

^d Department of Vegetable science, College of Horticulture, Bengaluru, Karnataka-560065, India.

^e Department of Social Science, ASPEE college of Horticulture, NAU, Navsari (Gujarat)-396450, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijecc/2024/v14i104501>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/123659>

Original Research Article

Received: 19/07/2024

Accepted: 21/09/2024

Published: 04/10/2024

ABSTRACT

The aim of the experiment is to screen twelve germplasm of *Aglaonema* for their suitability as pot plants and removal of particulate matter in air. The experiment was designed using Completely Randomized Design (CRD) at the greenhouse complex of the Department of Floriculture and Landscape Architecture, ASPEE College of Horticulture, Navsari (Gujarat), during 2023-24. Morphological parameters were measured using various methods, including a measuring tape for

*Corresponding author: E-mail: yuktharukma@gmail.com; yuktha251@gmail.com;

Cite as: R, Yukthamukhi, Alka Singh, Nikhitha, Anusha S R, HP Shah, and A. P. Chaudhary. 2024. "Assessment of Different Germplasm of *Aglaonema* for Pot and Removal of Particulate Matter in Air". *International Journal of Environment and Climate Change* 14 (10):490-95. <https://doi.org/10.9734/ijecc/2024/v14i104501>.

plant height, a digital leaf area meter for leaf area, and manual counting for leaves and branches. Leaf color was evaluated on visual grading system and the RHS Colour Chart (2015), to assess visual appeal and characteristics and percent removal of particulate matter (PM) by the plant was measured using digital air quality index meter. *Aglaonema* 'Emerald Bay' exhibited maximum plant height (73.18 cm), leaf area (311.66 cm²), leaf length (39.28 cm), leaf width (15.26 cm), internodal length (3.43 cm) and visual leaf colour grade (4.88) and dark green leaf colour (137A) as per RHS colour chart followed by *Aglaonema commutatum* with the plant height (69.58 cm), leaf area (298.09cm²), leaf length (34.28 cm), leaf width (11.10 cm), internodal length (3.34 cm) and visual leaf colour grade (4.80) and dark green leaf colour (139 A) . Maximum number of leaves were recorded in *Aglaonema* 'Cutlass' (43.90), while maximum number of branches per plant were observed in *Aglaonema brevispathum* (13.56). *Aglaonema* 'Emerald Bay' showed the maximum removal rate of PM₁ per 24 hours (92.16%) followed by *Aglaonema commutatum* (85.97%). Among all the germplasms, *Aglaonema* 'Emerald Bay' and *Aglaonema commutatum* have been highly suitable as pot plant owing to its good plant height, leaf area and their efficiency in removal of PM₁.

Keywords: *Aglaonema*; pot plant; morphological; visual leaf colour; particulate matter.

ABBREVIATION

Particulate matter 1 (PM₁): A type of air pollution that refers to particles that are less than 1 micron in diameter.

1. INTRODUCTION

Indoor gardening has been gaining high impetus with enhanced sense of emotional comfort, air quality and stress-free happy environment during the post COVID era [1]. Further, pot plant trading is growing with rapid urbanization and hasty life style [2,3,4].

Aglaonema, commonly known as Chinese evergreens, is a genus of 21 species within the Araceae family, native to Southeast Asia, Northeast India, and Southern China, with a distribution extending to Malaysia, New Guinea, and the Philippines [5,6]. This genus has been extensively studied for its adaptability to indoor environments and its potential uses [7,8]. It is known for its striking foliage and adaptability, is a widely cultivated genus of ornamental plants, particularly valued as pot plants, it encompasses a diverse range of species and cultivars, each exhibiting unique morphological traits that influence their suitability for pot cultivation. *Aglaonema* is ideal for indoor environments due to its air-purifying capabilities, adaptability to low light, aesthetic diversity, low maintenance, making it both functional and visually appealing [9,10]. It is essential to evaluate and screen various germplasms for desirable characteristics as pot plant and its ability to curb indoor pollution.

Aglaonema's contribution to indoor air quality is supported by its ability in removal of particulate matter and other pollutants, that makes it a valuable addition to indoor environments, promoting healthier and cleaner air [11]. However, there are different species and varieties of *Aglaonema* that needs to be studied for air quality purification.

Hence this study focused on the screening of twelve *Aglaonema* germplasms, namely *Aglaonema siam aurora* (Lipstick), *Aglaonema* 'Emerald Bay', *Aglaonema commutatum*, *Aglaonema alumina*, *Aglaonema* 'Key Lime', *Aglaonema* cutlass, *Aglaonema* 'White Rain', *Aglaonema brevispathum*, *Aglaonema* lime narrow, *Aglaonema commutatum* (variegated), *Aglaonema* white Chinese evergreen (Golden Bay), and *Aglaonema* Red Valentine. The evaluation is based on key morphological parameters and its ability to curb indoor pollution.

2. MATERIALS AND METHODS

The experiment was conducted during 2023-24 at the greenhouse complex of Department of Floriculture and Landscape Architecture, ASPEE College of Horticulture, Navsari (Gujarat), using a Completely Randomized Design (CRD) to screen twelve *Aglaonema* germplasms for their suitability as pot plants and removal of particulate matter in air. The plants were maintained under uniform growing conditions to ensure consistency in environmental factors such as light, water, and nutrients. Morphological parameters were recorded both at the start of the experiment and after two months. plant height was measured

from the soil level to the growing tip using a measuring tape. Leaf area was determined using a digital leaf area meter, while leaf length and width were measured using a meter scale. The number of leaves and branches per plant were counted manually. The visual colour grade of *Aglaonema* plants was assessed using a standardized grading system [12]. This system evaluates leaf colour grade based on greenness with freshness and luster, assigning scores 1= poor colour, 3= good, light green and 5= excellent dark green & silver contrast, and visual leaf color was evaluated using the RHS Colour Chart (2015). Internodal length was measured between nodes using a meter scale. Particulate matter (PM₁) level in air was recorded by using enclosed chamber using digital air quality index meter [13] before and after 24 hours and the PM₁ removal rate by the plant was calculated on per cent basis. The data on all the parameters was collected twice at an interval of two months were pooled and statistically analyzed as per the method given by Panse and Sukhatme [14] and the results were evaluated at 5 %level of significance.

3. RESULTS AND DISCUSSION

Observations on morphological parameters viz., plant height (cm), leaf area (cm²), leaf length (cm), leaf width(cm) and internodal length(cm) of twelve germplasm of *Aglaonema* are represented in the Table 1. Maximum plant height exhibited by *Aglaonema* 'Emerald Bay'(73.18 cm), closely followed by *Aglaonema commutatum* (69.58 cm). In contrast, *Aglaonema siam aurora* (lipstick) displayed the minimum plant height (30.68 cm). Maximum leaf area exhibited by *Aglaonema* 'Emerald Bay'(311.66 cm²), closely followed by *Aglaonema commutatum* (298.09 cm²). In contrast, *Aglaonema siam aurora* (lipstick) displayed the smallest leaf area (111.55 cm²). *Aglaonema* 'Emerald Bay' showed the maximum leaf length (39.28cm) followed by *Aglaonema commutatum* (34.28 cm) while it was minimum in *Aglaonema siam aurora* (lipstick) (15.25cm). *Aglaonema* 'Emerald Bay' exhibited maximum leaf width (15.26 cm), closely followed by *Aglaonema* 'white Rain' (13.30 cm), whereas *Aglaonema* 'cutlass' displayed the narrowest leaves (4.83 cm). Maximum internodal length was recorded in *Aglaonema* 'Emerald Bay' (3.43 cm), closely followed by *Aglaonema commutatum* (3.34 cm). In contrast, *Aglaonema* 'lime narrow'

exhibited the shortest internodal length (1.52 cm).

Observations on number of leaves, number of branches per plant, visual leaf color grade of twelve germplasm of *Aglaonema* are represented in the Table 2. Maximum leaf count was observed in *Aglaonema* 'Cutlass' (43.90), followed by *Aglaonema* 'Alumina' (25.08), while *Aglaonema* 'White Rain' had the minimum (11.16). *Aglaonema brevispathum* exhibited maximum number of branches per plant, with an average of 13.56, closely followed by *Aglaonema* 'Cutlass' (12.24 branches per plant). In contrast, *Aglaonema siam aurora*' (Lipstick) displayed the lowest branching frequency, with an average of 4.20 branches per plant. *Aglaonema* 'Emerald Bay' exhibited the highest visual leaf color grade (4.88), closely followed by *Aglaonema* 'Alumina' (4.80) on the basis of 5 point visual grade for freshness, greenness and luster. Variation with regard to colour as per RHS chart among different germplasms was observed, wherein *Aglaonema* 'Emerald Bay', *Aglaonema commutatum* and *Aglaonema alumina* exhibited dark green colour, *Aglaonema siam aurora* (Lipstick) exhibited dark purple red, *Aglaonema* 'Key Lime' and *Aglaonema* lime narrow exhibited medium yellow green, *Aglaonema* cutlass exhibited medium green, *Aglaonema* 'White Rain', *Aglaonema commutatum* (variegated) and *Aglaonema* white Chinese evergreen (Golden Bay) exhibited light green and *Aglaonema* Red Valentine exhibited light red pink colour (Fig. 1).

Observations on removal rate of Particulate Matter₁ per 24 h (PM₁) (%) is represented in the Table 3. *Aglaonema* 'Emerald Bay' showed the maximum removal rate of PM₁ (%) in 24 hours (92.16%) followed by *Aglaonema commutatum* (85.97%) while it was minimum in *Aglaonema* Red Valentine (63.06%). The results suggest that the morphological characteristics of plants play a crucial role in reducing PM concentration. Leaf traits, including shape, arrangement, and the leaf area and hair structure on the surface, are particularly important for PM accumulation. Plants remove PM through stomatal uptake and deposition on the leaf surface. PM particles can penetrate the stomata and accumulate beneath the epidermis or in the palisade layer of leaves [15,16,17,18,19,20]. Higher leaf area in *Aglaonema* 'Emerald Bay' and *Aglaonema commutatum* contributed to higher PM removal rate.

Table 1. Morphological characters of twelve germplasm of *Aglaonema*

Treatments	Morphological characters				
	Pooled data of two months				
	Plant Height (cm)	Leaf area (cm ²)	Leaf length (cm)	Leaf width (cm)	Internodal length (cm)
G ₁	30.68	111.55	15.25	6.28	1.92
G ₂	73.18	311.66	39.28	15.26	3.43
G ₃	69.58	298.09	34.28	11.10	3.34
G ₄	40.24	212.43	21.71	10.53	2.91
G ₅	54.74	212.26	23.22	11.69	3.08
G ₆	45.07	113.70	27.82	4.83	2.23
G ₇	62.75	236.36	33.94	13.30	2.06
G ₈	56.48	271.19	29.14	12.57	3.00
G ₉	56.15	269.81	25.40	7.57	1.52
G ₁₀	66.49	154.72	26.09	8.30	2.25
G ₁₁	35.42	252.93	25.58	12.21	2.80
G ₁₂	37.05	119.65	20.61	6.35	1.81
S.Em ±	0.39	1.76	0.45	0.15	0.04
CD at 5%	1.13	5.03	1.30	0.38	0.10

Table 2. Morphological characters of twelve germplasm of *Aglaonema*

Treatments	Morphological characters				
	Pooled data of two months				
	Number of leaves	Number of branches per plant	visual leaf colour grade (5-point scale)	RHS Colour group	colour code
G ₁	13.43	4.20	4.21	Dark purple red	46 A
G ₂	14.16	4.89	4.88	Dark green	137 A
G ₃	11.18	8.36	4.80	Dark green	139 A
G ₄	25.08	9.35	4.79	Dark green	131B
G ₅	19.78	6.58	4.34	Medium yellow green	151 D
G ₆	43.90	12.24	4.61	Medium green	144 A
G ₇	11.16	10.83	4.30	Light green	141D
G ₈	13.63	13.56	4.71	Medium grey green	191 C
G ₉	12.86	11.56	4.71	Medium yellow green	154 D
G ₁₀	15.85	9.11	4.68	Light green	144 B
G ₁₁	11.79	5.11	4.69	Light green	145 D
G ₁₂	12.93	5.00	4.06	Light red pink	49 D
S.Em ±	0.34	0.13	0.03		
CD at 5%	0.98	0.36	0.10		

Table 1 and 2. G₁: *Aglaonema siam aurora* (Lipstick), G₂: *Aglaonema 'Emerald Bay'*, G₃: *Aglaonema commutatum*, G₄: *Aglaonema alumina*, G₅: *Aglaonema 'Key Lime'*, G₆: *Aglaonema cutlass*, G₇: *Aglaonema 'White Rain'*, G₈: *Aglaonema brevispathum*, G₉: *Aglaonema lime narrow*, G₁₀: *Aglaonema commutatum (variegated)*, G₁₁: *Aglaonema white Chinese evergreen (Golden Bay)*, G₁₂: *Aglaonema Red Valentine*

Table 3. Effect of different germplasms of *Aglaonema* on Removal rate of PM₁ (%) in 24 h

Treatments	Removal rate of PM ₁ (%) / 24 h	Mean (T)
G ₁ : <i>Aglaonema siam aurora</i> (Lipstick)		64.96
G ₂ : <i>Aglaonema</i> 'Emerald Bay'		92.16
G ₃ : <i>Aglaonema commutatum</i>		85.97
G ₄ : <i>Aglaonema alumina</i>		80.05
G ₅ : <i>Aglaonema</i> 'Key Lime'		81.09
G ₆ : <i>Aglaonema</i> cutlass		70.03
G ₇ : <i>Aglaonema</i> 'White Rain'		85.11
G ₈ : <i>Aglaonema brevispathum</i>		81.13
G ₉ : <i>Aglaonema</i> lime narrow		81.15
G ₁₀ : <i>Aglaonema commutatum</i> (variegated)		72.15
G ₁₁ : <i>Aglaonema</i> white Chinese evergreen (Golden Bay)		72.15
G ₁₂ : <i>Aglaonema</i> Red Valentine		63.06
S.Em ±		0.12
CD at 5%		0.36




Fig. 1. Effect of different germplasms of *Aglaonema* on percent removal rate of PM₁ (%) / 24h

G₁: *Aglaonema siam aurora* (Lipstick), G₂: *Aglaonema* 'Emerald Bay', G₃: *Aglaonema commutatum*, G₄: *Aglaonema alumina*, G₅: *Aglaonema* 'Key Lime', G₆: *Aglaonema* cutlass, G₇: *Aglaonema* 'White Rain', G₈: *Aglaonema brevispathum*, G₉: *Aglaonema* lime narrow, G₁₀: *Aglaonema commutatum* (variegated), G₁₁: *Aglaonema* white Chinese evergreen (Golden Bay), G₁₂: *Aglaonema* Red Valentine

4. CONCLUSION

Based on the findings of the present investigation, *Aglaonema* 'Emerald Bay' and *Aglaonema commutatum* exhibited superior morphological characteristics with maximum values for plant height, leaf area, leaf length, leaf width, visual leaf colour grade with dark green and lustrous leaf colour and along with efficient removal rate of particulate matter and thus, making them the more suitable germplasms for pot plants. Overall, *Aglaonema* 'Emerald Bay' and *Aglaonema commutatum* are promising germplasms for pot plant production and indoor air purification.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models

(ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENTS

We acknowledge ASPEE college of Horticulture (NAU, Navsari, Gujrat) for providing the best platform and necessary infrastructure and support in conduction of this research endeavor.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Singh A. Indoor Gardening for Clean Air. Nursery Today. 2020, July-October;39-42.

2. Singh A, Ahlawat TR, Chavan S, Patel AI. Wonder World of adeniums. Floriculture Today. 2016;20(9):44-49.
3. Guddad M, Singh A, Shah HP, Chaudhari P and Ahlawat TR. Effect of Foliar Application of Chemicals on Plant Architecture in Potted *Ixora chinensis* var. 'Mini Double'. Curr J Appl Sci Tech. 2022; 41(32):9–15.
4. Sindhuja M, Singh A, Bhandari AJ, Shah HP, Patel Aiand, Parekh VB. Morphological characterization of different genotypes of adenium. Indian J Hort. 2022;79(3):296-304.
5. Nicolson DH. A revision of the genus *Aglaonema* (araceae). Smithsonian Contributions to Botany. 1969;1-66.
6. Govaerts R, Frodin DG. World checklist and bibliography of Araceae (and Acoraceae). Royal Botanic Gardens, Kew. 2002;1–560.
7. Henny RJ, Chen J, Mellich TA, Brennan MS. 'Moonlight Bay' *Aglaonema*. HortScience. 2008(5);43:1598–1599.
8. Li DM, Zhu GF, Yu B, Huang D. Comparative chloroplast genomes and phylogenetic relationships of *Aglaonema modestum* and five variegated cultivars of *Aglaonema*. Plos One. 2022;17(9): e0274067.
9. Chen J, Li Q, Dai X. Effect of different growing media on the growth of *Aglaonema*. Acta Hort. 2002;572:265–270.
10. Wolverson BC, Wolverson JD. Plants and soil microorganisms: Removal of formaldehyde, xylene, and ammonia from the indoor environment. J Missi Acad Sci. 1993;38(2):11–15.
11. Kim KJ, Kil MJ, Song JS, Yoo EH, Son KC, Kays SJ. Efficiency of volatile formaldehyde removal by indoor plants: Contribution of aerial plant parts versus the root zone. HortScience. 2010;45(10): 1489–1495.
12. Henny RJ, Poole RT, Conover CA. Visible leaf color grade. Hort Science. 1983; 18(3):374.
13. Yoo MH, Kwon YJ, Son KC, Kays SJ. Efficacy of indoor plants for the removal of single and mixed volatile organic pollutants and physiological effects of the volatiles on the plants. JASHS. 2006;131(4):452–458.
14. Panse VG, Sukhatme PV. Statistical methods for agricultural workers. 4th ed. New Delhi: Indian Council of Agricultural Research; 1985.
15. Chen H, Wang Z, Li X. Investigation of photo regulation in *Hedera helix* for improving indoor air quality. J Environ Manage. 2017;203:981–988.
16. Chen Q, Huang S, Shen L, Chen X. Mitigating indoor air pollutants using plant-based methods: A phytoremediation approach. Air Qual Atmos Health. 2017; 10(4):473–485.
17. Chen W, Liu Y, Tong D. Photoregulation of potted *Hedera helix* to remove volatile formaldehyde for improving indoor air quality. Build Environ. 2017;177:245–253.
18. Jang J, Park S, Lee J, Kim J. Effect of different houseplants on indoor air quality improvement. Environ Res. 2021;194: 110627.
19. Ryu Y, Jeon H, Lee J, Kim S. Evaluation of indoor air quality improvement by different types of houseplants. Indoor Built Environ. 2013;22(2):284–295.
20. Han J.A study on the effectiveness of various indoor plants in improving indoor air quality. J Environ Manage. 2019; 240:285–293.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/123659>