



# **Performance of Wheat (*Triticum aestivum* L.) Genotypes, their Growth and Yield under Irrigated Condition of Prayagraj**

**S. Prashanth<sup>a\*</sup>, C. Umesha<sup>a#</sup> and S. P. Vikas<sup>a</sup>**

<sup>a</sup> Department of Agronomy, Naini Agricultural Institute, SHUATS, Prayagraj, Uttar Pradesh, 211007, India.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/IJECC/2022/v12i1130936

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88739>

**Original Research Article**

**Received 18 April 2022**

**Accepted 27 June 2022**

**Published 01 July 2022**

## **ABSTRACT**

A field experiment was carried out during *Rabi* season, 2021 at Wheat Breeding experimental Field, Naini Agricultural institute, SHUATS, Prayagraj (U.P). The experiment was laid out in Randomized Block Design and nine genotypes were replicated fourfold. Study revealed that the genotype G<sub>2</sub> i.e., NETS-102 recorded significantly higher plant height (112.02 cm), number of tillers/hill (7.7/hill), plant dry weight (31.41 g/hill), length of the spike (12.79 cm), number of grains per spike (63.41/spike), grain yield (4.26 t/ha) and Straw yield (6.04 t/ha). It was evident that the performance of genotype NETS-102 was proved to be viable and productive.

**Keywords:** Genotype; viable; productive; performance.

## **1. INTRODUCTION**

Wheat is considered as one of the major cereal crops which belongs to the grass family [1]. Wheat is the direct source of food for human

beings and hence known as king of cereals. After rice and maize, wheat is the third most important crop under cultivation. But in terms of human consumption it ranks second. Wheat is the staple food in more than 40 countries of the world [2].

<sup>o</sup> M.Sc. Scholar,

<sup>#</sup>Assistant Professor,

\*Corresponding author: E-mail: prashanthshivarathri69@gmail.com;

Wheat is highly a diversified crop which is grown in different agro-ecological zones and environments with different production potentials. The three main species of wheat viz., *Triticum aestivum*, *Triticum durum* and *Triticum dicocum* L. are being cultivated in India.

Wheat is produced under irrigated conditions within the country, but low rainfall and late heat stress conditions are the constraints to achieve the specified results. Proper irrigation scheduling is crucial for the efficient use of water, energy and other production inputs. Irrigation water should be applied in numerous critical stages of wheat for successful wheat production. Availability of adequate amount of moisture at critical stages of plant growth not only optimizes the metabolism in plant cell but also increases the effectiveness of the mineral nutrients applied to the crop. Water requirements may vary depending on the stages of development.

Genotype plays an important role in producing high yield varieties of wheat. Different varieties responds differently for their genotypic characters, input requirement, growth process and the diversified environmental during growing season. Quality traits of the wheat also should be considered by breeders to improve high nutritional genotypes. These emerging varieties should be more productive with the limited inputs and should maintain the economical trend for the farmers.

Therefore, these promising genotypes were developed under prayagraj condition and the present investigation was carried out with the objective "Performance of Wheat (*Triticum aestivum* L.) Genotypes, their Growth and Yield under Irrigated Condition of Prayagraj".

## 2. MATERIALS AND METHODS

### 2.1 Site Selection

The experiment entitled "Performance of Wheat (*Triticum aestivum* L.) Genotypes, their Growth and Yield under Irrigated Condition of Prayagraj" was carried out during *Rabi* season of 2021 at Wheat Breeding Experimental Field, Naini Agricultural Institute, Sam Higginbottom University of Agriculture and Sciences, Prayagraj (U.P.) which is located at 25° 24' 33" N latitude, 81° 51' 12" E longitude (Google, 2022) and 96 m altitude above the mean sea level. All the facilities required for the crop cultivation were available.

### 2.2 Experimental Design

The experiment was conducted in Randomized block design consisting of nine genotypes replicated four times each and was allocated randomly in each replication. The soil of the experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 6.7). The wheat was sown on 20<sup>th</sup> November 2021 with plant geometry of 20 x 10 cm. The nutrient application was given @ 150:60:40 kg/ha (N:P:K). Pre-harvest observations were recorded from the tagged hills with time intervals of 25, 50, 75, 100 DAS and at harvest. Five plants per plot were uprooted randomly at each interval. The samples were oven dried for 72 h at 70-80°C and weights were taken with weighing balance. From the dry weight of different plants, crop growth rate and relative growth rate were calculated by classical techniques of growth analysis. And at the time of harvest only post-harvest observations were recorded. The genotypes were G<sub>1</sub> - NETS-101, G<sub>2</sub> - NETS-102, G<sub>3</sub> - NETS-103, G<sub>4</sub> - NETS-104, G<sub>5</sub> - NETS-105, G<sub>6</sub> - NETS-106, G<sub>7</sub> - NETS-107, G<sub>8</sub> - NETS-108, G<sub>9</sub> - NETS-109 respectively.

### 2.3 Statistical Analysis

The experimental data recorded was subjected to statistical analysis by adopting the Fishers method of analysis of variance (ANOVA) as described by Gomez and Gomez [3]. The data collected from the experiment was subjected to statistical analysis using ICAR WASP software. Critical difference (CD) and standard error of mean (SEm) values were calculated by the 'F' test was found significantly at 5% level.

## 3. RESULTS AND DISCUSSION

### 3.1 Growth Parameters

#### 3.1.1 Plant height (cm)

Data pertaining the plant height (cm) of wheat are presented in Table 1. Significantly higher plant height (112.02 cm) was recorded in the genotype G<sub>2</sub>. There was a significant increase in the growth in all the growth stages of the wheat genotype. However, the genotypes G<sub>9</sub> (111.5 cm), G<sub>1</sub> (108.89 cm) and G<sub>7</sub> (108.55 cm) were at par with the genotype G<sub>2</sub>. The higher plant height in the G<sub>2</sub> genotype was may be due to the maximum leaf area index, chlorophyll content and etc., The plant height is mostly associated with the genetic makeup of the parental material of each genotype. Each genotype has its own

feature from the growth viewpoint and variation in plant height was recorded as their genetic character. Similar findings were reported by Bhutto et al. [4].

### 3.1.2 Number of Tillers per hill

Genotype G<sub>2</sub> was recorded with higher number of tillers per hill (7.7) over all other genotypes and was tabulated in Table 1. However, the genotype G<sub>9</sub> (7.25) was at par with the genotype G<sub>2</sub>. Tillers ultimately affect the yield indirectly. In most of the cereals the yield is determined by the number of tillers. Maximum number of tillers was recorded at 75 DAS interval when compared to 100 DAS and at harvest. This is due to the perishable nature of the tillers after the vegetative phase (mainly non-productive tillers). These findings were found to be consistent with those of Ghanbari and Malidarreh [5].

### 3.1.3 Plant dry weight (g/hill)

Data on plant dry weight (g/hill) was presented in the Table 1. Significantly higher plant dry weight (31.41 g/hill) was recorded in the genotype G<sub>2</sub>. However, the genotype G<sub>9</sub> with the plant dry weight (29.86 g/hill) maintained to stay at par with the genotype G<sub>2</sub>. Initial growth stages exhibited low dry weight when compared with the later stages. Later on, the wheat crop gradually achieved the ultimate plant dry weight. This is due to the accumulation of dry matter and the further growth of the plant. These results are in agreement with the findings of Singh et al. [6].

### 3.1.4 Crop growth rate (g/m<sup>2</sup>/day)

Data regarding the crop growth rate is presented in the Table 1. The results pertaining to the study of crop growth rate indicated significant difference among the genotypes at 25, 50, 75, 100 DAS and at harvest. The genotype G<sub>2</sub> during 25 - 50 DAS interval dominated all the other genotypes with maximum crop growth rate (8.58 g/m<sup>2</sup>/day). The genotypes G<sub>9</sub>, G<sub>1</sub> and G<sub>7</sub> with respective crop growth rates (8.43 g/m<sup>2</sup>/day, 7.66 g/m<sup>2</sup>/day and 7.14 g/m<sup>2</sup>/day) were at par with the genotype G<sub>2</sub>. There was an increase in the crop growth rate of the wheat from 25 DAS to 100 DAS. But, later on it was a sudden decline in the crop growth rate after 100 DAS and is presented in Fig. 1. Initially, the increase in the grow rate is gradual and happened to be rapid later on. This is due to the maximum growth during the vegetative phase and maximum production of

dry matter in the early growth stages of the Plant. When the wheat crop achieved the maturity stage the growth rate declines immediately. These results were in match up with those reported by Alam [7].

### 3.1.5 Relative growth rate (g/g/day)

There was no significant variation among the genotypes at different intervals during all the crop growth stages. Higher relative growth rate was recorded in initial growth stages during 25 - 50 DAS in all the genotypes and is presented in Table 1. The genotype G<sub>9</sub> recorded maximum relative growth rate (0.087 g/g/day) during 25 - 50 DAS. During the further intervals there was more reduction trend observed in the relative growth rate when compared to the initial intervals. This is due to the crop maturity and sudden halting in the vegetative growth. There was a declining growth rate trend recorded in all the genotypes. The studies of Akhtar et al. [8] showed similar outcomes.

## 3.2 Yield and Yield Attributes

### 3.2.1 Length of the spike (cm)

The length of the spike was recorded and presented in Table 2. There was a significant variance observed in the spike length among all the wheat genotypes. The longer spike length (12.79 cm) was recorded in the genotype G<sub>2</sub>. However, the genotype G<sub>9</sub> with the spike length (12.19 cm) was at par with the genotype G<sub>2</sub>. Length of the spike is a genetic character and indirectly involves in the crop yield. Longer spike may aid in more number of grains that leads to higher yield. These results were in line with those reported by Gobinda et al. [9].

### 3.2.2 Number of Grains per Spike

Data related to number of grains/spike was embodied in Table 2. Significantly higher number of grains per spike (63.41) was recorded in the genotype G<sub>2</sub>. However, the genotypes G<sub>9</sub> (62.83) and G<sub>1</sub> (60.08) were at par with the number of grains per spike to the genotype G<sub>2</sub>. Increase in the number of irrigations favours the percentage increase in the filled grains per spike [10]. Number of grains per spike is a yield attributing character and affects the grain yield. Many factors influence the number of grains/spike mainly at genotypic level and also at environmental level. Similar findings were reported by Omar et al. [11].

Table 1. Evaluation of wheat genotypes on growth attributes

Genotypes	Plant height (cm)	Number of Tillers per hill	Plant Dry weight (g/hill)	Crop Growth Rate (g/m <sup>2</sup> /day)	Relative Growth Rate (g/g/day)
NETS-101	108.89 <sup>ab</sup>	6.2 <sup>b</sup>	27.82 <sup>bc</sup>	16.11 <sup>ab</sup>	0.085
NETS-102	112.02 <sup>a</sup>	7.7 <sup>a</sup>	31.41 <sup>a</sup>	16.5 <sup>a</sup>	0.079
NETS-103	99.9 <sup>c</sup>	5.25 <sup>c</sup>	22.24 <sup>e</sup>	9.69 <sup>d</sup>	0.066
NETS-104	100.96 <sup>c</sup>	5.47 <sup>c</sup>	24.58 <sup>cde</sup>	14.15 <sup>d</sup>	0.061
NETS-105	106.52 <sup>b</sup>	5.77 <sup>bc</sup>	24.74 <sup>cde</sup>	14.08 <sup>cd</sup>	0.067
NETS-106	100.25 <sup>c</sup>	5.6 <sup>bc</sup>	22.64 <sup>de</sup>	12.3 <sup>cd</sup>	0.066
NETS-107	108.55 <sup>ab</sup>	6.15 <sup>b</sup>	26.13 <sup>c</sup>	16.26 <sup>abc</sup>	0.078
NETS-108	107.78 <sup>b</sup>	5.85 <sup>bc</sup>	25.7 <sup>cd</sup>	16.46 <sup>bcd</sup>	0.075
NETS-109	111.5 <sup>a</sup>	7.25 <sup>a</sup>	29.86 <sup>ab</sup>	16.48 <sup>a</sup>	0.087
<b>SEm (±)</b>	1.24	0.23	1.13	1.58	0.00
<b>CD (P=0.05)</b>	3.62	0.67	3.31	4.61	-

Table 2. Evaluation of wheat genotypes on yield and yield attributes

Genotypes	Spike length (cm)	No. of Grains/spike	Grain Yield (t/ha)	Straw Yield (t/ha)
NETS-101	11.8 <sup>b</sup>	60.08 <sup>abc</sup>	3.96 <sup>abc</sup>	5.82 <sup>abc</sup>
NETS-102	12.79 <sup>a</sup>	63.41 <sup>a</sup>	4.26 <sup>a</sup>	6.04 <sup>a</sup>
NETS-103	10.38 <sup>i</sup>	51.49 <sup>e</sup>	3.3 <sup>e</sup>	5.5 <sup>d</sup>
NETS-104	10.88 <sup>def</sup>	54.99 <sup>cde</sup>	3.43 <sup>de</sup>	5.57 <sup>cd</sup>
NETS-105	11.15 <sup>cde</sup>	55.16 <sup>cde</sup>	3.66 <sup>cd</sup>	5.57 <sup>bcd</sup>
NETS-106	10.83 <sup>ef</sup>	53.99 <sup>de</sup>	3.39 <sup>de</sup>	5.56 <sup>d</sup>
NETS-107	11.69 <sup>bc</sup>	58.16 <sup>bcd</sup>	3.84 <sup>bc</sup>	5.76 <sup>bcd</sup>
NETS-108	11.47 <sup>bcd</sup>	55.16 <sup>cde</sup>	3.73 <sup>bcd</sup>	5.59 <sup>bcd</sup>
NETS-109	12.1 <sup>ab</sup>	62.83 <sup>ab</sup>	4.01 <sup>ab</sup>	5.83 <sup>ab</sup>
<b>SEm (±)</b>	0.22	1.74	0.12	0.09
<b>CD (P=0.05)</b>	0.65	5.09	0.35	0.26




Fig. 1. Crop growth rate (g/m<sup>2</sup>/day) in different wheat genotypes




Fig. 2. Relative growth rate (g/g/day) in different wheat genotypes

### 3.2.3 Grain Yield (t/ha)

The genotypic effect on grain yield is found to be significant in the wheat crop among all the genotypes and is presented in Table 2. The genotype G<sub>2</sub> dominated all the other genotypes

with higher grain yield (4.26 t/ha). However, the genotypes G<sub>9</sub> (4.01 t/ha) and G<sub>1</sub> (3.96 t/ha) with their respective grain yield were found to be at par with the genotype G<sub>2</sub>. Higher grain yields are usually associated with delayed maturity [12]. The Higher grain yield in the genotype G<sub>2</sub> was

due to the longer spike, maximum number of grains and other higher records of yield attributes. Karman and Akhtas [13] observed compatible findings.

### 3.2.4 Straw Yield (t/ha)

Data related to straw yield was evaluated and tabulated in Table 2. The genotype G<sub>2</sub> was recorded with higher straw yield (6.04 t/ha). However, the genotypes G<sub>9</sub> and G<sub>1</sub> with the straw yield (5.83 t/ha and 5.82 t/ha) were at par with the genotype G<sub>2</sub>. The better vegetative growth might have obviously resulted into higher straw yield. Higher plant height, maximum number of tillers and higher dry matter accumulation results in the higher straw yield. Maintenance of favourable soil moisture balance in the crop root zone may also results in higher biomass accumulation. These results were in similar to the observations recorded by Nayak et al. [14].

## 4. CONCLUSION

This study concluded that the wheat genotype NETS-102 was found to be productive and more effective in cultivation with higher growth and yield attributes. As the cost of cultivation is same for all the genotypes, NETS-102 genotype will be economically viable due to higher yield in wheat.

## ACKNOWLEDGEMENT

The authors are thankful to department of Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj – 211007, Uttar Pradesh, India for providing us necessary facilities to undertake the studies.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Shewry PR. Wheat. Journal of Experimental Botany. 2009;60(6):1573-1533.
2. Sharma D, Singh R, Tiwari R, Kumar R, Gupta V. Wheat Responses and Tolerance to Terminal Heat Stress: A Review. In Hasanuzzaman M, Nahar K, Hossain MA. (Eds.), Wheat Production in Changing Environments: Responses, Adaptation and Tolerance. 2019;149–173.
3. Gomez KA, Gomez AA. Statistical Procedures for Agricultural Research. (2 ed). John Wiley and Sons, New York. 1984;680.
4. Bhutto TA, Buriro M, Wahocho NA, Jakhro MI, Zulfiqar AA, Vistro R, Abbasi F, et al. Evaluation of Wheat Cultivars for Growth and Yield Traits under Agro-Ecological Condition of Tandojam. Pakistan Journal of Agricultural Research. 2020;34(1):136-143.
5. Ghanbari A, Malidarreh. The Effect of Complementary Irrigation in Different Growth Stages on Yield, Qualitative and Quantitative Indices of the Two Wheat (*Triticum aestivum* L.) Cultivars. International Journal of Agricultural and Bio-systems Engineering. 2010;4(5).
6. Sing H, Singh S, Lal M, Kumar S, Singh L. Response of irrigation schedule and nitrogen on growth and yield of wheat (*Triticum aestivum* L.) under irrigated conditions of Punjab. Journal of Pharmacognosy and Phytochemistry. 2020;9(3):1020-1023.
7. Alam MS. Growth and yield potentials of wheat as affected by management practices. African Journal of Agricultural Research. 2013;8(47):6068-6072.
8. Akhtar M, Mosleh U, Hossain I. Study on the Growth and Yield Potential of Promising Wheat Genotypes under Modified Agronomical Practices. World Journal of Research and Review. 2018;6(1):41-49.
9. Gobinda P, Yadav L, Tiwari A, Khatri HB, Basnet S, Bhattarai K, Gyawali B, Rawal N, Khatri N. Analysis of Yield Attributing Characters of Different Genotypes of Wheat in Rupandehi, Nepal. International Journal of Environment, Agriculture and Biotechnology. 2017;2(5).
10. Ahmad A, Kumar R. Effect of irrigation scheduling on the growth and yield of Wheat genotypes. Agricultural Science Digest. 2015;35(3):199-202.
11. Omar K, Rafiqul I, Hasanuzzaman MD, Imanur R, Shoebur R, Kajal MD, Datta T. Evaluation of Some Improved Wheat (*Triticum aestivum* L.) Genotypes for Growth and Yield Potential. International Journal of Plant & Soil Science. 2017; 16(3):1-7.
12. Zhang G, Wang Y, Guo Y, Zhao Y, Kong F, Li S. Characterization and mapping of

- QTLs on chromosome 2D for grain size and yield traits using a mutant line induced by EMS in wheat. *The Crop Journal*. 2015;1-10.
13. Karman M, Aktas H. Comparison of the Agricultural Characteristics of Bread Wheat (*Triticum aestivum L.*) Genotypes based on Irrigated Conditions in Different Locations. *Manas Journal of Agriculture Veterinary and Life Sciences*. 2020;10(1):33-42.
  14. Nayak MK, Patel HR, Prakash V, Kumar A. Influence of Irrigation Scheduling on Crop Growth Yield and Quality of Wheat. *Journal of Agri Search*. 2015;2(1):65-68.

---

© 2022 Prashanth et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/88739>