



## **Anti-diabetic and Antioxidant Activity of *Pterocarpus santalinus* and Stevia Herbal Formulation**

**V. M. Nivedha<sup>a</sup>, R. Priyadarshini<sup>a\*#≡</sup>, S. Rajeshkumar<sup>b#≡</sup> and Palati Sinduja<sup>a\*≡</sup>**

<sup>a</sup> Department of Pathology, Saveetha Dental College, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai – 600077, Tamil Nadu, India.

<sup>b</sup> Department of Pharmacology, Saveetha Dental College, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, India.

### **Authors' contributions**

This work was carried out in collaboration among all authors. Author VMN designed the study, performed the methods and wrote the first draft of the manuscript and author RP performed the statistical analysis and wrote the protocol, author SR managed the analyses of the study and managed the literature searches and author PS managed the final drafting and editing of the manuscript. All authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/JPRI/2021/v33i62B35178

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/81770>

**Original Research Article**

**Received 23 October 2021**  
**Accepted 28 December 2021**  
**Published 29 December 2021**

### **ABSTRACT**

**Introduction:** *Pterocarpus santalinus* have their application in the pharmaceutical, cosmetic, agricultural, and food industries. *Stevia* (*Stevia rebaudiana*) is a natural, non-caloric sugar substitute that is a rich source of a pharmacologically significant glycoside. Proper diet, exercise, and pharmacological interventions contribute to overcoming diabetes.

**Aim:** The present study aims to assess the anti-diabetic and antioxidant activity of *Pterocarpus santalinus* and *Stevia* herbal formulation.

**Materials and Methods:** Preparation of plant extract followed by antidiabetic and antioxidant activity.

**Results:** Results were tabulated and graphically analyzed using SPSS software. As the concentration increased the percentage of inhibition also increased in both antidiabetic and antioxidant activity.

**Conclusion:** The present study concluded that *Pterocarpus santalinus* and *Stevia* herbal formulation has antidiabetic and antioxidant activity.

<sup>#</sup> Assistant Professor;

<sup>≡</sup> Dr.;

\*Corresponding author: E-mail: Priyadarshinir.sdc@saveetha.com;

**Keywords:** *Pterocarpus santalinus*; *stevia*; antidiabetic; antioxidant; innovative method.

## 1. INTRODUCTION

Herbal drugs are commonly considered as less toxic and hence no side effects are associated [1]. Diabetes is a growing health concern and an emerging global epidemic. It is a major challenge to control diabetes. Natural herbs have always been a source of drugs for humans since time immemorial. Traditional medicine in India has used these natural herbs to control diabetics. According to the World Health Organization (WHO), 90% of the total population in most of the developing countries use plants and their products as traditional medicine for primary health care. There are about 800 plants with anti-diabetic potential. Thus there is a growing demand for research on natural products with anti-diabetic properties. Numerous studies have defined the benefits of medicinal plants with anti-hyper-glycaemic effects in controlling diabetes mellitus [2–4]. In the Past it is clearly shown that medicinal plants have been used in traditional healing globally for a long time to treat diabetes; this is because such herbal plants have hypoglycemic properties and other beneficial properties.

*Pterocarpus santalinus*: The medicinal plants have their application in the pharmaceutical, cosmetic, agricultural, and food industries. The use of medicinal herbs for curing disease has been marked in the past Saga of all civilizations. Humans in the prehistoric era were not aware of the health hazards and their association with irrational therapy. With the commencement of research in medicine, it was concluded that plants contain active principles, which are responsible, for the curative action of the disease [5,6]. Before the synthetic era, man was fully dependent on medicinal herbs for the prevention and treatment of various diseases. With the introduction of scientific procedures, medical practitioners were able to understand toxic principles present in the green flora. Medicinal plants serve as a natural source for most of the bioactive medicinal compounds [7,8]. Medicinal plants like *Pterocarpus santalinus* are commonly used in traditional medicine which is rich in phenols and flavonoids. *Pterocarpus santalinus* shows antidiabetic activity and helps in decreasing the increased glucose levels and improving hyperlipidemia and restoring the insulin levels [9].

*Stevia rebaudiana* (Bertoni) Bertoni (*Stevia*) is a natural, non-caloric sugar substitute that is a rich

source of a pharmacologically significant glycoside. Proper diet, exercise, and pharmacological interventions contribute to overcoming diabetes. The pharmacological drugs indicated for treating diabetes are expensive and have certain adverse side effects [10]. Therefore, herbs are considered a natural source of drugs that have strong antioxidant activities to be more effective against diabetes. *S. rebaudiana* (family Asteraceae) is a traditional plant that is famous because of its sweet taste and beneficial effects on blood glucose regulation. *S. rebaudiana* is popularly known as stevia, sweet weed, honey leaf, and the sweet herb of Paraguay [11]. *Stevia* leaves contain a complex mixture of diterpene glycosides including stevioside, steviolbioside, rebaudiosides (A, B, C, D, E), and dulcoside A, the major sweet constituents are stevioside and rebaudioside A. Natural non-caloric sweetener stevioside (a major component of stevia) is considered to be sweeter than sucrose and is extensively used as a non-caloric sugar substitute in many application like foods, medicine, beverage, cosmetics, winemaking, household chemical industry, and other food industries. It cuddles anti-hyperglycaemic, anti-hypertensive, anti-oxidant, anti-tumor, anti-diarrheal, diuretic, gastro and renal-protective, and immunomodulatory properties [12]. Herbal products like *Pterocarpus Santalinus* and *Stevia* are less toxic and have fewer side effects and so can be used daily. This study aims to assess the antidiabetic and antioxidant activity of *Pterocarpus santalinus* and *stevia* herbal formulation.

## 2. MATERIALS AND METHODS

### 2.1 Preparation of Plant Extract

Commercially available dry powder of *Pterocarpus santalinus* and *stevia* was readily available and was used for this experiment (Fig. 1). This experiment was conducted in Saveetha Dental College, Chennai, Tamilnadu. The experiment was carried out by dissolving 1 g of *Pterocarpus santalinus* and *stevia* in 100 ml of water. The moisture was then boiled in a heating mantle at 70°C for up to 10 minutes. The boiled mixture was then filtered using Whatman number 1 filter paper to obtain the plant extract. Then 40 µl of the plant extract was measured using a measuring cylinder which might not be that accurate and the mixture was added to 60 ml of 1 mM dissolved in 60 ml of distilled water (Fig. 2).



**Fig. 1** Weighing 1g of *Pterocarpus santalinus* and *Stevia* herbal formulation



**Fig. 2.** Heating *Pterocarpus santalinus* and *Stevia* extract at 70°C for 10 minutes for the preparation of plant extract

## 2.2 Antioxidant Activity

### 1. DPPH METHOD(2,2-diphenyl-1-picrylhydrazyl)

#### Antioxidant activity

DPPH assay was used to test the antioxidant activity of *Pterocarpus santalinus* and *Stevia* herbal formulation. Diverse concentrations (2-10 µg/ml) of plant extract were mixed with 1 ml of 0.1 mM DPPH in methanol and 450 µl of 50 mM Tris HCl buffer (pH 7.4) and incubated for 30 minutes. Later, the reduction in the quantity of DPPH free radicals was assessed dependent on the absorbance at 517 nm. BHT was employed as control. (Fig. 3) The percentage of inhibition was determined from the following equation,

$$\% \text{ inhibition} = \frac{\text{Absorbance of control} - \text{Absorbance of test sample} \times 100}{\text{Absorbance of control}}$$



**Fig. 3.** Antioxidant activity performed by adding 1ml of DPPH in methanol and then the reduction in the quantity of DPPH free radicals was assessed dependent on the absorbance at 517nm

## 2.3 In-vitro anti-diabetic Assay

The in Vitro anti-diabetic assay was performed using two different techniques:

## 2.4 Alpha-Amylase Inhibitory Assay

Alpha-amylase inhibition was determined by quantifying the amount of maltose liberated during the experiment. The method reported by Bhutkar and Bhise has been followed (Bhutkar and Bhise, 2012). Different concentrations of nanoparticles (20, 40, 60, 80, 100 µl) was pre-incubated with 100% α amylase solution (1U/mL) at room temperature for 30 minutes. 100µl of starch solution (1%w/v) was further added to it and the mixture was incubated at room temperature for 10 minutes. 100µl of 96mM (3,5-dinitrosalicylic acid solution)DNSA reagent was added to it stop the reaction and the solution was heated in a water bath for 5 minutes. (Fig. 4) Control was maintained where the equal quantity of enzyme extract was replaced by sodium phosphate buffer maintained at a pH value of 6.9. Reading was measured at 540nm. The experiment was performed in triplicate. Acarbose was used as a positive control.

The %inhibition was calculated using the formulae-

$$\% \text{ inhibition} = C - T / C * 100$$

Where, C=control, T=test sample

Validation was done with nano experts followed by correlation analysis using SPSS

software. Only antidiabetic and antioxidant activity was performed. In future studies, cytotoxicity and antimicrobial activity will be performed.




**Fig. 4. Antidiabetic activity was performed by adding DNSA reagent to stop the reaction and the solution was heated in a water bath for 5 minutes**

### 3. RESULTS

**Table 1. Antioxidant activity of *Pterocarpus santalinus* and *Stevia* herbal formulation. When the concentration was 10µl,20µl,30µl,40µl,50µl the percentage of inhibition was 0.16%,0.28%,0.42%,0.58%,0.73 respectively**


ANTIOXIDANT ACTIVITY		
Concentration	Percentage of inhibition	Wavelength
10µl	0.16	5.17nm
20µl	0.28	5.17nm
30µl	0.42	5.17nm
40µl	0.58	5.17nm
50µl	0.73	5.17nm



**Fig. 5.** The graph represents the antioxidant activity of the extract. The X-axis represents different concentrations and Y-axis represents the antioxidant activity of the plant extract. When the concentration was 10 $\mu$ l,20 $\mu$ l,30 $\mu$ l,40 $\mu$ l,50 $\mu$ l the percentage of inhibition was 0.16%,0.28%,0.42%,0.58%,0.73 respectively

**Table 2.** Antidiabetic activity of *Pterocarpus santalinus* and *Stevia* herbal formulation. When the concentration was 10 $\mu$ l,20 $\mu$ l,30 $\mu$ l,40 $\mu$ l,50 $\mu$ l the percentage of inhibition was 0.06%,0.15%,0.20%,0.24%,0.31 respectively

ANTIDIABETIC ACTIVITY		
Concentration	Percentage of inhibition	Wavelength
10 $\mu$ l	0.06	540nm
20 $\mu$ l	0.15	540nm
30 $\mu$ l	0.20	540nm
40 $\mu$ l	0.24	540nm
50 $\mu$ l	0.31	540nm



**Fig. 6.** The graph represents the antidiabetic activity of the extract. The X-axis represents different concentrations and the Y-axis represents the antidiabetic activity of the plant extract. When the concentration was 10 $\mu$ l, 20 $\mu$ l, 30 $\mu$ l, 40 $\mu$ l, 50 $\mu$ l the percentage of inhibition was 0.06%, 0.15%, 0.20%, 0.24%, 0.31 respectively

#### 4. DISCUSSION

In Table 1 & Fig. 5 when the concentration was 10 $\mu$ l, 20 $\mu$ l, 30 $\mu$ l, 40 $\mu$ l, 50 $\mu$ l the percentage of inhibition was 0.16%, 0.28%, 0.42%, 0.58%, 0.73 respectively. We can see that antioxidant activity increased with an increase in concentration. Streptozotocin-induced Diabetes mellitus is exemplified not only by impaired glucose tolerance and hyperglycemia, as well by low antioxidant activity. The *Pterocarpus santalinus* has a strong antioxidant activity which has been demonstrated via different in vitro assays, as well as by using liver-slice slides [13,14]. Even though the  $\beta$ -cell cytotoxic effect of Streptozotocin induction is not fully understood, it is considered to result in the inhibition of free radical scavenger enzyme production. Oxidative stress is held responsible for tissue damage and  $\beta$ -cell dysfunction. It is observed that an increase in HDL and reduction in LDL, has been observed in the combination with vitamin E and P. Body, muscle, heart, kidney, and liver weight are completely perceived to benefit from the combinational approach [15]. Weight loss due to excessive breakdown of tissue proteins is also a complication of Diabetes Mellitus. Weight loss in

treated groups either with *P. santalinus* or vitamin E alone or in combination was less significant, as compared to the untreated-diabetic group, which lost a considerable amount of weight. The pooled treatment with *Pterocarpus santalinus* and vitamin E exhibited promising results. An improvement in body weight, lipid profile, glucose tolerance, biochemical parameters such as urea, creatinine, and histological reversal of the nephropathy and activity of antioxidant enzymes were noticed. These results illustrated the anti-diabetic activity antioxidant action of *Pterocarpus santalinus*. The previous study was done by M. Eshrat Halim. stands in line with our present study stating the antidiabetic and antioxidant activity of *Pterocarpus santalinus*. [16].

Free radicals are electrically charged molecules, and a disproportionate generation of these free radicals is linked to many human diseases. Reactive oxygen species (ROS) along with hydroxyl radicals cause damage to the structure and function of cells, oxidation of lipids, proteins, and DNA; leads to the development of various diseases [17]. The free radical scavenging activity of extracts of the leaves of *Pterocarpus santalinus* has been evaluated through in vitro

studies. The methanolic extract of the leaves exhibited radical scavenging activity for diphenyl picrylhydrazyl (DPPH), nitric oxide, and hydrogen peroxide. Studies have proved that Fe<sup>3+</sup> reducing capacity and DPPH radical scavenging activity in the methanolic extract of heartwood. As the concentration increased the antidiabetic and antioxidant activity also increased which was compared with butylated hydroxyanisole (BHA). Few other studies also showed the strong antioxidant activity against free radicals like DPPH, 2,2'-azino-bis, hydroxyl, superoxide, and hydrogen peroxide. The present study is in line with the previous study done by Saradamma Bulle [18].

In Table 2 & Fig. 6 when the concentration was 10µl, 20µl, 30µl, 40µl, 50µl the percentage of inhibition was 0.16%, 0.28%, 0.42%, 0.58%, 0.73 respectively. We can see that antidiabetic activity increased with an increase in concentration. *Stevia* leaf extract has been conventionally used to treat diabetes. Their ingestion causes a slender suppression of plasma glucose levels and significantly increases the glucose tolerance in human beings. Steviol glycosides have an enhancing effect on insulin secretion by directly acting on β-cells without altering the K<sup>+</sup> - ATP channel activity and cAMP level in the islets, thus recording stevioside and steviol as potent antihyperglycemic agents. Stevioside regulates blood glucose levels by enhancing not only insulin secretion but also insulin utilization [18,19]. Overall, *Stevia* possesses the ability to boost the insulin effect on cell membranes, increase insulin production, stabilize glucagon secretion and blood sugar levels, when carbohydrates are ingested improves glucose tolerance and lowers postprandial blood sugar levels in both animals as well as humans. Alternatively, it is described that *Stevia* is shown to provide a comprehensive set of mechanisms that counter the mechanics of type II diabetes and its eventual complications. Thus, to support healthy glucoregulation sugars can be replaced with steviol glycosides or stevioside of *Stevia* leaf. Adding to it not only the leaves of *Stevia*, dried or in powder form in supplementary food products of diabetic patients aid in increasing the natural sweetness and also help in rejuvenating the pancreatic gland [20].

The serum insulin level in the diabetic control group decreased due to Streptozotocin that resulted in diabetes by the rapid depletion of β-cells, which reduced the insulin release. An inadequate release of insulin causes

hyperglycemia, which results in oxidative damage by the generation of reactive oxygen species and the development of diabetic complications [20,21]. When *Stevia* aqueous extracts at different dose levels the insulin levels improve significantly due to the presence of natural components (stevioside) in *Stevia* leaves that are related to inhibition of hepatic expression of phosphoenolpyruvate carboxykinase and gluconeogenesis together with stimulation of hepatic glycogen synthesis that increased insulin secretion and insulin sensitivity. Evidence from other studies revealed that through the PPARγ-dependent mechanism *Stevia* aqueous extract elevates the insulin level due to stevioside which acts on pancreatic tissue, exerts beneficial anti-hyperglycemic effects. The study done by Uswa Ahmad states that *Stevia* aqueous extract enhanced caloric management and weight control by decreasing the feed intake and body weight gain, as per the present Study [13].

The Intent of an antidiabetic therapy in insulin-dependent patients (Type 1 diabetes) and insulin-non dependent patients (Type 2 diabetes) is to attain normoglycemia and reduce insulin resistance to improve metabolic control and prevent future diabetic patients [22]. An effective strategy for the management of Type 2 diabetes is the inhibition of the two enzymes, alpha-amylase, and alpha-glucosidase, to slow the rate of absorption of carbohydrates, thus altering the postprandial rise of blood sugar and reducing the effects of diet on hyperglycemia. Undue inhibition of alpha-amylase due to alpha-glucosidase inhibitors such as acarbose carries certain side effects (bloating, flatulence, meteorism, and diarrhea) as a result of an abnormal bacterial fermentation of undigested carbohydrates in the colon [23,24].

According to a previous study, the alpha-amylase and alpha-glucosidase inhibitors derived from plants have mild inhibitory activity against alpha-amylase and strong inhibitory activity against alpha-glucosidase, which indicates their utilization in therapy for postprandial hyperglycemia with fewer side effects. This research reveals that the aqueous extracts of *Stevia rebaudiana* have significant inhibitory activity against the alpha-amylase and alpha-glucosidase enzymes and show capability for the treatment of both types of diabetes. Our team has extensive knowledge and research experience that has translate into high quality publications [23-44]. This activity can be performed with more concentration, the only anti-

diabetic, and the antioxidant property is analyzed, clinical trials can be done. Because of its antioxidant property, it can be used as an anti-cancer drug. As it also shows anti-diabetic properties it can be standardized and given for patients with all types of diabetes.

## 5. CONCLUSION

The present study suggests that aqueous extract from *Pterocarpus santalinus* and *Stevia* had shown better anti-diabetic and antioxidant activity by decreasing the random blood glucose level and fasting blood glucose and glycosylated (HbA1c) hemoglobin while insulin and liver glycogen levels significantly increased. It is implicit from the results that *Stevia* extract has anti-diabetic effects, and therefore could be used as a natural anti-diabetic drug for the treatment of diabetes and its associated complications.

## NOTE

The study highlights the efficacy of "Herbal" which is an ancient tradition, used in some parts of India. This ancient concept should be carefully evaluated in the light of modern medical science and can be utilized partially if found suitable.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

It is not applicable.

## ACKNOWLEDGEMENT

The authors are thankful to Saveetha Institute of Medical and Technical Sciences, Saveetha Dental College and Hospitals, Saveetha University for giving a platform to conduct the study.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Karimi A, Majlesi M, Rafieian-Kopaei M. Herbal versus synthetic drugs; beliefs and facts. *J Nephroarmacol* [Internet]. 2015;4(1):27–30. Available:

<https://www.ncbi.nlm.nih.gov/pubmed/28197471>

2. Kooti W, Farokhipour M, Asadzadeh Z, Ashtary-Larky D, Asadi-Samani M. The role of medicinal plants in the treatment of diabetes: a systematic review. *Electron Physician* [Internet]. 2016;8(1):1832–42. Available:<http://dx.doi.org/10.19082/1832>
3. Preethikaa S, Brundha MP. Awareness of diabetes mellitus among general population. *Research Journal of Pharmacy and Technology* [Internet]. 2018;11(5):1825–9. Available: <https://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=11&issue=5&article=024>
4. Timothy CN, Samyuktha PS, Brundha MP. Dental pulp stem cells in regenerative medicine-A literature review. *Research Journal of Pharmacy and Technology* [Internet]. 2019;12(8):4052–6. Available:<https://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=8&article=088>
5. Chen S-L, Yu H, Luo H-M, Wu Q, Li C-F, Steinmetz A. Conservation and sustainable use of medicinal plants: problems, progress, and prospects. *Chin Med* [Internet]. 2016;11:37. Available:<http://dx.doi.org/10.1186/s13020-016-0108-7>
6. Brundha MP. A comparative study-The role of skin and nerve biopsy in hansen's disease. *Res J Pharm Biol Chem Sci* [Internet]. 2015;7(10):837. Available: [https://www.researchgate.net/profile/Brundha-Mp-2/publication/283561218\\_A\\_comparative\\_study-\\_the\\_role\\_of\\_skin\\_and\\_nerve\\_biopsy\\_in\\_hansen's\\_disease/links/5892ba5d458515aeac946451/A-comparative-study-the-role-of-skin-and-nerve-biopsy-in-hansens-disease.pdf](https://www.researchgate.net/profile/Brundha-Mp-2/publication/283561218_A_comparative_study-_the_role_of_skin_and_nerve_biopsy_in_hansen's_disease/links/5892ba5d458515aeac946451/A-comparative-study-the-role-of-skin-and-nerve-biopsy-in-hansens-disease.pdf)
7. Ojuederie OB, Babalola OO. Microbial and plant-assisted bioremediation of heavy metal polluted environments: A review. *Int J Environ Res Public Health* [Internet]. 2017;14(12). Available:<http://dx.doi.org/10.3390/ijerph14121504>
8. Harsha L, Brundha MP. Prevalence of dental developmental anomalies among men and women and its psychological effect in a given population. *Journal of Pharmaceutical Sciences and Research; Cuddalore* [Internet]. 2017;9(6):869–73.

- Available:<https://www.proquest.com/scholarly-journals/prevalence-dental-developmental-anomalies-among/docview/1917938864/se-2>
9. El-Badawy RE, Ibrahim KA, Hassan NS, El-Sayed WM. ameliorates streptozotocin-induced diabetes mellitus via anti-inflammatory pathways and enhancement of insulin function. *Iran J Basic Med Sci* [Internet]. 2019;22(8):932–9. Available: <http://dx.doi.org/10.22038/ijbms.2019.34998.8325>
  10. Marín-Peñalver JJ, Martín-Timón I, Sevillano-Collantes C, Del Cañizo-Gómez FJ. Update on the treatment of type 2 diabetes mellitus. *World J Diabetes* [Internet]. 2016;7(17):354–95. Available:<http://dx.doi.org/10.4239/wjd.v7.i17.354>
  11. Nasri H, Baradaran A, Shirzad H, Rafieian-Kopaei M. New concepts in nutraceuticals as alternative for pharmaceuticals. *Int J Prev Med* [Internet]. 2014;5(12):1487–99. Available: <https://www.ncbi.nlm.nih.gov/pubmed/25709784>
  12. Ntchapda F, Barama J, Kemeta Azambou DR, Etet PFS, Dimo T. Diuretic and antioxidant activities of the aqueous extract of leaves of *Cassia occidentalis* (Linn.) in rats. *Asian Pac J Trop Med* [Internet]. 2015;8(9):685–93. Available: <http://dx.doi.org/10.1016/j.apjtm.2015.07.030>
  13. Ahmad U, Ahmad RS, Arshad MS, Mushtaq Z, Hussain SM, Hameed A. Antihyperlipidemic efficacy of aqueous extract of *Stevia rebaudiana* Bertoni in albino rats. *Lipids Health Dis* [Internet]. 2018;17(1):175. Available:<http://dx.doi.org/10.1186/s12944-018-0810-9>
  14. Kumar D. Anti-inflammatory, analgesic, and antioxidant activities of methanolic wood extract of *Pterocarpus santalinus* L. *J Pharmacol Pharmacother* [Internet]. 2011; 2(3):200–2. Available: <http://dx.doi.org/10.4103/0976-500X.83293>
  15. Davidson LE, Kelley DE, Heshka S, Thornton J, Pi-Sunyer FX, Boxt L, et al. Skeletal muscle and organ masses differ in overweight adults with type 2 diabetes. *J Appl Physiol* [Internet]. 2014;117(4):377–82. Available:<http://dx.doi.org/10.1152/jappphysiol.01095.2013>
  16. M. Eshrat Halim AM. The effects of the aqueous extract of *Pterocarpus santalinus* heartwood and vitamin E supplementation in streptozotocin-induced diabetic rats. *Journal of Medicinal Plants* [Internet]. 2011;5(3):398–409. Available: <https://academicjournals.org/journal/JMPR/article-full-text-pdf/736711624896>
  17. Phaniendra A, Jestadi DB, Periyasamy L. Free radicals: properties, sources, targets, and their implication in various diseases. *Indian J Clin Biochem* [Internet]. 2015;30(1):11–26. Available: <http://dx.doi.org/10.1007/s12291-014-0446-0>
  18. Bulle S, Reddyvari H, Nallanchakravarthula V, Vaddi DR. Therapeutic potential of *Pterocarpus santalinus* L.: An update. *Pharmacogn Rev* [Internet]. 2016;10(19):43–9. Available: <http://dx.doi.org/10.4103/0973-7847.176575>
  19. Ajami M, Seyfi M, Abdollah Pouri Hosseini F, Naseri P, Velayati A, Mahmoudnia F, et al. Effects of stevia on glycemic and lipid profile of type 2 diabetic patients: A randomized controlled trial. *Avicenna J Phytomed* [Internet]. 2020;10(2):118–27. Available:<https://www.ncbi.nlm.nih.gov/pubmed/32257884>
  20. Gechev TS, Hille J, Woerdenbag HJ, Benina M, Mehterov N, Toneva V, et al. Natural products from resurrection plants: Potential for medical applications. *Biotechnol Adv* [Internet]. 2014; 32(6):1091–101. Available:<http://dx.doi.org/10.1016/j.biotechadv.2014.03.005>
  21. Zhang C, Caldwell TA, Mirbolooki MR, Duong D, Park EJ, Chi N-W, et al. Extracellular CADM1 interactions influence insulin secretion by rat and human islet  $\beta$ -cells and promote clustering of syntaxin-1. *Am J Physiol Endocrinol Metab* [Internet]. 2016;310(11):E874–85. Available: <http://dx.doi.org/10.1152/ajpendo.00318.2015>
  22. Otto-Buczowska E, Dryżałowski M. Metabolicsyndrome in youngpatients. *Pediatr Endocrinol Diabetes Metab* [Internet]. 2015;21(1):32–6. Available:<http://dx.doi.org/10.18544/PEDM-21.01.0022>
  23. Al-Massarani SM, El-Gamal AA, Parvez MK, Al-Dosari MS, Al-Said MS, Abdel-Kader MS, et al. New cytotoxic seco-type triterpene and labdane-type diterpenes

- from nuxia oppositifolia. *Molecules* [Internet]. 2017;22(3). Available:<http://dx.doi.org/10.3390/molecules22030389>
24. Hannah R, Ramani P, Brundha MP, Sherlin HJ, Ranjith G, Ramasubramanian A, et al. Liquid paraffin as a rehydrant for air dried buccal smear. *Research Journal of Pharmacy and Technology* [Internet]. 2019;12(3):1197–200. Available: <https://www.indianjournals.com/ijor.aspx?target=ijor:rjpt&volume=12&issue=3&article=038>
  25. JC, Ruiz-Ruiz YB, Moguel-Ordoñez AJ, Matus-Basto, Segura-Campos MR. Antidiabetic and antioxidant activity of *Stevia rebaudiana* extracts (Var. Morita) and their incorporation into a potential functional bread. *J Food Sci Technol* [Internet]. 2015;52:7893–903. Available: <https://link.springer.com/article/10.1007/s13197-015-1883-3>
  26. Jayaseelan VP, Paramasivam A. Emerging role of NET inhibitors in cardiovascular diseases [Internet]. *Hypertension Research*. 2020;43:1459–61. Available: <http://dx.doi.org/10.1038/s41440-020-0527-9>
  27. Anita R, Paramasivam A, Priyadharsini JV, Chitra S. The m6A readers and aberrations associated with metastasis and predict poor prognosis in breast cancer patients. *Am J Cancer Res* [Internet]. 2020;10(8):2546–54. Available: <https://www.ncbi.nlm.nih.gov/pubmed/32905518>
  28. Sivakumar S, Smiline Girija AS, Vijayashree Priyadharsini J. Evaluation of the inhibitory effect of caffeic acid and gallic acid on tetR and tetM efflux pumps mediating tetracycline resistance in *Streptococcus* sp., using computational approach [Internet]. *Journal of King Saud University - Science*. 2020;32:904–9. Available: <http://dx.doi.org/10.1016/j.jksus.2019.05.003>
  29. Girija ASS, Smiline Girija AS. Delineating the Immuno-dominant antigenic vaccine peptides against gacs-sensor kinase in *acinetobacter baumannii*: An *In silico* investigational approach [Internet]. *Frontiers in Microbiology*. 2020;11. Available: <http://dx.doi.org/10.3389/fmicb.2020.02078>
  30. Girija ASS, Smiline Girija AS. Fox3 CD25 CD4 T-regulatory cells may transform the nCoV's final destiny to CNS! [Internet]. *Journal of Medical Virology*. 2021; 93:5673–5. Available: <http://dx.doi.org/10.1002/jmv.26482>
  31. Jayaseelan VP, Ramesh A, Arumugam P. Breast cancer and DDT: Putative interactions, associated gene alterations, and molecular pathways. *Environ Sci Pollut Res Int* [Internet]. 2021; 28(21):27162–73. Available:<http://dx.doi.org/10.1007/s11356-021-12489-6>
  32. Arumugam P, George R, Jayaseelan VP. Aberrations of m6A regulators are associated with tumorigenesis and metastasis in head and neck squamous cell carcinoma. *Arch Oral Biol* [Internet]. 2021;122:105030. Available:<http://dx.doi.org/10.1016/j.archoralbio.2020.105030>
  33. Kumar SP, Praveen Kumar S, Smiline Girija AS, Vijayashree Priyadharsini J. Targeting NM23-H1-mediated Inhibition of tumour metastasis in viral hepatitis with bioactive compounds from *ganoderma lucidum*: A computational study [Internet]. *Indian Journal of Pharmaceutical Sciences*. 2020;82. Available:<http://dx.doi.org/10.36468/pharmaceutical-sciences.650>
  34. Girija SA, Priyadharsini JV, Paramasivam A. Prevalence of carbapenem-hydrolyzing OXA-type  $\beta$ -lactamases among *acinetobacter baumannii* in patients with severe urinary tract infection. *Acta Microbiol Immunol Hung* [Internet]. 2019;67(1):49–55. Available: <http://dx.doi.org/10.1556/030.66.2019.030>
  35. Priyadharsini JV, Paramasivam A. RNA editors: Key regulators of viral response in cancer patients. *Epigenomics* [Internet]. 2021;13(3):165–7. Available:<http://dx.doi.org/10.2217/epi-2021-0001>
  36. Mathivadani V, Smiline AS, Priyadharsini JV. Targeting epstein-barr virus nuclear antigen 1 (EBNA-1) with *Murraya koengii* bio-compounds: An *In-silico* approach. *Acta Virol* [Internet]. 2020;64(1):93–9. Available: [http://dx.doi.org/10.4149/av\\_2020\\_111](http://dx.doi.org/10.4149/av_2020_111)
  37. Girija As S, Priyadharsini J V, A P. Prevalence of and complex in elderly population with urinary tract infection (UTI). *Acta Clin Belg* [Internet]. 2021;76(2):106–12. Available:

- <http://dx.doi.org/10.1080/17843286.2019.1669274>
38. Anchana SR, Girija SAS, Gunasekaran S, Priyadharsini VJ. Detection of gene in carbapenem-resistant strains and targeting with biocompounds. Iran J Basic Med Sci [Internet]. 2021;24(5):690–8. Available:<http://dx.doi.org/10.22038/IJBMS.2021.52852.11917>
39. Girija ASS, Smiline Girija AS, Shoba G, Vijayashree Priyadharsini J. Accessing the T-Cell and B-Cell immuno-dominant peptides from a.baumannii biofilm associated protein (bap) as vaccine candidates: A computational approach [Internet]. International Journal of Peptide Research and Therapeutics. 2021;27:37–45. Available:<http://dx.doi.org/10.1007/s10989-020-10064-0>
40. Arvind PTR, Jain RK. Skeletally anchored forsus fatigue resistant device for correction of Class II malocclusions-A systematic review and meta-analysis. Orthod Craniofac Res [Internet]. 2021;24(1):52–61. Available:<http://dx.doi.org/10.1111/ocr.12414>
41. Venugopal A, Vaid N, Jay Bowman S. Outstanding, yet redundant? After all, you may be another Choluteca Bridge! [Internet]. Seminars in Orthodontics. 2021;27:53–6. Available: <http://dx.doi.org/10.1053/j.sodo.2021.03.007>
42. Ramadurai N, Gurunathan D, Samuel AV, Subramanian E, Rodrigues SJL. Effectiveness of 2% articaine as an anesthetic agent in children: Randomized controlled trial. Clin Oral Investig [Internet]. 2019;23(9):3543–50. Available:<http://dx.doi.org/10.1007/s00784-018-2775-5>
43. Varghese SS, Ramesh A, Veeraiyan DN. Blended module-based teaching in biostatistics and research methodology: A retrospective study with postgraduate dental students. J Dent Educ [Internet]. 2019;83(4):445–50. Available:<http://dx.doi.org/10.21815/JDE.019.054>
44. Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of Streptococcus mutans, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial [Internet]. Clinical Oral Investigations. 2020;24:3275–80. Available:<http://dx.doi.org/10.1007/s00784-020-03204-9>

© 2021 Nivedha et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/81770>