



Study on Approximate Solution of Fractional Order Biological Population Model

Yunqi Xu^{1*} and Dianchen Lu¹

¹Faculty of Science, Jiangsu University, Zhenjiang 212013, Jiangsu, P.R. China.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ACRI/2018/46155

Editor(s):

(1) Dr. Nguyen Van Toan, School of Biotechnology, Vietnam National University, Ho Chi Minh City, Vietnam.

Reviewers:

(1) Omenyi, O. Louis, Alex Ekwueme Federal University, Nigeria.

(2) Abdullah Sonmezoglu, Bozok University, Turkey.

(3) Youwei Zhang, Hexi University, China.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/46155>

Received 06 October 2018

Accepted 27 December 2018

Published 22 January 2019

Method Article

ABSTRACT

In this paper, we present an algorithm of the homotopy analysis Elzaki transform method (HAETM) which is a combination of Elzaki transform method and homotopy analysis method (HAM) to solve a more general biological population model. The fractional derivatives are described by Caputo sense. The proposed method presents a procedure of constructing the set of base functions and gives the high-order deformation equations in a simple form and provides the solution in the form of a convergent series. Three examples are used to illustrate the preciseness and effectiveness of the proposed method.

Keywords: Fractional calculus; Elzaki transform; biological population model; homotopy analysis method.

2010 Mathematics Subject Classification: 35C10; 35D99.

*Corresponding author: E-mail: yqxu1018@qq.com

1 INTRODUCTION

Fractional calculus have gained importance and popularity due to its various applications in fluid mechanics, visco-elasticity, biology, electrical network, optics and signal processing and so on [1],[2],[3],[4],[5],[6],[7],[8],[9],[10]. Except in a limited number of these problems, we have difficulty to find their exact analytic solutions. An effective and easy method for solving such equations is needed.

Various powerful methods such as Differential Transform method (DTM) [11],[12],[13], Adomian Decomposition method (ADM) [14],[15],[16], Homotopy Perturbation method (HPM) [17],[18],[19], Variational Iteration method (VIM) [20],[21],[22] and other methods have been proposed to solve linear and nonlinear problems. Another analytical approach that can be applied to solve many types of nonlinear fractional differential equation is Homotopy Analysis method (HAM) [23],[24],[25],[26],[27],[28], put forward by Liao in 1992. Very recently, the Homotopy Analysis method is combined with Elzaki transform to produce a highly effective technique called Homotopy Analysis Elzaki Transform method (HAETM) for handling some differential equations [29],[30].

In this paper, we consider the nonlinear fractional order biological population model in the form:

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + f(u), \quad 0 < \alpha \leq 1 \tag{1.1}$$

with the initial condition

$$u(x, y, 0) = f_0(x, y) \tag{1.2}$$

where u denotes the population density, and $f(u)$ represents the population supply due to birth and deaths, α is a parameter describing the order of the fractional derivative. The derivatives in (1.1) are understood in the Caputo sense. In this paper, further we apply the homotopy analysis Elzaki transform method (HAETM) to solve the fractional biological population models.

The advantage of this method is its capability of combing two powerful methods for obtaining exact and approximate analytical solutions for nonlinear equations. The fact that the HAETM solves nonlinear problems without using Adomian's polynomials and He's polynomials is a clear advantage over the Adomian decomposition method (ADM) and He's perturbation transform method (HPTM) [31],[32]. The plan of our paper is as follows: Brief definition of fractional calculus are given in Section 2. Some theorems of Elzaki transform are given in Section 3. The homotopy analysis Elzaki transform method is presented in Section 4. In Section 5, three numerical examples are solved to illustrate the applicability of the considered method. Conclusions are presented in Section 6.

2 BASIC DEFINITIONS

In this section, we mention the following basic definitions of fractional calculus.

Definition 2.1. A real function $f(x)$, $x > 0$ is said to be in the space $C_{\mu}, \mu \in R$, if there exist a real number $p(> \mu)$ such that $f(x) = x^p f_1(x)$, where $f_1(x) \in C[0, +\infty)$, and it is said to be in the space C_{μ}^m if $f^{(m)} \in C_{\mu}, m \in N \cup \{0\}$.

Definition 2.2. The Riemann-Liouville fractional integral operator of order $\alpha > 0$, of a function $f(x) \in C_{\mu}, \mu \geq -1$ is defined as

$$J^\alpha f(x) = \frac{1}{\Gamma(\alpha)} \int_0^x (x-t)^{\alpha-1} f(t) dt, \quad \alpha > 0, x > 0 \tag{2.1}$$

$$J^0 f(x) = f(x) \tag{2.2}$$

For the Riemann-Liouville fractional integral we have:

$$J^\alpha J^\beta f(x) = J^{\alpha+\beta} f(x) \tag{2.3}$$

$$J^\alpha J^\beta f(x) = J^\beta J^\alpha f(x) \tag{2.4}$$

$$J^\alpha x^\gamma = \frac{\Gamma(\gamma + 1)}{\Gamma(\alpha + \gamma + 1)} x^{\alpha+\gamma} \tag{2.5}$$

For $f \in C_\mu$, $\mu \geq -1$, $\alpha, \beta \geq 0$ and $\gamma > -1$.

Definition 2.3. The fractional derivatives of $f(x)$ in the Caputo sense is defined:

$$D_t^\alpha f(x) = J^{m-\alpha} D^m f(x) = \frac{1}{\Gamma(m-\alpha)} \int_0^x (x-t)^{m-\alpha-1} f^{(m)}(t) dt \tag{2.6}$$

For $m-1 < \alpha \leq m$, $m \in N$, $\mu \geq -1$, and $f \in C_\mu^m$.

Definition 2.4. The Laplace transform of Caputo derivative is:

$$L(D_t^\alpha f(x)) = s^\alpha L[f(x)] - \sum_{k=0}^{m-1} s^{\alpha-k-1} f^{(k)}(0^+), \quad m-1 < \alpha \leq m. \tag{2.7}$$

3 ELZAKI TRANSFORM

Recently, Tarig Elzaki introduced a new integral transform, named Elzaki transform [30], and further applied it to the solution of ordinary and partial differential equations. The Elzaki transform is defined over the set of functions

$$A = \{f(t) : \exists M, k_1, k_2 > 0, |f(t)| < M e^{\frac{|t|}{k_j}}, t \in (-1)^j \times [0, +\infty)\} \tag{3.1}$$

by the following formula

$$T(v) = E[f(t)] = v \int_0^{+\infty} e^{-\frac{t}{v}} f(t) dt, \quad v \in [k_1, k_2]. \tag{3.2}$$

Theorem 3.1. If $f(t) = t^\alpha$,

$$E[t^\alpha] = v \int_0^{+\infty} e^{-\frac{t}{v}} t^\alpha dt = v^{\alpha+2} \Gamma(\alpha + 1).$$

Theorem 3.2. Elzaki transform on the Riemann-Liouville fractional integral operator of a function $f(t)$:

$$E[J^\alpha f(t)] = v^{\alpha+1} T(v).$$

Theorem 3.3. Elzaki transform on the Caputo fractional derivative of $f(t)$:

$$E[D_x^{n\alpha} u(x, t)] = \frac{T(v)}{v^{n\alpha}} - \sum_{k=0}^{n-1} v^{2-n\alpha} u(0, t), \quad n-1 < n\alpha \leq n$$

4 HAETM FOR GENERALIZED BIOLOGICAL POPULATION MODEL

To illustrate the basic idea of this method, let us consider the generalized biological population model:

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + ku^a(1 - ru^b) \quad (4.1)$$

$t > 0, x, y \in R, 0 < \alpha \leq 1, k, a, r, b$ are real numbers, with the initial condition

$$u(x, y, 0) = f_0(x, y) \quad (4.2)$$

Now taking the Elzaki transform of both side of (4.1), we get

$$\frac{E[u(x, y, t)]}{v^\alpha} - v^{2-\alpha}u(x, y, 0) - E[\frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + ku^a(1 - ru^b)] = 0 \quad (4.3)$$

We define the nonlinear operator

$$N[\phi(x, y, t; q)] = E[\phi(x, y, t; q)] - v^2u(x, y, 0) - v^\alpha E[\frac{\partial^2}{\partial x^2}(\phi^2(x, y, t; q)) + \frac{\partial^2}{\partial y^2}(\phi^2(x, y, t; q)) + k\phi^a(x, y, t; q)(1 - r)\phi^b(x, y, t; q)] \quad (4.4)$$

where $q \in [0, 1]$ and $\phi(x, y, t; q)$ is a real function of x, y, t, q .

The zero-order deformation equation of the (4.3) has the form

$$(1 - q)E[\phi(x, y, t; q) - u_0(x, y, t)] = \hbar q H(t) N[\phi(x, y, t; q)] \quad (4.5)$$

where E is the Elzaki transform, $q \in [0, 1]$ is the embedding parameter, $H(t)$ denotes a nonzero auxiliary function, $\hbar \neq 0$ is an auxiliary parameter, $u_0(x, y, t)$ is an initial guess of $u(x, y, t)$ and $\phi(x, y, t; q)$ is an unknown function. Obviously, when the parameter $q = 0$ and $q = 1$, it holds

$$\phi(x, y, t; 0) = u_0(x, y, t), \quad \phi(x, y, t; 1) = u(x, y, t) \quad (4.6)$$

respectively. Thus, as q increase from 0 to 1, the solution $\phi(x, y, t; q)$ varies from the initial guess $u_0(x, y, t)$ to the solution $u(x, y, t)$. Expanding $\phi(x, y, t; q)$ in Taylor series with respect to q , we have

$$\phi(x, y, t; q) = u_0(x, y, t) + \sum_{m=1}^{\infty} u_m(x, y, t) q^m \quad (4.7)$$

where

$$u_m(x, y, t) = \frac{1}{m!} \frac{\partial^m \phi(x, y, t; q)}{\partial q^m} \Big|_{q=0} \quad (4.8)$$

Differentiating the zero-order deformation equation (4.5) m times with respect to q and then dividing by $m!$ and finally setting $q = 0$ we get the following m^{th} -order deformation equation:

$$E[u_m(x, y, t) - \chi_m u_{m-1}(x, y, t)] = \hbar q H(t) R_m(\vec{u}_{m-1}(x, y, t)). \quad (4.9)$$

Define the vectors

$$\vec{u}_m = \{u_0(x, y, t), u_1(x, y, t), u_2(x, y, t), \dots, u_m(x, y, t)\} \quad (4.10)$$

Applying the inverse Elzaiki transform on (4.9), we have

$$u_m(x, y, t) = \chi_m u_{m-1}(x, y, t) + \hbar q E^{-1}[H(t) R_m(\vec{u}_{m-1}(x, y, t))] \quad (4.11)$$

where

$$R_m(\vec{u}_{m-1}(x, y, t)) = \frac{1}{(m-1)!} \frac{\partial^{m-1} N[\phi(x, y, t; q)]}{\partial q^{m-1}} \Big|_{q=0} \quad (4.12)$$

and

$$\chi_m = \begin{cases} 0 & m \leq 1 \\ 1 & m > 1 \end{cases}$$

If the auxiliary linear operator, the initial guess, the auxiliary parameter h , and the auxiliary function are properly chosen, the series (4.7) converges at $q = 1$, then we have

$$u(x, y, t) = u_0(x, y, t) + \sum_{m=1}^{\infty} u_m(x, y, t) \quad (4.13)$$

which must be one of the solutions of the original nonlinear equation.

5 NUMERICAL RESULTS

In this section, we use homotopy analysis Elzaki transform method to solve nonlinear fractional biological population equation.

Example 5.1. Consider the equation (4.1) with $k = 1, a = 1, r = 0$, we have the following fractional biological population equation:

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + u \quad (5.1)$$

with the initial condition

$$u(x, y, 0) = \sqrt{\sin x \cdot \sinh y} \quad (5.2)$$

Applying the Elzaki transform to (5.1) and using (5.2), we have

$$\frac{E[u(x, y, t)]}{v^\alpha} - v^{2-\alpha} \sqrt{\sin x \cdot \sinh y} - E\left[\frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + u\right] = 0 \quad (5.3)$$

The nonlinear operator is

$$N[\phi(x, y, t; q)] = E[\phi(x, y, t; q)] - v^2 \sqrt{\sin x \cdot \sinh y} - v^\alpha E\left[\frac{\partial^2}{\partial x^2}(\phi^2(x, y, t; q)) + \frac{\partial^2}{\partial y^2}(\phi^2(x, y, t; q)) + \phi(x, y, t; q)\right] \quad (5.4)$$

and thus

$$R_m(\vec{u}_{m-1}) = E[u_{m-1}] - (1 - \chi_m)v^2 \sqrt{\sin x \cdot \sinh y} - v^\alpha E\left[\frac{\partial^2}{\partial x^2}\left(\sum_{r=0}^{m-1} u_r u_{m-1-r}\right) + \frac{\partial^2}{\partial y^2}\left(\sum_{r=0}^{m-1} u_r u_{m-1-r}\right) + u_{m-1}\right] \quad (5.5)$$

The m^{th} -order deformation equation is given by

$$E[u_m(x, y, t) - \chi_m u_{m-1}(x, y, t)] = \hbar R_m(\vec{u}_{m-1}(x, y, t)) \quad (5.6)$$

Applying the inverse Elzaki transform, we have

$$u_m(x, y, t) = \chi_m u_{m-1}(x, y, t) + \hbar E^{-1}[R_m(\vec{u}_{m-1}(x, y, t))] \quad (5.7)$$

Solving the above equation (5.7), for $m = 1, 2, 3, \dots$, we get

$$\begin{aligned}
 u_1(x, y, t) &= -\hbar \sqrt{\sin x \cdot \sinh y} \frac{t^\alpha}{\Gamma(\alpha + 1)} \\
 u_2(x, y, t) &= -\hbar(1 + \hbar) \sqrt{\sin x \cdot \sinh y} \frac{t^\alpha}{\Gamma(\alpha + 1)} \\
 &\quad + \hbar^2 \sqrt{\sin x \cdot \sinh y} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} \\
 u_3(x, y, t) &= -\hbar(1 + \hbar)^2 \sqrt{\sin x \cdot \sinh y} \frac{t^\alpha}{\Gamma(\alpha + 1)} \\
 &\quad + 2\hbar^2(1 + \hbar) \sqrt{\sin x \cdot \sinh y} \frac{t^{2\alpha}}{\Gamma(2\alpha + 1)} \\
 &\quad - \hbar^3 \sqrt{\sin x \cdot \sinh y} \frac{t^{3\alpha}}{\Gamma(3\alpha + 1)}
 \end{aligned} \tag{5.8}$$

and so on.

Substituting u_1, u_2, u_3, \dots into equation (4.13) gives the solution in series form, by $\hbar = -1$, we have

$$u(x, y, t) = \sqrt{\sin x \cdot \sinh y} \left[1 + \frac{t^\alpha}{\Gamma(1 + \alpha)} + \frac{t^{2\alpha}}{\Gamma(1 + 2\alpha)} + \frac{t^{3\alpha}}{\Gamma(1 + 3\alpha)} + \dots \right] \tag{5.9}$$

If we put $\alpha = 1$, we have

$$u(x, y, t) = \sqrt{\sin x \cdot \sinh y} \left[1 + t + \frac{t^2}{2!} + \frac{t^3}{3!} + \dots \right] \tag{5.10}$$

$$u(x, y, t) = \sqrt{\sin x \cdot \sinh y} \cdot e^t \tag{5.11}$$

which is an exact solution.

Example 5.2. Consider the equation (4.1) with $a = 1, r = 0$, we have the following fractional biological population equation:

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + ku \tag{5.12}$$

with the initial condition

$$u(x, y, 0) = \sqrt{xy} \tag{5.13}$$

Applying the Elzaki transform to (5.12) and using (5.13), we have

$$\frac{E[u(x, y, t)]}{v^\alpha} - v^{2-\alpha} \sqrt{xy} - E\left[\frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + ku\right] = 0 \tag{5.14}$$

The nonlinear operator is

$$\begin{aligned}
 N[\phi(x, y, t; q)] &= E[\phi(x, y, t; q)] - v^2 \sqrt{xy} - v^\alpha E\left[\frac{\partial^2}{\partial x^2}(\phi^2(x, y, t; q))\right. \\
 &\quad \left. + \frac{\partial^2}{\partial y^2}(\phi^2(x, y, t; q)) + k\phi(x, y, t; q)\right]
 \end{aligned} \tag{5.15}$$

and thus

$$\begin{aligned}
 R_m(\vec{u}_{m-1}) &= E[u_{m-1}] - (1 - \chi_m) v^2 \sqrt{xy} \\
 &\quad - v^\alpha E\left[\frac{\partial^2}{\partial x^2} \left(\sum_{r=0}^{m-1} u_r u_{m-1-r}\right) + \frac{\partial^2}{\partial y^2} \left(\sum_{r=0}^{m-1} u_r u_{m-1-r}\right) + k u_{m-1}\right]
 \end{aligned} \tag{5.16}$$

The m^{th} -order deformation equation is given by

$$E[u_m(x, y, t) - \chi_m u_{m-1}(x, y, t)] = \hbar R_m(\vec{u}_{m-1}(x, y, t)) \quad (5.17)$$

Applying the inverse Elzaki transform, we have

$$u_m(x, y, t) = \chi_m u_{m-1}(x, y, t) + \hbar E^{-1}[R_m(\vec{u}_{m-1}(x, y, t))] \quad (5.18)$$

Solving the above equation (5.18), for $m = 1, 2, 3, \dots$, we get

$$\begin{aligned} u_1(x, y, t) &= -k\hbar\sqrt{xy}\frac{t^\alpha}{\Gamma(\alpha+1)} \\ u_2(x, y, t) &= -k\hbar(1+\hbar)\sqrt{xy}\frac{t^\alpha}{\Gamma(\alpha+1)} \\ &\quad + k^2\hbar^2\sqrt{xy}\frac{t^{2\alpha}}{\Gamma(2\alpha+1)} \\ u_3(x, y, t) &= -k\hbar(1+\hbar)^2\sqrt{xy}\frac{t^\alpha}{\Gamma(\alpha+1)} \\ &\quad + 2k^2\hbar^2(1+\hbar)\sqrt{xy}\frac{t^{2\alpha}}{\Gamma(2\alpha+1)} \\ &\quad - k^3\hbar^3\sqrt{xy}\frac{t^{3\alpha}}{\Gamma(3\alpha+1)} \end{aligned} \quad (5.19)$$

and so on.

Substituting u_1, u_2, u_3, \dots into equation (4.13) gives the solution in series form, by $\hbar = -1$, we have

$$u(x, y, t) = \sqrt{xy}\left[1 + k\frac{t^\alpha}{\Gamma(1+\alpha)} + k^2\frac{t^{2\alpha}}{\Gamma(1+2\alpha)} + k^3\frac{t^{3\alpha}}{\Gamma(1+3\alpha)} + \dots\right] \quad (5.20)$$

If we put $\alpha = 1$, we have

$$u(x, y, t) = \sqrt{xy}\left[1 + kt + \frac{k^2t^2}{2!} + \frac{k^3t^3}{3!} + \dots\right] \quad (5.21)$$

$$u(x, y, t) = \sqrt{xy} \cdot e^{kt} \quad (5.22)$$

which is an exact solution.

The evolution results for the exact solution (5.22) and the approximate solution (5.20), for the case $\alpha = 1$, are shown in Figure 1. It can be seen from Figure 1 that the solution obtained by the HAETM is nearly identical with the exact solution.




Figure 1: Exact and numerical solutions of Example 2

Example 5.3. Consider the equation (4.1) with $a = -1, b = 1$, we have the following fractional biological population equation:

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + ku^{-1} - kr \quad (5.23)$$

with the initial condition

$$u(x, y, 0) = \sqrt{\frac{kr}{4}x^2 + \frac{kr}{4}y^2 + y + 5} \quad (5.24)$$

Applying the Elzaki transform to (5.23) and using (5.24), we have

$$\frac{E[u(x, y, t)]}{v^\alpha} - v^{2-\alpha} \sqrt{\frac{kr}{4}x^2 + \frac{kr}{4}y^2 + y + 5} - E\left[\frac{\partial^2 u^2}{\partial x^2} + \frac{\partial^2 u^2}{\partial y^2} + ku^{-1} - kr\right] = 0 \quad (5.25)$$

The nonlinear operator is

$$\begin{aligned} N[\phi(x, y, t; q)] = & E[\phi(x, y, t; q)] - v^2 \sqrt{\frac{kr}{4}x^2 + \frac{kr}{4}y^2 + y + 5} - v^\alpha E\left[\frac{\partial^2}{\partial x^2}(\phi^2(x, y, t; q))\right. \\ & \left. + \frac{\partial^2}{\partial y^2}(\phi^2(x, y, t; q)) + k\phi^{-1}(x, y, t; q) - kr\right] \end{aligned} \quad (5.26)$$

and thus

$$\begin{aligned} R_m(\vec{u}_{m-1}) = & E[u_{m-1}] - (1 - \chi_m)v^2 \sqrt{\frac{kr}{4}x^2 + \frac{kr}{4}y^2 + y + 5} \\ & - v^\alpha E\left[\frac{\partial^2}{\partial x^2}\left(\sum_{r=0}^{m-1} u_r u_{m-1-r}\right) + \frac{\partial^2}{\partial y^2}\left(\sum_{r=0}^{m-1} u_r u_{m-1-r}\right) + \frac{k}{u_{m-1}} - kr\right] \end{aligned} \quad (5.27)$$

The m^{th} -order deformation equation is given by

$$E[u_m(x, y, t) - \chi_m u_{m-1}(x, y, t)] = \hbar R_m(\vec{u}_{m-1}(x, y, t)) \quad (5.28)$$

Applying the inverse Elzaki transform, we have

$$u_m(x, y, t) = \chi_m u_{m-1}(x, y, t) + \hbar E^{-1}[R_m(\vec{u}_{m-1}(x, y, t))] \quad (5.29)$$

Solving the above equation (5.29), for $m = 1, 2, 3, \dots$, we get

$$\begin{aligned} u_1(x, y, t) &= \frac{-k\hbar t^\alpha}{\Gamma(1 + \alpha)\sqrt{\frac{kr}{4}x^2 + \frac{kr}{4}y^2 + y + 5}} \\ u_2(x, y, t) &= \frac{-2k^2\hbar^2 t^{2\alpha}}{\Gamma(1 + 2\alpha)(\sqrt{\frac{kr}{4}x^2 + \frac{kr}{4}y^2 + y + 5})^3} \\ u_3(x, y, t) &= \frac{-3k^3\hbar^3 t^{3\alpha}}{\Gamma(1 + 3\alpha)(\sqrt{\frac{kr}{4}x^2 + \frac{kr}{4}y^2 + y + 5})^5} \end{aligned} \quad (5.30)$$

and so on.

Substituting u_1, u_2, u_3, \dots into equation (4.13) gives the solution in series form, by $\hbar = -1$, we have

$$u(x, y, t) = u_0 + \frac{kt^\alpha}{u_0} \left[\frac{1}{\Gamma(1 + \alpha)} + \frac{2}{\Gamma(1 + 2\alpha)} \frac{-kt^\alpha}{u_0^2} + \frac{3}{\Gamma(1 + 3\alpha)} \frac{k^2 t^{2\alpha}}{u_0^4} + \dots \right] \quad (5.31)$$

namely

$$u(x, y, t) = u_0 + \frac{kt^\alpha}{u_0} \sum_{n=0}^{\infty} \frac{n+1}{\Gamma(1 + (n+1)\alpha)} \left(\frac{-kt^\alpha}{u_0^2}\right)^n \quad (5.32)$$

If we put $\alpha = 1$, we have

$$u(x, y, t) = u_0 + \frac{kt}{u_0} e^{\frac{-kt}{u_0}} \quad (5.33)$$

which is an exact solution.

The evolution results for the exact solution (5.33) and the approximate solution (5.31), for the special case $\alpha = 1, k = 0.05, r = 45, t = 10$ are shown in Figure 2. It can be seen from Figure 2 that the solution obtained by the HAETM is nearly identical with the exact solution.




Figure 2: Exact and numerical solutions of Example 3

6 CONCLUSIONS

In this paper, the homotopy analysis Elzaki transform method has been successfully applied to derive approximate solutions of the fractional

order biological population equations subject to some initial conditions. The results obtained by this method agree well with the results obtained by ADM, VIM, HPM. The reliability of HAETM and reduction in computations give this method

a wider applicability. Finally, we conclude that the HAETM is very powerful and efficient in finding analytical and numerical solutions for wider classes of linear and nonlinear fractional differential equations. It provides us with a simple way to adjust and control the convergence region of solution series by choosing proper values for auxiliary parameter \hbar and auxiliary function $H(t)$. The corresponding solutions and 3D graphs are obtained according to the recurrence relation using Mathematica.

ACKNOWLEDGEMENT

The authors are grateful to the handling editor and the anonymous referees for their careful reading and critical comments which have helped them to improve the presentation of this work significantly.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

References

- [1] Young GO. Definition of physical consistent damping laws with fractional derivatives. *Z. Angew. Math. Mech.* 1995;20:623-635.
- [2] He Jihuan. Some applications of nonlinear fractional differential equations and their approximations. *Bull.Sci.Technol.* 1999;15(2):86-90.
- [3] He Jihuan. Approximate analytic solution for seepage flow with fractional derivatives in porous media. *Comput. Methods Appl. Mech. Eng.* 1998;167(1-2):57-68.
- [4] Mainardi F, Luchko Y, Pagnini G. The fundamental solution of the space-time fractional diffusion equation. *Fractional Calculus and Applied Analysis.* 2001;4(2):153-192.
- [5] Radi SZ, El-Sayed AMA, Arafa AMA. On the solutions of time-fractional reaction-diffusion equations. *Communications in Nonlinear Science and Numerical Simulation.* 2010;15(2):3847-3854.
- [6] Yildirim A. He's homotopy perturbation method for solving the space- and time-fractional telegraph equations. *International Journal of Computer Mathematics.* 2010;87(13):2998-3006.
- [7] Debnath L. Fractional integrals and fractional differential equations in fluid mechanics. *Frac. Calc. Appl. Anal.* 2003;6(2):119-155.
- [8] Muslih SI, Baleanu D, Rabei E. Hamiltonian formulation of classical fields within Riemann-Liouville fractional derivatives. *Physica Scripta.* 2006;73(5):436-438.
- [9] Baleanu D. About fractional quantization and fractional variational principles. *Communications in Nonlinear Science and Numerical Simulation.* 2009;14(6):2520-2523.
- [10] Herzallah MAE, El-Sayed AMA, Baleanu D. On the fractional-order diffusion-wave process. *Romanian Journal of Physics.* 2010;55(3-4):274-284.
- [11] Al-Rabtah A, Erturk VS, Momani S. Solutions of a fractional oscillator by using differential transform method. *Computers and Mathematics with Applications.* 2010;59(3):1356-1362.
- [12] Zhou JK. *Differential transformation and its applications for electrical circuits.* Huazhong University Press, Wuhan, China; 1986.
- [13] Pukhov GE. Computational structure for solving differential equations by Taylor transformation. *Cybernetics and System Analysis.* 1978;14(3):383-390.
- [14] Adomian G. *Solving frontier problems of physics: The decomposition method.* Kluwer Acad. Publ. Boston; 1994.
- [15] Jafari H, Daftardar-Gejji V. Solving linear and nonlinear fractional diffusion and wave equations by Adomian-decomposition. *Appl. Math. Comput.* 2006;180(2):488-497.
- [16] Momani S, Al-Khaled A. Numerical solutions for system of fractional differential equations by decomposition method. *Appl. Math. Comput.* 2005;196:644-665.
- [17] He Jihuan. Homotopy perturbation technique. *Computer Methods in Applied Mechanics and Engineering.* 1999;178(3-4):257-262.

- [18] He Jihuan. New interpretation of homotopy perturbation method. *Int. J. Mod. Phys. B.* 2006;20:2561-2568.
- [19] Ganji DD. The applications of He's homotopy perturbation method to nonlinear equation arising in heat transfer. *Physics Letters A.* 2006;335(4-5):337-341.
- [20] He Jihuan. Variational iteration method—a kind of nonlinear analytical technique: Some examples. *International Journal of Nonlinear Mechanics.* 1999;34(4):699-708.
- [21] He Jihuan, Wu XH. Variational iteration method: New development and applications. *Computers and Mathematics with Applications.* 2007;54(7-8):881-894.
- [22] He Jihuan, Wu GC, Austin F. The variational iteration method which should be followed. *Nonlinear Science Letters A-Mathematics, Physics and Mechanics;* 2010.
- [23] Liao Shijun. The proposed homotopy analysis technique for the solution of nonlinear problems. PhD Thesis, Shanghai Jiao Tong University; 1992.
- [24] Liao Shijun, Sherif SA. Beyond Perturbation: Introduction to homotopy analysis method. *Applied Mechanics Reviews.* 2004;57(5):25.
- [25] Liao Shijun, Sherif SA. On the homotopy analysis method for nonlinear problems. *Applied Mathematics and Computation.* 2004;147(2):499-513.
- [26] Liao Shijun, Sherif SA. A new branch of solutions of boundary-layer flows over an impermeable stretched plate. *Int. J. Heat Mass Transfer.* 2005;48(12):2529-2539.
- [27] Wang Qi. Application of homotopy analysis method to solve Relativistic Toda Lattice System. *Communication in Theoretical Physics.* 2010;53(6):1111-1116.
- [28] Shidfar A, Molabahrami A. A weighted algorithm based on the homotopy analysis method: Application to inverse heat conduction problems. *Communications in Nonlinear Science and Numerical Simulation.* 2010;15(10):2908-2915.
- [29] Elzaki TM, Elzaki SM. Application of new transform "Elzaki Transform" to partial differential equations. *Global Journal of Pure and Applied Mathematics.* 2011;7(1):65-70.
- [30] Elzaki TM. The new integral transform "ELzaki Transform". *Global Journal of Pure and Applied Mathematics.* 2011;7(1):57-64.
- [31] Khan Y, Wu Q. Homotopy perturbation transform method for nonlinear equations using He's polynomials. *Computers and Mathematics with Applications.* 2011;61(8):1963-1967.
- [32] Ghorbani A. Beyond adomian's polynomials: He polynomials. *Chaos Solitons and Fractals.* 2009;39(3):1486-1492.

© 2018 Xu and Lu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/46155>