



Network-based Toxicogenomics Approach to Unraveling the Influence of Lead and Cadmium Mixture on the Development and Progression of Hypertension

Oluwatoyin Folake Olukunle^{a*} and Victor Omoboyede^b

^a *Department of Biotechnology, School of Life Sciences (SLS), Federal University of Technology Akure, P.M.B 704, Akure, Nigeria.*

^b *Department of Biochemistry, School of Life Sciences (SLS), Federal University of Technology Akure, P.M.B 704, Akure, Nigeria.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/BJI/2023/v27i2676

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95613>

Original Research Article

Received: 06/01/2023

Accepted: 25/02/2023

Published: 05/05/2023

ABSTRACT

Heavy metals, generally characterized by high densities and atomic weights, are ubiquitous in the environment and are a public health concern due to the several health issues they pose to humans. Of all heavy metals, lead and cadmium among others are known to be capable of inducing multiple health effects even at a low rate of exposure. Hypertension (HYP), a major cause of death and a risk factor for other cardiovascular diseases, is known to be caused by both lead and cadmium. While the mechanism underlying the development of HYP induced by independent exposure to

*Corresponding author: E-mail: ofolukunle@gmail.com;

lead and cadmium has been well studied, the mechanism underlying the induction and progression of HYP upon lead and cadmium co-exposure remains mildly explored. Hence, this study aimed to elucidate the mechanism using a toxicogenomic approach. The set of genes affected by both heavy metals was identified using the comparative toxicogenomics database (CTD) while HYP targets were retrieved from the Gene Cards database. The shared genes between the heavy metals and the disease were identified and subjected to further analysis. The results of our analysis revealed the signaling pathways that are dysregulated by lead and cadmium co-exposure while oxidative stress, inflammation, and endothelial dysfunction were revealed as processes pertinent to the induction and progression of HYP by lead and cadmium co-exposure. Biomarkers that could be used for prognosis evaluation were also identified. Ultimately, this study supports and advances the growing body of finding on the roles played by lead and cadmium co-exposure in inducing HYP.

Keywords: Hypertension (HYP); heavy metals; lead; cadmium; signaling pathways.

1. INTRODUCTION

Over the years, the risk of exposure to various toxic substances has tremendously increased due to various industrial, geogenic, and anthropogenic activities. These toxic substances include heavy metals such as Arsenic (As), Lead (Pb), Cadmium (Cd), and Mercury (Hg). Heavy metals are ubiquitous and are widely distributed in the environment due to technological applications among other factors [1]. Based on biological utility, heavy metals are classified into essential and non-essential. The essential heavy metals are required by living organisms in minimal concentrations and they play important biochemical and physiological functions such as serving as cofactors for several enzymes. Conversely, non-essential heavy metals are known to possess no known beneficial effect on living organisms [1,2]. Both essential and non-essential heavy metals are known to pose dose- and duration-dependent detrimental effects to human health; this is due to their ability to interact with cellular organelles and components such as the mitochondria, deoxyribonucleic acid (DNA), and proteins. Hence, this results in the disruption of normal biological and physiological processes including DNA damage, cytotoxicity, apoptosis, and carcinogenesis [3]. Notably, heavy metal toxicity has been implicated in the etiology of cardiovascular, neurological, autoimmune, and respiratory diseases [4].

Of all heavy metals, Pb and Cd pose a great challenge to human health due to their concentrations in the environment and non-biodegradability [5]. Interestingly, Pb and Cd are ranked second and seventh respectively on the priority list of dangerous substances by the United States Agency for Toxic Substances and Disease Registry (ATSDR) [6]. Pb and Cd, which exist in water, soil, and air, are transferred to the

human body via ingestion and inhalation of contaminated food and air. Upon entrance to the human body, they are distributed by the red blood cells and proteins which cause various damages including blood pressure changes that result in HYP [5]. Noteworthy, while some areas possess low individual concentrations of heavy metals which correspond to lower chances of inducing toxicity, a combined exposure to a mixture of heavy metals could still result in detrimental effects [7]. For example, Zhouab et al. [8] demonstrated that Pb, Cd, arsenic, and mercury co-exposure led to higher neurotoxic effects compared to the effects induced by co-exposure to three or fewer of the same metals. Similarly, Javorac et al. [9] reported that oxidative status induced by exposure of rats to metal mixtures was absent during treatment with the individual metals.

Several studies have investigated the roles and mechanisms of Pb and Cd in inducing blood pressure changes and HYP. Chen *et al.* reported that blood Cd level was associated with elevated blood pressure. They also reported that an increase in blood Cd level was concomitant with an increase in the prevalence of HYP in the studied population [10]. Furthermore, Eum et al. [11] reported that Cd toxicity results in HYP via the inhibition of endothelial nitric oxide synthase and the suppression of vascular relaxation induced by acetylcholine. They also reported that Cd mediates the production of cytokines and induces endothelial damage. Similarly, Robles et al. [12] reported that Pb treatment was associated with increased generation of reactive oxygen species (ROS) which might have led to the inactivation of nitric oxide (NO) in Pb-induced hypertensive animals. This is corroborated by a study by Alghasham et al. [13], in which they reported that Pb decreased the plasma levels of nitric oxide, which functions as a vasodilator, in

hypertensive patients. They also reported a positive correlation between blood pressure and blood Pb levels. Studies have also investigated the effects of Pb and Cd co-exposure. Andjelkovic et al. [5] reported that a mixed treatment of wistar rats with Pb and Cd resulted in a high plasma advanced oxidation protein product level (AOPP), signifying oxidative damages. They also reported that the co-exposure of Cd and Pb resulted in a higher negative effect in plasma compared to the individual metals. While numerous studies have been conducted to demystify the roles of Pb and Cd in the etiology of cardiovascular diseases, the mechanism by which the co-exposure to both metals induces HYP in humans remains a mystery [13]. Hence, this necessitates further attempts aimed at unraveling their mechanism as etiological agents of HYP.

Advances in bioinformatics have powered the breaking of limitations of conventional toxicology through the ability to investigate the impact of toxicants at the molecular level. Hence, providing biomarkers of toxicants for further *In vitro* and *In vivo* evaluation. Using a toxicogenomic approach, this study aims to unravel the molecular mechanism via which Pb and Cd co-exposure induce and aggravate HYP.

2. MATERIALS AND METHODS

2.1 Retrieval of Pb and Cd Gene Targets

The genes that are affected by Pb and Cd toxicity were retrieved from the comparative toxicogenomics database (CTD, <http://ctdbase.org/>) [14–16] (accessed May 8, 2022) using the keywords “lead” and “cadmium” respectively.

2.2 Retrieval of HYP Gene Targets

The gene targets of HYP were retrieved from the GeneCards database (<https://www.genecards.org/>) [17].

2.3 Identification of Shared Targets

The My Venn analysis tool of the CTD database (<http://ctdbase.org/tools/myVenn.go>) was utilized to identify the gene targets of lead and cadmium relevant to HYP.

2.4 Construction of Protein-protein Interaction (PPI) Network

To explore the interaction between the shared targets of lead, cadmium, and HYP, the string

database (<https://string-db.org/>) [18] was employed to construct the PPI network of the shared targets, with *Homo sapiens* selected as the organism of interest and a medium confidence score of 0.4, as the minimum required interaction score.

2.5 Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) Pathway Analysis

To access the biological roles and signaling pathways that are disrupted due to co-exposure to both Pb and Cd, the ShinyGo v 0.76 web server (<https://bioinformatics.sdstate.edu/go/>) [19] was employed to perform the GO and KEGG pathways analysis of the common targets. Furthermore, the enrichment plots of the significant GO and pathways of the target genes were generated using the graphing tool on the same web server.

3. RESULTS

3.1 Retrieval of Pb and Cd Gene Targets

Following extensive data mining activity on the CTD, the gene targets of lead and cadmium were curated. Interestingly, 3, 175 genes and 3, 124 genes were retrieved as Cd and Pb targets respectively.

3.2 Retrieval of HYP Gene Targets

Using the keyword “HYP” on the Gene Cards database, 9, 408 unique genes that have been implicated in the development of HYP were retrieved. Subsequently, the RNA genes and other non-genes were filtered out. This resulted in the identification of 8, 727 unique genes with standard IDs.

3.3 Identification of Shared Targets

Pb and Cd genes that are disrupted for the induction of HYP following co-exposure were identified using the My Venn tool of CTD. The gene IDs of HYP, lead, and cadmium was uploaded to the server. The identified 512 shared targets (Fig. 1) were considered the genes that are disrupted for the induction of HYP following Pb and Cd co-exposure (Supplementary file 1).

3.4 Construction of PPI Network

To understand the relationship between the shared genes in the biological milieu, StringDB

was employed to generate a PPI network of the 512 common targets. Interestingly, only 496 targets showed interactions based on the PPI network analysis. Visualization and further analysis of the generated PPI network using Cytoscape v 3.9.1 showed the network possessed 496 nodes and 10,985 edges (Fig. 2).

Subsequently, the CytoHubba plug-in of Cytoscape was utilized to derive the top 10 targets of the network based on the degree of interactions (Fig. 3). These targets can be considered the essential targets via which co-exposure to lead and cadmium disrupt normal biological and physiological activities of the body.




Fig. 1. A Venn diagram depicting the number of genes that were retrieved for Pb, Cd, HYP, and the number of interacting genes




Fig. 2. PPI network of the 512 shared targets constructed using string

3.5 GO and KEGG Pathway Analysis

Enrichment analysis of the common targets was performed using the ShinyGo v 0.76 webserver. The GO analysis revealed the various functions that are played by the genes at the molecular level. Interestingly, the genes were enriched in biological processes such as response to lead ion, response to cadmium ion, response to cell death, response to oxygen-containing compound, and response to oxidative stress. Similarly, the molecular functions of the genes include oxidoreductase activity, antioxidant activity, ubiquitin-like protein ligase binding, hormone activity, and protein kinase activity. Furthermore, the targets were also found to be cellular components of the mitochondrion, blood microparticle, cell surface, lysosome, and platelet alpha granule. Based on the FDR cut-off of 0.05, the top 10 results of the GO analysis were further selected for visualization (Fig. 4).

Identification of the potential signaling pathways disrupted as a result of the co-exposure revealed

the HIF-1 signaling pathway, chemical carcinogenesis, PI3K-Akt signaling pathway. Mineral absorption pathway, and P53 signaling pathway as the potential pathways in which the shared are enriched. The top 20 pathways were identified and visualized based on an FDR cut-off of 0.05 (Fig. 5).

4. DISCUSSION

Heavy metals are highly notorious for their ability to induce numerous forms of toxicity-associated burdens in humans, especially at elevated levels of exposure. While several experimental and computational studies have accessed the effects and mode of action of many heavy metals including lead and cadmium on the biological system, studies on the mechanism and the ability of co-exposure to these metals to result in some diseases and trigger an additive or synergistic effect have been mildly explored. To this end, this study was aimed at unraveling the molecular mechanism via which Pb and Cd co-exposure results in the development of HYP.




Fig. 3. The top ten targets of the PPI network selected based on the degree of interactions using CytoHubba plug-in of cytoscape




Fig. 4. The Gene ontology functional annotation analysis of the targets of Pb and Cd in HYP. A. Biological process in which the targets are enriched. (b) Molecular function of the targets. (c) Cellular components of the targets




Fig. 5. The top 20 KEGG pathways in which the targets are enriched

The set of genes whose biological activities or expression are affected by Pb and Cd were retrieved from the earlier stated CTD while the set of genes implicated in the development and progression of HYP were also retrieved from the Gene Cards database. Further analysis performed revealed the 512 genes that are co-affected by Pb and Cd in their pathogenesis of HYP. The set of genes includes genes that control important biological processes such as cell cycle, cell death, inflammation, protein folding, and many others. Interestingly, the set of the mutual genes are genes that are genes affected by independent exposure to Pb and Cd. Hence, this suggests that co-exposure to Pb and Cd could pose an additive or synergistic effect on the pathogenesis of HYP. For example, the activity of blood ALAD, which provides instructions for making heme, is known to be inhibited by lead and contributes to the retention of lead in the blood [20,21].

ACE plays a significant role in the renin-angiotensin system which is responsible for the control of blood pressure via the regulation of body fluids volume. This enzyme converts angiotensin I to angiotensin II, a process that leads to the constrictions of blood vessels [22]. Júnior et al. reported that cadmium poses a dose/duration-dependent effect on ACE activity. They reported that acute exposure to low doses of cadmium increases ACE activity while chronic exposure to low doses reduces its activity [23].

Contrastingly, a study by Broseghini-Filho et al. showed that acute cadmium exposure inhibits the activity of ACE in the serum, aorta, and lung while promoting the retention of the metal in the associated tissues. The inhibition was proposed to be via competitive replacement of zinc at the active site of ACE [24]. Furthermore, Carmignani et al. [25] reported an increase in the activity of ACE following acute lead exposure, a result which is also consistent with the findings of Sharifi et al. [26]. While no study has reported the effect of Pb and Cd co-exposure on ACE activity, from the results of this study, it can be inferred that co-exposure to both metals could be capable of altering the expression and activity of ACE based on the dose and duration of exposure.

Co-exposure to Pb and Cd is known to promote systemic inflammation [27]. They also induce oxidative stress via the generation of free radicals and depletion of the body's antioxidant level [28, 29]. Several studies have reported an increased level of inflammation in hypertensive patients. Xiao et al. reported that several biomarkers of inflammation are elevated in HYP [30]. Our study revealed several pro- and anti-inflammatory markers which are targeted by Pb and Cd in HYP. These markers include IL10, TNF, IL2, IL4, IL6, CRP, IL1B, and IL17 among many others [31]. Cadmium activates inflammatory response via the upregulation of NF-K β , IL6, TNF α , and IL-1 β among others.

Conversely, the ability of lead to alter the expression of pro- and anti-inflammatory cytokines is heavily dependent on the dose [32]. It could be inferred that Pb and Cd co-exposure would both contribute to the increased progression of HYP either via the stimulation of the expression of pro-inflammatory cytokines or/and the increased generation of ROS either directly or indirectly. This claim is supported by a study by Zhang et al. [33] in which they reported that Pb and Cd co-exposure were associated with increased systemic immune inflammation. They also reported that both Cd and Pb can be immune inflammation biomarkers.

Additionally, Cd and Pb are capable of disrupting the balance between oxidant and antioxidant levels which play a vital role in the remission of oxidative stress [33]. While a study has reported that co-exposure to Cd and Pb increased the expression of oxidative stress genes such as SOD, GPx, and HSP, this study found that co-exposure to Cd and Pb could potentially alter the expression and activities of GSTA1, GSTM1, GSTO1, GSTP1, GSTT1, HSP90AB1, HSP90B1, HSPA1A, HSPA5, HSPA8, HSPA9, HSPB1, and HSPD1 in the context of HYP development and progression.

To further understand the complex relationship between genes dysregulated Pb and Cd co-exposure in HYP, the PPI network of the 512 shared targets was constructed and analyzed while GO and KEGG analyses were also performed. The top 10 targets via which Pb and Cd co-exposure disrupt the normal biological and physiological activities of the body were then identified and the biological activities of the targets among others were also revealed. The top 10 targets include GAPDH, ALB, AKT1, TP53, TNF, IL6, EGFR, CTNNB1, MYC, and CASP3.

Based on FDR values, Lipid and atherosclerosis, pathways in cancer, AGE-RAGE signaling pathway, HIF-1 signaling pathway, and human cytomegalovirus infection, fluid shear stress and atherosclerosis, and apoptosis were the top five pathways that are dysregulated by Pb and Cd co-exposure in HYP development and progression.

Both advanced glycation end products (AGEs) and their cell receptor (RAGE) have been reported to be well involved in the stiffening of the arteries while the expression of AGEs and RAGE are also known to be elevated in hypertensive

patients [34,35]. The results of this study show that AGEs and RAGE are also involved in the development and progression of HYP as a result of Pb and Cd co-exposure. The interaction of AGEs with RAGE facilitates the expression of inflammatory cytokines, increase the production of ROS via the nicotinamide adenine dinucleotide phosphate (NADPH)-oxidase, and activates the nuclear factor kappa-B (NF- κ B) which results in the contraction of arterial walls. The activation of NF- κ B leads to the generation of pro-inflammatory cytokines, which results in an elevated inflammatory response, a biological process that has been reported to be dysregulated in hypertensive patients [30].

Several accumulating evidence suggest that the activation of Hypoxia-inducible factor -1 α (HIF-1 α) is associated with arterial HYP and acts as a pathogenic mechanism for several diseases [36]. HIF plays an essential role in the response of cells to low oxygen levels and is responsible for the activation of specific genes in hypoxic conditions [36]. The deletion of HIF-1 α has been reported to combat HYP induced as a result of hypoxia. Hence, the results of this study reveal HIF signaling pathway can be regarded as a dysregulated pathway in Pb and Cd-induced HYP. Other pathways including. There exists an association between signaling pathways that are dysregulated by Cd and Pb co-exposure. This is evident from the number of genes that intercorrelate between the pathways, hence, depicting that the development and progression of Pb and Cd co-exposure-induced HYP are dependent on the deregulation of many interacting pathways.

5. CONCLUSION

Summarily, this study attempted to elucidate the molecular mechanism via which Pb and Cd co-exposure induce HYP. This study reveals the set of genes whose dysregulation by Pb and Cd co-exposure results in the induction and progression of HYP. Our study also reveals the signaling pathways via which the HYP-inducing effect of Pb and Cd co-exposure is mediated, with the most significant signaling pathways being the lipid and atherosclerosis signaling pathways. Furthermore, the stimulation of inflammation, oxidative stress, and mediating endothelial dysfunction was revealed to be an important process in Pb and Cd co-exposure mediated HYP while the inflammatory and stress markers that could be used for prognosis evaluation were revealed. While the results of this study provide

insights into some of the dysregulation caused by Pb and Cd co-exposure for HYP induction and progression. However, further *In vitro* and *In vivo* studies are needed to support the results.

SUPPLEMENTARY MATERIALS

Supplementary materials available in this link: <https://journalbjl.com/index.php/BJL/libraryFiles/downloadPublic/18>

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tchounwou PB, Yedjou CG, Patlolla AK, Sutton DJ. Heavy Metal Toxicity and the Environment, in: 2012;133–164. DOI:https://doi.org/10.1007/978-3-7643-8340-4_6.
2. Seraj F, Rahman T. Heavy metals, metalloids, their toxic effect and living systems. *Am J Plant Sci.* 2018;09:2626–2643. DOI:<https://doi.org/10.4236/ajps.2018.913191>.
3. Mehrendish R, Rahimian A, Shahriary A. Heavy metals detoxification: A review of herbal compounds for chelation therapy in heavy metals toxicity. *Journal of Herbmед Pharmacology.* 2019;8:69–77. DOI:<https://doi.org/10.15171/jhp.2019.12>.
4. Rana SVS. Perspectives in endocrine toxicity of heavy metals—a review. *Biol Trace Elem Res.* 2014;160:1–14. DOI:<https://doi.org/10.1007/s12011-014-0023-7>.
5. Andjelkovic M, Buha Djordjevic A, Antonijevic E, Antonijevic B, Stanic M, Kotur-Stevuljevic J, Spasojevic-Kalimanovska V, Jovanovic M, Boricic N, Wallace D, Bulat Z. Toxic effect of acute cadmium and lead exposure in rat blood, liver, and kidney. *Int J Environ Res Public Health.* 2019;16:274. DOI:<https://doi.org/10.3390/ijerph16020274>.
6. Substance Priority List | ATSDR, (n.d.). Available:<https://www.atsdr.cdc.gov/spl/#2019spl> (accessed September 23, 2022).
7. Živančević K, Baralić K, Jorgovanović D, Buha Djordjević A, Čurčić M, Antonijević Miljaković E, Antonijević B, Bulat Z, Đukić-Ćosić D. Elucidating the influence of environmentally relevant toxic metal mixture on molecular mechanisms involved in the development of neurodegenerative diseases: *In silico* toxicogenomic data-mining. *Environ Res.* 2021;194:110727. DOI:<https://doi.org/10.1016/j.envres.2021.110727>.
8. Zhou F, Xie J, Zhang S, Yin G, Gao Y, Zhang Y, Bo D, Li Z, Liu S, Feng C, Fan G. Lead, cadmium, arsenic, and mercury combined exposure disrupted synaptic homeostasis through activating the Snk-SPAR pathway. *Ecotoxicol Environ Saf.* 2018;163:674–684. DOI:<https://doi.org/10.1016/j.ecoenv.2018.07.116>.
9. Javorac D, Đorđević AB, Anđelković M, Tatović S, Baralić K, Antonijević E, Kotur-Stevuljević J, Đukić-Ćosić D, Antonijević B, Bulat Z. Redox and essential metal status in the brain of Wistar rats acutely exposed to a cadmium and lead mixture, *Archives of Industrial Hygiene and Toxicology.* 2020;71:197–204. DOI:<https://doi.org/10.2478/aiht-2020-71-3425>.
10. Chen X, Zhu G, Lei L, Jin T. The association between blood pressure and blood cadmium in a Chinese population living in cadmium polluted area. *Environ Toxicol Pharmacol.* 2013;36:595–599. DOI:<https://doi.org/10.1016/j.etap.2013.06.006>.
11. Eum K, Lee M, Paek D. Cadmium in blood and hypertension. *Science of The Total Environment.* 2008;407:147–153. DOI:<https://doi.org/10.1016/j.scitotenv.2008.08.037>.
12. Robles HV, Romo E, Sanchez-Mendoza A, Rios A, Soto V, Avila-Casado MC, Medina A, Escalante B. Lead exposure effect on angiotensin II renal vasoconstriction. *Hum Exp Toxicol.* 2007;26:499–507. DOI:<https://doi.org/10.1177/0960327106077597>.
13. Alghasham AA, Meki ARMA, Ismail HAS. Association of blood lead level with elevated blood pressure in hypertensive patients. *Int J Health Sci (Qassim).* 2011;5:17–27.
14. Davis AP, Grondin CJ, Johnson RJ, Sciaky D, McMorran R, Wiegers J, Wiegers TC, Mattingly CJ. The comparative toxicogenomics database: update 2019. *Nucleic Acids Res.* 2019;47:D948–D954. DOI:<https://doi.org/10.1093/nar/gky868>.

15. Davis AP, Murphy CG, Saraceni-Richards CA, Rosenstein MC, Wieggers TC, Mattingly CJ. Comparative toxicogenomics database: a knowledgebase and discovery tool for chemical-gene-disease networks, *Nucleic Acids Res.* 2009;37:D786–D792. DOI:<https://doi.org/10.1093/nar/gkn580>.
16. Davis AP, Murphy CG, Rosenstein MC, Wieggers TC, Mattingly CJ. The comparative toxicogenomics database facilitates identification and understanding of chemical-gene-disease associations: arsenic as a case study. *BMC Med Genomics.* 2008;1:48. DOI:<https://doi.org/10.1186/1755-8794-1-48>.
17. Stelzer G, Rosen N, Plaschkes I, Zimmerman S, Twik M, Fishilevich S, Stein T, Nudel R, Lieder I, Mazor Y, Kaplan S, Dahary D, Warshawsky D, Guan-Golan Y, Kohn A, Rappaport N, Safran M, Lancet D. Genecards suite: from gene data mining to disease genome sequence analyses. *Curr Protoc Bioinformatics.* 2016;54:1.30.1-1.30.33. DOI:<https://doi.org/10.1002/cpbi.5>.
18. Szklarczyk D, Gable A, Lyon D, Junge A, Wyder S, Huerta-Cepas J, Simonovic M, Doncheva N, Morris J, Bork P, Jensen L, Mering C. Protein-protein association networks with increased coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* 2019;47:D607–D613. DOI:<https://doi.org/10.1093/nar/gyk1131>.
19. Ge S, Jung D, Yao R, Shiny GO. A graphical gene-set enrichment tool for animals and plants. *Bioinformatics.* 2020;36:2628–2629. DOI:<https://doi.org/10.1093/bioinformatics/btz931>.
20. Mani MS, Kunnathully V, Rao C, Kabekkodu SP, Joshi MB, D'Souza HS. Modifying effects of δ -Aminolevulinic acid dehydratase polymorphism on blood lead levels and ALAD activity, *Toxicol Lett.* 2018;295:351–356. DOI:<https://doi.org/10.1016/j.toxlet.2018.07.014>.
21. Zhao Y. Structure and function of angiotensin converting enzyme and its inhibitors. *Chin J Biotechnol.* 2008;24:171–176. DOI:[https://doi.org/10.1016/S1872-2075\(08\)60007-2](https://doi.org/10.1016/S1872-2075(08)60007-2).
22. Pinheiro Júnior JEG, Moraes PZ, Rodriguez MD, Simões MR, Cibin F, Pinton S, Barbosa F, Junior FM, Peçanha, Vassallo DV, Miguel M, Wiggers GA. Cadmium exposure activates NADPH oxidase, renin–angiotensin system and cyclooxygenase 2 pathways in arteries, inducing hypertension and vascular damage. *Toxicol Lett.* 2020;333:80–89. DOI:<https://doi.org/10.1016/j.toxlet.2020.07.027>.
23. Broseghini-Filho GB, Almenara CCP, Vescovi MVA, de O. Faria T, Vassallo DV, Angeli JK, Padilha AS. Acute cadmium exposure reduces the local angiotensin I converting enzyme activity and increases the tissue metal content. *Biol Trace Elem Res.* 2015;166:149–156. DOI:<https://doi.org/10.1007/s12011-015-0250-6>.
24. Carmignani M, Boscolo P, Poma A, Volpe AR. Kininergic system and arterial hypertension following chronic exposure to inorganic lead. *Immunopharmacology.* 1999;44:105–110. DOI:[https://doi.org/10.1016/S0162-3109\(99\)00115-0](https://doi.org/10.1016/S0162-3109(99)00115-0).
25. Sharifi AM, Akbarloo N, Darabi R, Larijani B. Study of correlation between elevation of blood pressure and tissue ACE activity during development of hypertension in 1K1C rats. *Vascul Pharmacol.* 2004;41:15–20. DOI:<https://doi.org/10.1016/j.vph.2004.03.002>.
26. Zhang H, Yan J, Niu J, Wang H, Li X. Association between lead and cadmium co-exposure and systemic immune inflammation in residents living near a mining and smelting area in NW China. *Chemosphere.* 2022;287:132190. DOI:<https://doi.org/10.1016/j.chemosphere.2021.132190>.
27. Morcillo P, Esteban MÁ, Cuesta A. Heavy metals produce toxicity, oxidative stress and apoptosis in the marine teleost fish SAF-1 cell line. *Chemosphere.* 2016;144:225–233. DOI:<https://doi.org/10.1016/j.chemosphere.2015.08.020>.
28. Zhang Y, Huo X, Lu X, Zeng Z, Faas MM, Xu X. Exposure to multiple heavy metals associate with aberrant immune homeostasis and inflammatory activation in preschool children. *Chemosphere.* 2020;257:127257. DOI:<https://doi.org/10.1016/j.chemosphere.2020.127257>.

29. Xiao L, Harrison DG. Inflammation in hypertension. *Canadian Journal of Cardiology*. 2020;36: 635–647.
DOI:<https://doi.org/10.1016/j.cjca.2020.01.013>.
30. Das SC, Al-Naemi HA. Cadmium toxicity: Oxidative stress, inflammation and tissue injury. *Occupational Diseases and Environmental Medicine*. 2019;07:144–163.
DOI:<https://doi.org/10.4236/odem.2019.74012>.
31. Machoń-Grecka A, Dobrakowski M, Boroń M, Lisowska G, Kasperczyk A, Kasperczyk S. The influence of occupational chronic lead exposure on the levels of selected pro-inflammatory cytokines and angiogenic factors. *Hum Exp Toxicol*. 2017;36:467–473.
DOI:<https://doi.org/10.1177/0960327117703688>.
32. Zhang H, Yan J, Niu J, Wang H, Li X. Association between lead and cadmium co-exposure and systemic immune inflammation in residents living near a mining and smelting area in NW China, *Chemosphere*. 2022;287:132190.
DOI:<https://doi.org/10.1016/j.chemosphere.2021.132190>.
33. Prasad K, Mishra M. Do advanced glycation end products and its receptor play a role in pathophysiology of hypertension? *International Journal of Angiology*. 2017;26:001–011.
DOI:<https://doi.org/10.1055/s-0037-1598183>.
34. McNulty M, Mahmud A, Feely J. Advanced Glycation end-products and arterial stiffness in hypertension. *Am J Hypertens*. 2007;20:242–247.
DOI:<https://doi.org/10.1016/j.amjhyper.2006.08.009>.
35. Jiao Y, Jiang H, Lu H, Yang Y, Zhang Y, Zhang K, Liu H. Deficiency of hypoxia inducible factor-1 α promoted progression of diabetic nephropathy with hypertension. *Exp Ther Med*; 2018.
DOI:<https://doi.org/10.3892/etm.2018.6621>.
36. McGettrick AF, O'Neill LAJ. The role of hif in immunity and inflammation. *Cell Metab*. 2020;32: 524–536.
DOI:<https://doi.org/10.1016/j.cmet.2020.08.002>.

© 2023 Olukunle and Omoboyede; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95613>