



Research Article

Fasudil Attenuated 6-OHDA Cytotoxicity in PC12 Cells through Inhibition of JAK/STAT and Apoptosis Pathways

Samira Barangi¹, Seyyed Masoud Hosseini², Gholamreza Karimi^{1,2}, Soghra Mehri^{1,2*}

¹Pharmaceutical Research Center, Pharmaceutical Technology Institute, Mashhad University of Medical Sciences, Mashhad, Iran.

²Department of Pharmacodynamics and Toxicology, School of Pharmacy, Mashhad University of Medical Sciences, Mashhad, Iran.

Article Info

Article History:

Received: 24 May 2022

Accepted: 15 Jul 2022

ePublished: 21 Jul 2022

Keywords:

- Apoptosis
- Fasudil
- 6-Hydroxydopamine
- JAK/STAT
- Neurotoxicity
- Oxidative stress

Abstract

Background: 6-Hydroxydopamine (6-OHDA) is widely used to induce neurotoxicity and investigate the mechanisms of Parkinson disease. 6-OHDA causes cell injury through various mechanisms including oxidative stress, inflammation and apoptosis. The selective Rho-kinase inhibitor, fasudil displays neuroprotective effects in several neurodegenerative disorders. The aim of this study was to assess the protective effect of fasudil in PC12 cytotoxicity induced by 6-OHDA.

Methods: PC12 cells were exposed to 5, 10, 25, and 50 μM of fasudil concentrations. After 24 h, the IC_{50} value of 6-OHDA (150 μM) was added. Twenty-four hours later, the viability of cells was evaluated via MTT assay and the formation of reactive oxygen species (ROS) was measured by the fluorimetric method. At the 50 μM concentration of fasudil, with or without 6-OHDA, the changes of protein levels including STAT3, P-STAT3, JAK2, P-JAK2, and caspase-3 were determined via western blotting.

Results: Our results showed that 6-OHDA increased the intracellular level of ROS, reduced cell viability, upregulated p-STAT3/STAT3 and p-JAK2/JAK2 ratios and significantly raised cleaved caspase-3 in comparison to control group. Furthermore, pretreatment of cells with fasudil (50 μM) for 24 h could reverse all changes induced by 6-OHDA.

Conclusion: 6-OHDA caused cytotoxicity in PC12 cells through inducing oxidative stress and activating JAK/STAT and apoptosis pathways, while pretreatment with fasudil exhibited protective effect on 6-OHDA-induced neurotoxicity via the inhibition of oxidative stress and prevention of these pathways.

Introduction

6-Hydroxydopamine (6-OHDA), as a neurotoxin agent, is commonly utilized to develop Parkinson disease (PD) in experimental models. It has a similar structure to dopamine and norepinephrine and enters to the neurons through catecholamine receptors of cytoplasmic membrane.¹ In dopaminergic neurons, 6-OHDA produces free radicals, induces oxidative stress and finally causes inflammation via disturbance of dopaminergic pathway.² Additionally, the aggregation of this compound in mitochondria can induce the inhibition of electron transport chain and the loss of mitochondrial membrane potential, resulting mitochondrial dysfunction and nerve cells death.^{3,4} Moreover, free radicals produced by 6-OHDA damage DNA, cell skeletal structure and macromolecules like lipids and proteins, which leads to cell death and apoptosis of neurons.⁵ Thus, the formation of free radicals as well as reactive oxygen species (ROS), mitochondrial dysfunction and apoptosis are the main mechanisms of 6-OHDA neurotoxicity.

JAK/STAT is the one of cell signaling pathways that involves in several biological processes such as cell proliferation, hematopoiesis, immune system regulation and apoptosis.⁶ In CNS, this pathway mainly correlates to gene regulation during evolution, hormone secretion, inflammation or tumorigenesis.⁷ It is reported that JAK/STAT pathway has a pathological importance in neuroinflammation diseases like PD and leads to active innate and adaptive immune responses in these disorders.⁸ Therefore, the regulation of JAK/STAT pathway provides a suitable treatment option for Parkinson's patients.⁸

Fasudil, as a Rho-Kinase (ROCK) inhibitor, is a synthetic isoquinoline sulfonamide derivative that has an extensive range of pharmacological effects such as vasodilation and the protection of liver, kidney and brain.⁹⁻¹² It is shown that fasudil exerts its beneficial effects in various tissue damage including ischemic neuronal damage,¹³ acute kidney injury,¹⁴ myocardial infarction,¹⁵ and diabetic peripheral neuropathy¹⁶ through the regulation of different signaling pathways such as inflammation, oxidative stress

*Corresponding Author: Soghra Mehri, E-mail: mehri@mums.ac.ir

©2023 The Author(s). This is an open access article and applies the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>). Non-commercial uses of the work are permitted, provided the original work is properly cited.

and apoptosis. Moreover, in a mouse model of PD, fasudil protected dopaminergic neurons against MPTP-induced degeneration.¹⁷

Therefore, the aim of this research was to investigate whether fasudil had a protective effect against 6-OHDA induced cytotoxicity in PC12 cells and whether this effect is applied through JAK/STAT and apoptosis pathways.

Materials and Methods

Chemicals and reagents

Monohydrochloride fasudil was purchased from the Santa Cruze Biotechnology Company, USA. 6-OHDA, MTT reagent, DCF and penicillin-streptomycin (PS) were bought from Sigma (Germany). RPMI1640 medium was made a purchase from Bioidea Company (Iran). Dimethyl sulfoxide (DMSO) from Merck (Germany) and fetal bovine serum (FBS) from Gibco (USA) were obtained. Moreover, all of used antibodies were taken from Cell Signaling Company (USA).

Cell Culture

PC12, derived from rat pheochromocytoma cell line, was obtained from Pasteur Institute (Tehran, Iran). Complete culture media containing RPMI 1640, 10% heat-inactivated FBS, and 1% penicillin-streptomycin was used to cultivate the cells. Then, cells were maintained in 37 °C incubator accompanied by 95% humidified atmosphere and 5% CO₂. The cells were passage at the confluence of 80%.¹⁸

MTT assay

To evaluate the cell viability by MTT assay and calculate the IC₅₀ value (the inhibitory concentration of 50%), PC12 cells (4000 cells/well) were planted in microplate for 24 h. Then, the cells were exposed to various concentration of 6-OHDA (50, 100, 150, 200, 300 μM) and fasudil (5, 10, 25, 50, 100 μM) for 24 h and 48 h, respectively. When the exposure time is finished, the final concentration of MTT solution (0.5 mg/mL) was added to each well and the cells were incubated for 3 h at 37 °C. The upper medium was then removed from each well, and DMSO (150 μl) was added to dissolve the purple formazan product. A microplate reader (Start Fax-2100, UK) was used to read the absorbance at 545 and 630 nm. Then, the proportion of viable cells and the IC₅₀ value were calculated.¹⁸

In order to assess the fasudil effect on 6-OHDA-induced cytotoxicity, the various concentrations of fasudil (5, 10, 25, 50, 100 μM) were exposed to PC12 cells in 96-well microplate for 24 h and then 6-OHDA (150 μM; the concentration which determined based on the concentration assay of IC₅₀) was added. After 24 h, MTT solution (0.5 mg/mL) was added to each well and the cells were incubated for 3 h at 37 °C. The upper medium was then removed from each well, and DMSO (150 μl) was added to dissolve the purple formazan product and the absorbance was read at 545 and 630 nm. The cell viability was represented as a percentage of the value in control cultures.

Measurement of intracellular ROS production

DCFH-DA reagent was applied to evaluate the amount of ROS. In this line, PC12 cells (4000 cells/well) were seeded in 96-well microplate for 24 h. At first, the cells were incubated by fasudil (5, 10, 25, 50, 100 μM) before exposed to 6-OHDA (150 μM) for 24 h. After removing the upper mediate, the final concentration of DCFH-DA (10 μM) was added to the cultured cells for 30 min in a darkness at 37 °C in an incubator. After that, cells were washed with PBS and the fluorescence intensity of DCF was read at 485 and 538 nm for the excitation and emission wavelength, respectively by a microplate-reader.¹⁹

Western blot analysis

This test was used to investigate the changes in the protein levels including JAK2, P-JAK2, STAT3, P-STAT3 and caspase-3 in PC12 cells treated by 6-OHDA and fasudil. For this purpose, PC12 cells were exposed to fasudil (50 μM) for 24 h before 6-OHDA (150 μM). After 24 h, the cells were separated with trypsin and washed by PBS buffer. After centrifuge and removing supernatant liquid, lysis buffer containing 2 mM EDTA, 2 mM EGTA, 50 mM Tris-HCl, 0.2% W/V sodium deoxycholate, 10 mM NaF, 10 mM β-glycerophosphate, 1 mM sodium orthovanadate, along with protease inhibitor cocktail and phenylmethylsulfonyl fluoride (PMSF) were added to the cells. The concentration of proteins in samples were measured based on Bradford protein assay with some modification.²⁰

Afterward, the separation of proteins were done by means of SDS-PAGE electrophoresis technique and transferred to a PVDF membrane. Blocking step was done on PVDF membranes applying by 5% skimmed milk or 5% BSA (Bovine Serum Albumin) for non-phosphorylated or phosphorylated proteins, respectively for 2 h at room temperature. Thereupon, membranes were incubated with appropriate primary rabbit monoclonal antibodies against caspase-3 (Cell Signaling, #9665), STAT3 (Cell Signaling, #12640), P-STAT3 (Cell Signaling, #9145), JAK2 (Cell Signaling, #3230), P-JAK2 (Cell Signaling, #3776) and mouse monoclonal antibody against β-actin (Cell Signaling, #3700). After membranes washing, they were incubated with anti-rabbit (Cell Signaling, #7074) or anti-mouse (Cell Signaling, #7076) IgG antibody labeled with horseradish peroxidase as secondary antibodies. To visualize the protein bands, enhanced chemiluminescence was used by Alliance 4.7 Gel doc (UK) and all of bonds were analyzed by UVtec software (UK). Then, the protein expression levels were normalized with their corresponding β-actin level.

Statistics analysis

All data analyzed by means of GraphPad Prism software (version 8). All results are exhibited as means ± SD. The comparison among different groups were done by one-way analysis of variance (ANOVA) and Tukey-Kramer post hoc statistical test. P<0.05 was considered as statistically significant levels.




Figure 1. The effect of 6-OHDA (50-300 µM; A), fasudil (5-100 µM; B), and the effect of fasudil (F) on 6-OHDA-induced cytotoxicity (C) on PC12 cells. Cell viability was evaluated by MTT test. Data are expressed as means \pm SD from three independent experiments. Data were analyzed by one-way ANOVA following Tukey-kramer post test for multiple comparisons. ** $P < 0.01$ and *** $P < 0.001$ vs. control group, # $P < 0.05$ vs. 6-OHDA treated group.

Results

6-OHDA effect in PC12 cell viability

The cells were treated 24 h by various concentrations of 6-OHDA (50, 100, 150, 200, 300 µM). The result of MTT assay showed that the exposure to all 6-OHDA concentrations decreased cell viability compared to control (Figure 1A). Additionally, the calculated IC_{50} for 6-OHDA after 24 h exposure was 150 ± 26.46 µM in PC12 cells.

Fasudil effect in PC12 cell viability

Figure 1B shows that the exposing PC12 cells to fasudil at doses of 5, 10, 25, and 50 µM for 48 h had no effect on cell viability as compared to untreated cells, while 100 µM concentration of fasudil could significantly reduce the viability of cells compared with control group ($P < 0.01$).

Fasudil effect on 6-OHDA-induced cytotoxicity in PC12 cells

To investigate the fasudil protective effect on cytotoxicity by 6-OHDA, the cells were treated 24 h by various concentrations of fasudil (5-50 µM) before 24 h incubated by 6-OHDA (150 µM). Figure 1C shows that the 6-OHDA exposure resulted in cell viability reduction compared to control group, while the pretreatment of cells with 50 µM fasudil markedly elevated cell viability compared with 6-OHDA exposure group ($P < 0.05$). The pretreatment of

cells with other concentrations of fasudil had no significant effect on PC12 cell viability compared with 6-OHDA group (Figure 1C).

Fasudil effect on ROS production by 6-OHDA in PC12 cells

The result showed that after 24 h exposure to 6-OHDA (150 µM), the production of intracellular ROS increased in comparison to control ($P < 0.001$). Remarkably, the fasudil pretreatment at 5, 10, 25 and 50 µM concentrations could reduce the intracellular ROS generation compared with 6-OHDA group (Figure 2).

Fasudil and 6-OHDA effects on JAK2 and STAT3 proteins levels in PC12 cells

The PC12 cells treatment by 6-OHDA enhanced P-JAK2/JAK2 and P-STAT3/STAT3 ratios in comparison to untreated cells ($P < 0.05$ and $P < 0.01$, respectively), whereas the fasudil pretreatment reduced these ratios as compared with 6-OHDA treated cells ($P < 0.01$) (Figure 3).

Fasudil and 6-OHDA effects on caspase-3 protein level in PC12 cell

As shown in Figure 4, the PC12 cells treatment with 6-OHDA markedly enhanced the levels of caspase-3 cleaved compared with control ($P < 0.01$), while pretreatment of





Figure 2. The effect of fasudil (F) on 6-OHDA-induced ROS production in PC12 cells. ROS generation was determined using the DCFH-DA reagent. Data are expressed as means \pm SD from three independent experiments. Data were analyzed by one-way ANOVA following Tukey-kramer post test for multiple comparisons. *** $P < 0.001$ vs. control group, ### $P < 0.001$ vs. 6-OHDA group.

the cells with fasudil could decrease the cleaved caspase-3 level in PC12 cells compared to 6-OHDA group ($P < 0.05$). In addition, the results demonstrated no significant differences between groups in the level of pro caspase-3.



Discussion

The purpose of this study was to evaluate the fasudil protective effect on 6-OHDA-induced cytotoxicity through JAK/STAT and apoptosis pathways. The results showed that 6-OHDA exposure reduced cell viability, induced oxidative stress and caused cell death through the enhancement of apoptosis and the phosphorylated form of JAK2 and STAT3 proteins in PC12 cells. The fasudil pretreatment reversed the 6-OHDA-induced alterations in oxidative stress, JAK/STAT and apoptosis proteins.

6-OHDA is one of the neurotoxins that induce neurotoxicity in diverse models of PD. Oxidative stress as well as apoptosis are considered as main mechanisms of neurotoxicity by 6-OHDA in both *in vitro* and *in vivo* models.²¹⁻²³ In this regard, 6-OHDA diminished cell viability, increased ROS and induced cell death via the enhancement of apoptosis proteins such as Bax and caspase-3 in PC12 cells.²²⁻²⁴ The neurotoxicity induced by 6-OHDA attenuated by medicinal plants or synthetic compounds which have a potent antioxidant property.²²⁻²⁴ For instance, carvacrol is a monoterpene phenol and main constituents of many aromatic plants that could increase cell viability and reduce intracellular lipid peroxidation and ROS as well as apoptosis in PC12 cells.²⁵ Moreover, 6-OHDA induced mitochondrial dysfunction, ROS production, and p38 MAPK activation and augmented the level of cleaved caspase-9 and caspase-3 in SH-SY5Y cells.²⁶




Figure 3. The effect of fasudil (F) and 6-OHDA on protein level of JAK2, phosphor (P)-JAK2, STAT3 and phosphor (P)-STAT3 proteins in PC12 cells. The western blot bands (A and C), the bars (B and D) which exhibiting the densitometry analysis of blots for P-JAK2/JAK2 and P-STAT3/STAT3 ratio. Equal loading of proteins is illustrated by β -actin bands. Data are expressed as means \pm SD from three independent experiments. Data were analyzed by one-way ANOVA following Tukey-kramer post test for multiple comparisons. * $P < 0.05$ and ** $P < 0.01$ vs. control group, ## $P < 0.01$ vs. 6-OHDA group.




Figure 4. The effect of fasudil (F) and 6-OHDA on protein level of caspase-3 (pro and cleaved) protein in PC12 cells. The western blot bands (A), the bars (B) which exhibiting the densitometry analysis of bands for the level of caspase-3 (pro and cleaved) protein. Equal loading of proteins is illustrated by β -actin bands. Data are expressed as means \pm SD from three independent experiments. Data were analyzed by one-way ANOVA following Tukey-kramer post test for multiple comparisons. * $P < 0.01$ vs. control group, # $P < 0.05$ vs. 6-OHDA group.

Similar result of 6-OHDA neurotoxicity was reported by Hanrott *et al.*²⁷ in which 6-OHDA initiated mitochondrial dysfunction, oxidative stress and apoptosis through the caspases-3 and caspase-7 activating in PC12 cells.

Our results indicated that the exposure to 6-OHDA significantly reduced cell viability and raised ROS production in PC12 cells. In addition, 6-OHDA could induce apoptosis pathway through the cleaved caspase-3 elevation in PC12 cells. Furthermore, fasudil concentration was able to prevent cytotoxicity induced by 6-OHDA. The ROS production as well as caspase-3 expression were decreased following pretreatment with fasudil in PC12 cell. Kianfar *et al.*¹⁸ reported the fasudil protective effect in acrylamide-induced toxicity in PC12 cells through the decreased ROS production and downregulation of apoptosis proteins such as Bax/Bcl2 ratio, caspase-3 and caspase-9. Zhang *et al.*²⁸ reported that fasudil attenuated myocardial infarction size and apoptosis in cardiomyocytes and diminished the p-JAK level in ischemia/reperfusion injury model of rat heart. Besides, fasudil exhibited the protective effect in ischemic neuronal damage by increasing ROCK activity and inhibition of cell death.¹³ Also, Song *et al.*²⁹ demonstrated that fasudil could protect the rat retina against I/R injury through the reduction of

Bax/Bcl-2 mRNA ratio, the caspase-3 level and inducible nitric oxide synthase (iNOS) expression. In addition, fasudil mesylate can be effectively improved oxidative stress-induced neuronal damage through the decrease of ROS accumulation as well as downregulation of Bax/Bcl-2 ratio in PC12 cells.³⁰

JAK/STAT, as an important intracellular signaling pathway, has an essential role in different biological processes such as cell proliferation and differentiation, immune response, apoptosis and inflammation.^{6,7} The role of JAK/STAT pathway has been reported in neuroinflammation diseases and several models of PD. For instance, the administration of AZD1480, a JAK1/JAK2 inhibitor, suppressed JAK, STAT1 and STAT3 activation in microglia and macrophages cells and prevented the degeneration of dopaminergic neurons in rats treated by α -synuclein as a model of PD.³¹ Wang *et al.*³² reported that the phosphorylation of JAK2 and STAT3 significantly elevated in LPS-treated microglia BV2 cells. Moreover, the neurotoxin MPP⁺ treatment increased the expression level of STAT1 as well as STAT1 phosphorylation and induced apoptosis in cerebellar granule neurons cells.³³ In this regard, the type I interferons (IFN- α and IFN- β) treatment induced neurotoxicity in mouse primary neurons and

SH-SY5Y neuroblastoma cells. In this study, the apoptotic cell death by cleavage of caspase-3, caspase-9, caspase-7 and cytochrome C release as well as STAT1 and STAT3 phosphorylation were considered as main mechanisms of IFNs. Interestingly, the effects of IFNs reversed by pyridone 6 as a JAK inhibitor.³⁴ In this matter, our results exhibited that in PC12 cells, the 6-OHDA exposure markedly enhanced the phosphorylation levels of JAK2 and STAT3. It seems that there is an interplay between Rho-kinase and JAK/STAT signaling pathways. Loucks *et al.*³⁵ mentioned that JAK/STAT is a pro-apoptotic pathway, which regulates the downstream of Rho family in cerebellar granule neurons. Moreover, the activation of Rho family indirectly stimulates the STAT transcriptional activity as well as STAT3 phosphorylation in cytosolic and nuclear fractions through the enhancement of JAK2 phosphorylation in HEK293 cells.³⁶ Besides, Rho-kinase inhibitor could prevent STAT3 phosphorylation and activation.³⁶ Interestingly, Jaffe *et al.*³⁷ demonstrated that the RhoA activation by leptin was inhibited following the treatment of human colonic epithelial cell line with JAK2 inhibitor (AG490). Furthermore, it has been demonstrated that CPL409116, as a novel dual JAK/ROCK inhibitor, could block the STAT1 and STAT3 phosphorylation and reduce pro-inflammatory cytokines expression in systemic lupus erythematosus.³⁸ These evidence suggest that Rho-kinase pathway may be an effective regulator in the JAK2/STAT3 pathway activation and inhibition. In this regard, our results showed that the fasudil pretreatment declined the phosphorylation levels of JAK2 and STAT3 and decreased P-JAK2/JAK2 and P-STAT3/STAT3 ratios in PC12 cells treated with 6-OHDA.

Conclusion

In summary, present study exhibited that 6-OHDA induced cytotoxicity in PC12 cells via increased oxidative stress and the activation of JAK/STAT and apoptosis pathways. On the other hand, the Rho-kinase inhibitor, fasudil, displayed a neuroprotective effect against 6-OHDA in PC12 cells through the inhibition of oxidative stress and suppression of JAK/STAT and apoptosis pathways, which are important mechanisms of 6-OHDA neurotoxicity. However, more experimental and clinical studies are suggested to investigate the role of fasudil in people who are suffering from PD.

Acknowledgements

Authors are thankful to the Vice Chancellor of Research, Mashhad University of Medical Sciences, Mashhad, Iran for financial support (Grant Number: 970797).

Author Contributions

All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by SMH and SB. The first draft of the manuscript was written by SB and all authors commented on previous versions of the manuscript. All authors read

and approved the final manuscript.

Conflict of Interest

The authors report no conflicts of interest.

References

1. Bove J, Prou D, Perier C, Przedborski S. Toxin-induced models of parkinson's disease. *NeuroRx*. 2005;2(3):484-94. doi:10.1602/neurorx.2.3.484
2. Mehan S, Kaur G, Dudi R, Rajput M, Kalra S. Restoration of mitochondrial dysfunction in 6-hydroxydopamine induced parkinson's disease: A complete review. *Open J Park Dis Treat*. 2017;1(1):001-26. doi:10.17352/ojpd.000001
3. Ren J, Yuan L, Wang W, Zhang M, Wang Q, Li S, et al. Tricetin protects against 6-OHDA-induced neurotoxicity in parkinson's disease model by activating NRF2/HO-1 signaling pathway and preventing mitochondria-dependent apoptosis pathway. *Toxicol Appl Pharmacol*. 2019;378:114617. doi:10.1016/j.taap.2019.114617
4. Iglesias Gonzalez PA, Conde MA, Gonzalez-Pardo V, Uranga RM, Salvador GA. In vitro 6-hydroxydopamine-induced neurotoxicity: New insights on nfkappab modulation. *Toxicol In Vitro*. 2019;60:400-11. doi:10.1016/j.tiv.2019.06.019
5. Tabatabaei Mirakabad FS, Khoramgah MS, Tahmasebinia F, Darabi S, Abdi S, Abbaszadeh HA, et al. The effect of low-level laser therapy and curcumin on the expression of LC3, ATG10 and bax/bcl2 ratio in pc12 cells induced by 6-hydroxide dopamine. *J Lasers Med Sci*. 2020;11(3):299-304. doi:10.34172/jlms.2020.50
6. Bolli R, Dawn B, Xuan YT. Role of the jak-stat pathway in protection against myocardial ischemia/reperfusion injury. *Trends Cardiovasc Med*. 2003;13(2):72-9. doi:10.1016/s1050-1738(02)00230-x
7. Nicolas CS, Amici M, Bortolotto ZA, Doherty A, Csaba Z, Fafouri A, et al. The role of JAK-STAT signaling within the CNS. *JAKSTAT*. 2013;2(1):e22925. doi:10.4161/jkst.22925
8. O'Shea JJ, Pesu M, Borie DC, Changelian PS. A new modality for immunosuppression: Targeting the jak/stat pathway. *Nat Rev Drug Discov*. 2004;3(7):555-64. doi: 10.1038/nrd1441
9. Ho TJ, Huang CC, Huang CY, Lin WT. Fasudil, a rho-kinase inhibitor, protects against excessive endurance exercise training-induced cardiac hypertrophy, apoptosis and fibrosis in rats. *Eur J Appl Physiol*. 2012;112(8):2943-55. doi:10.1007/s00421-011-2270-z
10. Ikeda H, Kume Y, Tejima K, Tomiya T, Nishikawa T, Watanabe N, et al. Rho-kinase inhibitor prevents hepatocyte damage in acute liver injury induced by carbon tetrachloride in rats. *Am J Physiol Gastrointest Liver Physiol*. 2007;293(4):G911-7. doi:10.1152/ajpgi.00210.2007
11. Hayashi K, Wakino S, Kanda T, Homma K, Sugano

- N, Saruta T. Molecular mechanisms and therapeutic strategies of chronic renal injury: Role of rho-kinase in the development of renal injury. *J Pharmacol Sci.* 2006;100(1):29-33. doi:10.1254/jphs.fmj05003x6
12. Zhao Y, Liu B, Wang Y, Xiao B. Effect of fasudil on experimental autoimmune neuritis and its mechanisms of action. *Braz J Med Biol Res.* 2020;53(1):e8669. doi:10.1590/1414-431X20198669
 13. Yamashita K, Kotani Y, Nakajima Y, Shimazawa M, Yoshimura S, Nakashima S, et al. Fasudil, a rho kinase (rock) inhibitor, protects against ischemic neuronal damage in vitro and in vivo by acting directly on neurons. *Brain Res.* 2007;1154:215-24. doi:10.1016/j.brainres.2007.04.013
 14. Wang Y, Zhang H, Yang Z, Miao D, Zhang D. Rho kinase inhibitor, fasudil, attenuates contrast-induced acute kidney injury. *Basic Clin Pharmacol Toxicol.* 2018;122(2):278-87. doi:10.1111/bcpt.12895
 15. Zhou FT, Ma K. Fasudil protects against isoproterenol-induced myocardial infarction in mice via inhibiting rho/rock signaling pathway. *Eur Rev Med Pharmacol Sci.* 2020;24(10):5659-67. doi:10.26355/eurev_202005_21357
 16. Jiang DQ, Xu LC, Jiang LL, Li MX, Wang Y. Fasudil combined with methylcobalamin or lipoic acid can improve the nerve conduction velocity in patients with diabetic peripheral neuropathy: A meta-analysis. *Medicine (Baltimore).* 2018;97(27):e11390. doi:10.1097/MD.00000000000011390
 17. Chen M, Liu A, Ouyang Y, Huang Y, Chao X, Pi R. Fasudil and its analogs: A new powerful weapon in the long war against central nervous system disorders? *Expert Opin Investig Drugs.* 2013;22(4):537-50. doi:10.1517/13543784.2013.778242
 18. Kianfar M, Nezami A, Mehri S, Hosseinzadeh H, Hayes AW, Karimi G. The protective effect of fasudil against acrylamide-induced cytotoxicity in PC12 cells. *Drug Chem Toxicol.* 2018;43(6):595-601. doi:10.1080/01480545.2018.1536140
 19. Li Y, Shi W, Li Y, Zhou Y, Hu X, Song C, et al. Neuroprotective effects of chlorogenic acid against apoptosis of PC12 cells induced by methylmercury. *Environ Toxicol Pharmacol.* 2008;26(1):13-21. doi:10.1016/j.etap.2007.12.008
 20. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976;72:248-54. doi:10.1006/abio.1976.9999
 21. Lee J, Song K, Huh E, Oh MS, Kim YS. Neuroprotection against 6-OHDA toxicity in PC12 cells and mice through the NRF2 pathway by a sesquiterpenoid from *tussilago farfara*. *Redox Biol.* 2018;18:6-15. doi:10.1016/j.redox.2018.05.015
 22. Hadipour E, Fereidoni M, Tayarani-Najaran Z. Betanin attenuates oxidative stress induced by 6-OHDA in PC12 cells via s SAPK/JNK and PI3 K pathways. *Neurochem Res.* 2020;45(2):395-403. doi:10.1007/s11064-019-02927-w
 23. Ramazani E, YazdFazeli M, Emami SA, Mohtashami L, Javadi B, Asili J, et al. Protective effects of cinnamomum verum, cinnamomum cassia and cinnamaldehyde against 6-OHDA-induced apoptosis in PC12 cells. *Mol Biol Rep.* 2020;47(4):2437-45. doi:10.1007/s11033-020-05284-y
 24. Ramazani E, Fereidoni M, Tayarani-Najaran Z. Protective effects of vitamin k2 on 6-OHDA-induced apoptosis in PC12 cells through modulation bax and caspase-3 activation. *Mol Biol Rep.* 2019;46(6):5777-83. doi:10.1007/s11033-019-05011-2
 25. Manouchehrabadi M, Farhadi M, Azizi Z, Torkaman-Boutorabi A. Carvacrol protects against 6-hydroxydopamine-induced neurotoxicity in vivo and in vitro models of parkinson's disease. *Neurotox Res.* 2020;37(1):156-70. doi:10.1007/s12640-019-00088-w
 26. Ikeda Y, Tsuji S, Satoh A, Ishikura M, Shirasawa T, Shimizu T. Protective effects of astaxanthin on 6-hydroxydopamine-induced apoptosis in human neuroblastoma SH-SY5Y cells. *J Neurochem.* 2008;107(6):1730-40. doi:10.1111/j.1471-4159.2008.05743.x
 27. Hanrott K, Gudmunsen L, O'Neill MJ, Wonnacott S. 6-hydroxydopamine-induced apoptosis is mediated via extracellular auto-oxidation and caspase 3-dependent activation of protein kinase cdelta. *J Biol Chem.* 2006;281(9):5373-82. doi:10.1074/jbc.M511560200
 28. Zhang J, Li XX, Bian HJ, Liu XB, Ji XP, Zhang Y. Inhibition of the activity of Rho-kinase reduces cardiomyocyte apoptosis in heart ischemia/reperfusion via suppressing JNK-mediated AIF translocation. *Clin Chim Acta.* 2009;401(1-2):76-80. doi:10.1016/j.cca.2008.11.016
 29. Song H, Gao D. Fasudil, a rho-associated protein kinase inhibitor, attenuates retinal ischemia and reperfusion injury in rats. *Int J Mol Med.* 2011;28(2):193-8. doi:10.3892/ijmm.2011.659
 30. Li Q, Liu D, Huang X, Guo L. Fasudil mesylate protects PC12 cells from oxidative stress injury via the bax-mediated pathway. *Cell Mol Neurobiol.* 2010;31(2):243-50. doi:10.1007/s10571-010-9614-9
 31. Qin H, Buckley JA, Li X, Liu Y, Fox TH, 3rd, Meares GP, et al. Inhibition of the JAK/STAT pathway protects against alpha-synuclein-induced neuroinflammation and dopaminergic neurodegeneration. *J Neurosci.* 2016;36(18):5144-59. doi:10.1523/JNEUROSCI.4658-15.2016
 32. Wang B, Wang X-q, Yang S-s, Liu X, Feng D-y, Lu F-f, et al. Neuroprotective effects of nitidine in parkinson's disease models through inhibiting microglia activation: Role of the JAK2-STAT3 pathway. *RSC Adv.* 2016;6(75):71328-37. doi: 10.1039/c6ra11759g
 33. Junyent F, Alvira D, Yeste-Velasco M, de la Torre AV, Beas-Zarate C, Sureda FX, et al. Prosurvival role of JAK/STAT and AKT signaling pathways in mpp+-induced

- apoptosis in neurons. *Neurochem Int.* 2010;57(7):774-82. doi:10.1016/j.neuint.2010.08.015
34. Dedoni S, Olianias MC, Onali P. Interferon- β induces apoptosis in human SH-SY5Y neuroblastoma cells through activation of JAK-STAT signaling and down-regulation of PI3K/Akt pathway. *J Neurochem.* 2010;115(6):1421-33. doi:10.1111/j.1471-4159.2010.07046.x
35. Loucks FA, Le SS, Zimmermann AK, Ryan KR, Barth H, Aktories K, et al. Rho family gtpase inhibition reveals opposing effects of mitogen-activated protein kinase kinase/extracellular signal-regulated kinase and janus kinase/signal transducer and activator of transcription signaling cascades on neuronal survival. *J Neurochem.* 2006;97(4):957-67. doi:10.1111/j.1471-4159.2006.03802.x
36. Reipschläger S, Kubatzky K, Taromi S, Burger M, Orth J, Aktories K, et al. Toxin-induced rhoa activity mediates ccl1-triggered signal transducers and activators of transcription protein signaling. *J Biol Chem.* 2012;287(14):11183-94. doi:10.1074/jbc.M111.313395
37. Jaffe T, Schwartz B. Leptin promotes motility and invasiveness in human colon cancer cells by activating multiple signal-transduction pathways. *Int J Cancer.* 2008;123(11):2543-56. doi:10.1002/ijc.23821
38. Dulak-Lis M, Bujak A, Gala K, Banach M, Kędzierska U, Miszkiel J, et al. A novel JAK/ROCK inhibitor, CPL409116, demonstrates potent efficacy in the mouse model of systemic lupus erythematosus. *J Pharmacol Sci.* 2021;145(4):340-8. doi:10.1016/j.jphs.2021.02.002