



Assessment of Proximate and Sensory Properties of Gari and Fufu Flour from Cassava Roots Stored for 14 Days

Nwadili, V. U ^{a,b,*}, Ahemen, S. A. ^{b,c}, Yusufu, M. I. ^{b,d},
Yande, J. T. ^{b,e}, Ndi, B.B ^{b,e} and Yepshak, N ^{b,f}

^a National Root Crops Research Institute (NRCRI), PMB 7006, Umuahia, Abia State, Nigeria.

^b Center for Food Technology and Research, Benue State University Makurdi, Benue State, Nigeria.

^c Akperan Orshi Polytechnic, Yandev, PMB 181, Gboko, Benue State, Nigeria.

^d Department of Food Science and Technology, Federal Polytechnic, Idah, Kogi State, Nigeria.

^e Department of Chemistry, Benue State University PMB 102119, Makurdi, Benue, Nigeria.

^f College of Agriculture, Garkawa, Plateau State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors NVU and ASA designed the study. Author NVU wrote the protocol, managed the literature searches and wrote the first draft of the manuscript. Author NBB performed the statistical analysis. Authors NBB, YJT and YN did the manuscript review. Authors ASA and YMI made corrections of the drafts and general supervision of the study. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/118329>

Original Research Article

Received: 02/04/2024

Accepted: 06/06/2024

Published: 21/06/2024

ABSTRACT

Aims: To investigate the proximate composition *gari* and *fufu* flour and the sensory attributes of reconstituted *gari* and *fufu* dough processed from cassava roots stored for a period of 14 days.

Study Design: Complete randomized design (CRD)

Place and Duration of Study: Product Development Programme, National Root Crops Research Institute, Umudike, Umuahia, Abia State, Nigeria, between October, 2022 and March, 2023.

*Corresponding author: E-mail: vickyzeigbo@yahoo.com;

Cite as: V. U, Nwadili, Ahemen, S. A., Yusufu, M. I., Yande, J. T., Ndi, B.B, and Yepshak, N. 2024. "Assessment of Proximate and Sensory Properties of Gari and Fufu Flour from Cassava Roots Stored for 14 Days". *Asian Journal of Food Research and Nutrition* 3 (3):544-53. <https://www.journalajfrn.com/index.php/AJFRN/article/view/154>.

Methodology: Four cassava varieties, two white – fleshed (TME 419, NR 87/184), and two bio-fortified varieties (TMS 01/1368 and TMS 07/0539) were stored at ambient conditions for a period of 14 days on shelves. *Gari* and *fufu* flour were processed on days 0, 7 and 14 using standard methods. The acceptability of reconstituted *gari* and *fufu* dough was determined using a nine-point hedonic ranking scale ranging from 9 (liked extremely) to 1 (disliked extremely).

Results: Proximate composition of the stored roots showed significant decrease in moisture contents with highest percentage of 70.12 ± 0.04 at day 0 and 50.05 ± 0.04 at day 14 across varieties. Opposite trend was observed in carbohydrate contents of the varieties across the period of storage with values in the range 24.03 ± 0.01 to 44.45 ± 0.01 . *Fufu* flour recorded moisture contents between 7.44 ± 0.03 - 9.03 ± 0.03 while *gari* had moisture values between 4.12 ± 0.01 to 12.15 ± 0.05 % across all four varieties. Moisture and ash contents of *fufu* flour and *gari* were within acceptable maximum limits of 12 % and 3 % Codex Alimentarius standards. Sensory evaluation results showed that storage of cassava roots of varieties NR 87/184, and TME 419 for up to 14 days gave acceptable *gari* quality while *fufu* from variety NR 87/184 was most acceptable.

Conclusion: Cassava varieties NR 87/184 and TME 419 could be promoted for *gari* and *fufu* processing for longer period up to 14 days after harvest.

Keywords: Cassava varieties; proximate; sensory; *gari*; *fufu* flour; processing.

1. INTRODUCTION

Cassava (*Manihot esculenta* Crantz) is grown extensively in tropical and subtropical regions of the world. It is a vital staple food for millions in Africa, including Nigeria [1,2]. Post-harvest physiological deterioration is a major barrier to the use of cassava roots. It results in a considerable loss of storage roots and lowers the crop's food, feed, and market value [3,4]. *Gari* and *fufu* are products derived from cassava roots and are popularly consumed in West Africa. *Gari* is the most commercial and useful product from cassava roots, it is granular, partially gelatinized by toasting, has a slightly sour taste and fermented flavor [5,6]. Cassava *fufu* is a fermented white paste made from cassava and it is ranked next to *gari* as an indigenous food of most Nigerians in the South West and South East [7]. The shelf life of cassava *fufu* is short and deterioration rate is high because it is processed as a wet paste with moisture content of about 50 % therefore, not suitable for large-scale and commercial purposes [8]. A modern technology developed to extend the shelf life and market quality of cassava *fufu* is obtained by drying at high temperature of about 60 – 65°C to produce flour which can be further reconstituted with hot water [9]. *Gari* quality: taste (acidity or sourness), swelling capacity, color, texture, crispiness, and absence of foreign matter (cleanliness) is based on certain indices judged by both processors and consumers alike [10,11]. The quality of *fufu* is known to vary from one location to another and processing technique plays a vital role in the variation in quality [12]. Factors such as cassava variety, differences in

dry matter content and the quality of the roots influence the quality of *fufu* [13]. Other factors that could affect the physical quality of *fufu* are delay in processing cassava roots after harvest and increase in the length of soaking time during processing [13].

Freshly harvested cassava roots deteriorate two to three days after harvest as a result of post-harvest physiological deterioration (PPD), this deterioration has been linked with microbiological and physiological changes within the root [14,15,16,17]. Studies have shown that the physical and sensory attributes of *gari* from cassava roots stored in trench and polythene bags were acceptable and comparable to *gari* from freshly harvested cassava roots [18]. Another study showed that in terms of composition, insignificant changes were found in the acidity and moisture content of *gari* and *fufu* from cassava roots stored for varying period of storage ranging from 0–4 days [19]. Recent research have shown that certain cassava varieties have improved postharvest storage characteristics such as increased shelf-life for up to fourteen days after harvest [15].

There is no much information reported on the quality of *gari* and *fufu* flour from cassava roots tolerant to PPD stored for up to 14 days. The main focus of this investigation was to study the quality of processed *gari*, *fufu* flour made from cassava roots that were kept for 14 days. Hence determining the proximate and sensory properties were crucial. Availability of such information would guide in selection of cassava varieties that are suitable for marketing and

utilization in enterprises requiring fresh cassava roots.

2. MATERIALS AND METHODS

2.1 Material Selection

Four cassava varieties were used for this study. Two white fleshed cassava roots (NR87/184 and TME 419) and two yellow fleshed cassava roots (TMS 01/1368 and TMS 07/0539). Root samples were obtained from plants grown at the Cassava Programme Research field of the National Root Crops Research Institute Umudike, Umuahia, Abia State, Nigeria. The cassava roots were manually harvested at 12 months after planting and great during harvesting was taken to avoid injuries to the roots as root injury accelerates the onset of PPD [20]. 30 kg each of cassava roots of the four varieties NR87/184, TME 419, TMS 07/0539 and TMS 01/1368 without mechanical damage or pre-harvest rot were selected. Each of the 30 kg harvested cassava roots was divided into 3 equal parts of 10 kg each which represented day 0, day 7 and day 14. Samples were drawn from each of the 10 kg on each of the sampling days (0, day 7 and day 14) for laboratory analysis and the remaining roots were subsequently processed into *gari* and *fufu* flour on the respective days.

2.2 Cassava Roots Processing

2.2.1 Processing cassava roots into *Gari*

Gari was processed on days 0, 7 and 14 using the method as described by Abass et al. [10]. The roots of each variety were peeled, washed and grated into a mash. The mash was placed in a hessian sack, and left for 48 hours to ferment at ambient temperature. After 48 hours, the fermented mash was dewatered and the resulting cake was then sieved. The pulverized cake was toasted. The resulting toasted granules (*gari*) were allowed to cool and packaged in polyethylene bags for further analysis.

2.2.2 Processing of Cassava roots into *fufu*

Fufu flour was processed on days 0, 7 and 14 using the method as described by Aniedu & Omodamiro, [21]. The roots of each variety were peeled and washed. The washed roots soaked in clean water and left for 48 h after which the submerged roots were removed and grated into a fine mash. The grated mash was left for 24 h before dewatering. The dewatered mash was sifted, dried and allowed to cool. The resulting granules were milled and sieved to obtain fine

flour which were subsequently packed in polyethylene bags for further analysis

2.3 Proximate Composition of Cassava Roots, *gari* and *fufu* Flour

The proximate composition of cassava roots on days 0, 7 and 14, *gari* and *fufu* flour processed at the same days 0, 7 and 14 were determined using standard methods [22].

2.4 Reconstitution of *gari* and *fufu* Flour into Dough

Gari and *fufu* flour processed at days 0, 7 and 14 were reconstituted into a dough using the modified method described by Udoro et al. [23]. The *Gari* dough was prepared by adding about 100 g of *gari* to 195 mL of boiled water (100 °C) and continuously stirred to form a smooth thick paste, while the *fufu* dough was prepared by adding about 200 g of *fufu* flour to 195 mL of boiling water (100 °C) while on the heat source and continuously stirred for two minutes to form a smooth thick paste.

2.5 Sensory Evaluation

Sensory evaluation was carried out on reconstituted *gari* and *fufu* (from *gari* and *fufu* flour). A twenty-man panelist familiar with *gari* and *fufu* was drawn from staff of NRCRI, Umudike and students of Michael Okpara University of Agriculture, Umudike. The criteria for selection was that panelist were 18 years of age, regular consumers of *gari* and *fufu* and were neither sick nor allergic to any of the foods. The panelists were trained in the use of sensory evaluation procedures. *Gari* and *fufu* samples were served on white disposable plates, properly coded with 3-digit random numbers to prevent bias. The reconstituted *gari* and *fufu* flours samples were assessed for aroma, taste, texture, colour, mouldability, and general acceptability. In each case, the samples were rated according to a 9-point hedonic scale of preference with ratings ranging from 1 (dislike extremely) and 9 (like extremely) as described by Iwe, [24]. Precautions were taken to avoid transfer of flavour during the analysis by ensuring that panelist rinse their mouth with potable water after each evaluation.

2.6 Statistical Analysis

Analyses were done in triplicate and all data obtained were subjected to Analysis of Variance (ANOVA) R- Statistics, (R- programming language version 3.4.4). Statistical significance was established using analysis of variance

(ANOVA) and means were separated using least significant difference (LSD) at $p \leq 0.05$.

3. RESULTS AND DISCUSSION

Table 1 shows the proximate composition of four cassava root varieties stored at ambient conditions for 14 days. Moisture content of cassava roots ranged from 50.05 to 70.12% across the storage period. TMS 07/0539 had the highest initial moisture content (70.12%) while NR 87/184 had the lowest initial moisture content (54.35%). All varieties showed a significant decrease in moisture content throughout storage (from Day 0 to Day 14). This decrease could be attributed to respiration and drying processes of the roots after harvest. This observation agrees with the reports of Omosuli et al. [25] who had moisture of 68.40% in fresh cassava (day 0) and 58.80% on day 10.

Commercial cassava varieties have poor protein of about 1-2% [26]. The protein content of cassava varieties varied from 0.42 to 1.10 % across the storage period (Day 0 to 14). Protein content of roots for all varieties increased from day 0 to 7, and then decreased on day 14. The increase in protein from day 0 to 7 might have been slightly enhanced by fermentation as reported by Tivana et al. [27]. All varieties showed a decrease in protein content by Day 14. The decrease in protein content of cassava varieties during storage could be due to enzymatic activity as cassava naturally contains enzymes that can degrade proteins over time [28]. Protein content obtained in this research is within range of 0.4 and 1.5 g/100 g fresh weight reported by Bradbury & Holloway, [29], Charles et al. [30].

Crude fiber content of cassava roots varied from 1.71 to 3.20 % across storage days. The white fleshed varieties NR87/184 and TME 419 had higher crude fibre values than the yellow fleshed varieties TMS 07/0593 and TMS 01/1368. Reports have shown that compared to yellow- and cream-flesh cassava varieties, the roots of white-flesh variants have a comparatively greater fiber content of 0.62-2.92 % [31]. Little increase in fiber content was observed for all varieties by day 7 and then decreased by day 14. The results obtained for this work is within the range of value obtained from results of 2.08 and 4.53 reported by Tivana et al. [27].

Cassava roots are known to have low fat content [26]. Fat content obtained in this study varied between 0.72 to 1.24 %. The fat content in roots

slightly decreased consistently across storage period for all varieties. The results of fat content obtained in this study are higher than results of 0.1 - 0.3 % reported by Bayata, [26]. This could be attributed to cassava varietal differences.

Ash content of cassava roots varied from 1.58 to 2.12 % across the storage period. There was a trend of consistent decrease of ash content during the storage period. This could be attributed to the fact that during the period of storage, cassava roots continue to respire during which they burn sugars and starches for energy, releasing carbon dioxide and water vapor. This process can lead to the breakdown of some minerals that contribute to the ash content. The results of ash content reported in this study are higher compared to those reported by Ikujenlola & Opawale, [32] who had ash content values between the range (0.60 – 1.30 %) and lower than ash content values of 2.29 to 2.67 % reported by Koko et al. [33]. The differences in ash content of cassava roots could be attributed to varietal differences.

Carbohydrates the dominant component was observed to increase throughout storage. This likely reflects the decrease in moisture content.

Tables 2 and 3 show the proximate composition of *fufu* flour and *gari* samples processed at day 0, 7 and 14 from stored cassava roots.

The moisture content of *fufu* flour and *gari* processed from the cassava varieties ranged from 7.44 % to 9.03 % and 4.12 to 12.15 % respectively. There was decrease in moisture content of both *fufu* flour and *gari* in all varieties throughout processing days. This confirms the moisture loss observed in roots during storage period of roots. Moisture content of cassava products determine to a large extent the ability of floury product to store well. The moisture content of *fufu* flour and *gari* were within acceptable limits of 12%. With this moisture content, *fufu* flour and *gari* processed from stored roots will store for a reasonably long period.

The protein content showed some variation across varieties and processing days. The protein content of *fufu* flour was in the range of 1.06 to 1.71 % while that of *gari* was in the range of 0.23 to 2.16 % across processing days. Generally, the protein content for both *fufu* flour and *gari* increased from day 0 to 7 and then decreased on day 14. This was observed also observed in the roots during storage. *Fufu* and *gari* are not protein-rich foods, but this variation might be relevant for dietary considerations.

Table 1. Proximate composition of cassava roots during storage

Variety	Storage Period (Days)	Moisture	Crude protein	Crude fibre	Fat	Ash	CHO
TMS 01/1368	0	68.42 ± 0.03	0.86 ± 0.01	2.04 ± 0.01	1.17 ± 0.02	2.11 ± 0.01	25.42± 0.01
	7	57.56± 0.03	0.91± 0.01	2.11 ± 0.01	1.03 ±0.01	2.05 ± 0.01	36.35 ±0.01
	14	54.75 ± 0.42	0.71 ± 0.01	1.76 ± 0.21	0.72 ± 0.01	1.71 ± 0.01	40.37± 0.01
TMS 07/0539	0	70.12 ± 0.04	0.97 ±0.01	1.93 ± 0.02	1.07 ± 0.01	1.89 ± 0.01	24.03± 0.01
	7	65.62 ± 0.02	1.10 ± 0.01	1.98 ± 0.01	0.98 ± 0.01	1.83 ± 0.02	28.50 ±0.01
	14	53.33± 0.42	0.75 ± 0.01	1.71 ± 0.01	0.79± 0.01	1.61 ± 0.01	41.82 ± 0.01
NR 87/184	0	54.35 ± 0.35	0.75 ± 0.07	3.05 ± 0.01	1.24 ± 0.01	2.12 ± 0.01	38.50± 0.08
	7	52.37 ± 0.03	0.88 ±0.01	3.20 ± 0.01	1.18 ± 0.01	2.06 ± 0.01	40.31 ± 0.00
	14	51.12 ± 0.02	0.51 ±0.01	2.27± 0.01	0.82 ± 0.01	1.79 ± 0.01	43.49 ± 0.01
TME 419	0	64.05 ±0.35	0.73 ± 0.02	2.58 ± 0.01	0.98 ± 0.01	2.02 ±0.01	30.10 ± 0.72
	7	60.45 ±0.04	0.78 ±0.01	3.14 ± 0.70	0.92 ± 0.03	1.85 ± 0.02	32.72 ± 0.70
	14	50.05± 0.04	0.42 ± 0.02	2.41 ± 0.01	0.85 ± 0.01	1.58± 0.71	44.45 ± 0.01
LSD		5.01	NS	NS	0.11	NS	4.52

*NS= Not Significant

Table 2. Proximate composition of fufu flour

Variety	Dayof processing	Moisture	Crude protein	Crude fibre	Fat	Ash	CHO
NR87/184	0	9.03 ± 0.03	1.66 ±0.02	2.38 ±0.01	0.63 ±0.01	1.52 ±0.01	84.79±0.01
	7	8.02 ±0.03	1.74 ±0.01	2.31 ±0.01	0.83 ±0.01	1.48 ±0.01	85.64±0.02
	14	7.44 ±0.03	1.06 ±0.01	2.05 ±0.02	0.37 ±0.03	1.31 ±0.04	87.67±0.05
TME419	0	8.72 ±0.03	1.51 ±0.01	2.45 ±0.02	0.58 ±0.02	1.51 ±0.01	85.25±0.01
	7	8.38 ±0.02	1.71 ±0.01	2.38 ±0.01	0.78 ±0.02	1.45 ±0.01	85.31±0.00
	14	8.02 ±0.03	1.12 ±0.01	2.02 ±0.01	0.43 ±0.01	1.22 ±0.02	87.21±0.00
TMS 01/1368	0	8.93 ± 0.03	1.47 ± 0.01	2.46 ±0.01	0.73 ±0.02	1.38 ±0.01	85.71±0.00
	7	8.24 ± 0.03	1.31 ± 0.01	2.61 ±0.01	0.61 ±0.01	1.27 ±0.03	85.29±0.02
	14	7.83 ± 0.03	1.16 ±0.02	2.18 ±0.01	0.44 ±0.01	1.27 ±0.01	87.13±0.00
TMS 07/0539	0	8.82 ±0.04	1.59 ±0.01	2.23 ±0.02	0.82 ±0.02	1.46 ±0.01	85.36±0.02
	7	8.57 ±0.03	1.28 ± 0.01	2.74 ±0.01	0.52 ±0.02	1.33 ±0.01	85.32±0.01
	14	7.87± 0.04	1.12 ±0.02	2.13 ±0.02	0.47 ±0.01	1.23 ±0.02	87.20±0.01
LSD		0.06	0.03	0.31	0.34	0.04	0.04

Table 3. Proximate composition of *gari*

Variety	Day of processing	Moisture	Crude protein	Crude fibre	Fat	Ash	CHO
NR87/184	0	9.64±0.01	0.52 ±0.01	2.06 ± 0.02	0.50 ±0.01	1.31 ±0.04	87.85±0.01
	7	4.21±0.02	2.08 ±0.01	1.21 ±0.01	0.70 ±0.02	1.52 ±0.01	89.80±0.01
	14	4.14±0.03	2.05 ±0.01	2.31 ±0.01	0.82 ±0.01	1.48 ±0.01	89.66±0.00
TME419	0	10.05 ±0.03	0.57 ±0.01	0.72 ±0.02	0.54 ±0.01	0.62 ±0.01	87.50±0.01
	7	4.56 ±0.01	2.16 ±0.07	1.22 ±0.01	0.74 ±0.01	2.15 ±0.01	89.18±0.42
	14	4.12 ±0.01	2.01 ±0.01	1.22 ±0.01	0.76 ±0.01	1.89 ±0.01	90.02 ±0.04
TMS 01/1368	0	10.36±0.02	0.49 ±0.01	0.73 ±0.02	0.62 ±0.02	0.78±0.01	87.05±0.02
	7	4.47 ±0.03	1.93 ±0.01	1.12 ±0.01	0.78 ±0.13	1.90 ±0.01	90.07±0.01
	14	4.22 ±0.02	2.11 ±0.01	1.12 ±0.02	0.85 ±0.02	1.96 ±0.01	89.51 ±0.03
TMS 07/0539	0	12.15±0.05	0.23 ±0.03	0.81 ±0.01	0.37 ±0.04	0.38 ±0.03	86.07±0.01
	7	4.86 ±0.02	2.06 ±0.01	1.18 ±0.02	0.86 ±0.1	2.02 ±0.01	89.04±0.01
	14	4.53 ±0.03	2.14 ±0.02	1.16 ±0.02	0.91 ±0.01	2.03 ±0.01	89.25±0.04
LSD		0.06	0.03	0.04	0.04	0.03	0.05

Fat content of *fufu* flour and *gari* were in the ranges 0.37 to 0.83 % and 0.37 to 0.91 %. Fat content of *fufu* flour and *gari* remained fairly stable across processing days.

Crude fiber content varied significantly ($p \geq 0.05$). The crude fibre content of *fufu* flour was in the range 2.02 to 2.74 % while the crude fibre content of *gari* was in the range 0.72 to 2.06 %. The Codex Alimentarius maximum values for crude fibre of edible flour and *gari* is 2 % [10]. The crude fibre content values of *gari* obtained in this study were within the codex Alimentarius maximum value while *fufu* flour crude fibre values were slightly above the standards. Ash content values of *fufu* flour and *gari* varied significantly ($p \geq 0.05$). The ash content values of *fufu* flour and *gari* obtained in this study are within 3 % Codex Alimentarius standards. Carbohydrates content had values ranging from 84.79% to 87.67% for *fufu* flour and 86.07 to 90.07 %. There was a general increase in carbohydrate content of *fufu* flour for all varieties across the period of storage. The increase in carbohydrate content likely reflects the decrease in moisture content during storage.

Table 4 shows the results of the sensory evaluation of reconstituted *fufu* at days (0, 7, and 14) from stored cassava varieties. Scores ranged from 1 (disliked extremely) to 9 (liked extremely) for appearance, aroma, texture, and general acceptability. All varieties showed a decrease in hedonic scores (sensory liking) as storage time increased (from Day 0 to Day 14). This suggests a decline in *fufu* quality during storage. At (Day 0) scores for most varieties fell between "liked

slightly" (6) and "neither liked nor disliked" (5), indicating a more neutral initial perception. Scores decreased further towards "disliked slightly" (4) or even "disliked moderately" (3) by Day 14 for most sensory attributes. Variety NR87/184 maintained the highest hedonic scores throughout storage. TME419 showed a significant decline, especially after Day 7. By Day 14, most attributes scored below "neither liked nor disliked" (5), indicating a clear negative perception. This suggests a very short shelf life for TME419 *fufu* based on consumer preference. TMS01/1368 and TMS07/0539 had the lowest hedonic scores across all storage days, with scores dropping to "disliked moderately" (3) or even "disliked very much" (2) by Day 14 for some aspects. This indicates that consumers generally disliked these varieties, especially after storage.

The results for sensory evaluations of *gari* prepared from cassava varieties stored for 0, 7 and 14 days are presented in Table 5. It was observed that storage of cassava roots of varieties NR 87/184, TME 419 and TMS 01/1368 for up to 14 days gave acceptable *gari* characteristics above 5 points (hedonic scale) in terms of colour, aroma, texture and general acceptability, though the colour for TMS 01/1368 was below 5. Variety TMS 07/0953 gave acceptable *gari* characteristics only after 7 days of roots storage. The results also showed that as the period of cassava roots storage progressed, there was a gradual decrease in scores the of the quality attributes. Nevertheless, the scores were within the acceptable range for a good quality *gari* characteristics.

Tale 4. Sensory qualities of reconstituted *fufu* flour

Variety	Day of Processing	Appearance	Aroma	Texture	General Acceptability
NR87/184	0	6.60	5.70	5.60	6.00
	7	5.70	5.10	5.20	6.00
	14	5.80	5.60	5.80	6.20
TME419	0	6.20	6.20	6.30	6.60
	7	3.90	5.30	5.10	5.10
	14	2.50	2.60	4.30	3.50
TMS01/1368	0	6.50	5.90	5.90	6.50
	7	4.80	5.70	4.80	5.70
	14	5.60	5.80	3.30	5.50
TMS07/0539	0	5.60	5.30	4.50	5.40
	7	3.80	4.70	4.90	5.00
	14	1.90	4.20	1.90	2.90
LSD		0.70	0.70	0.80	0.70

Table 5. Sensory qualities of reconstituted gari

Variety	Days of Processing	Appearance	Aroma	Taste	Texture	General Acceptability
NR87/184	0	7.40	7.20	6.80	7.70	7.30
	7	6.30	7.40	7.30	7.60	7.80
	14	6.40	6.40	5.60	6.40	6.40
TME419	0	7.70	7.90	7.40	7.90	7.90
	7	6.60	6.60	7.10	7.30	6.80
	14	5.50	5.40	5.40	5.50	5.70
TMS01/1368	0	7.30	6.90	7.00	7.20	7.30
	7	3.90	5.80	5.80	6.20	5.90
	14	5.90	6.40	6.40	6.30	6.30
TMS07/0539	0	7.20	7.10	7.00	7.40	7.50
	7	6.10	6.30	5.80	5.20	6.30
	14	2.50	2.60	5.00	4.10	4.40
	LSD	0.50	0.50	0.40	0.50	0.40

4. CONCLUSION

The analysis of cassava roots, *gari*, and *fufu* revealed that cassava roots are a carbohydrate-rich food source with moderate protein content and low fat content. During ambient storage, the roots experienced significant moisture loss, increasing carbohydrate content being the dominant nutrient of the roots.

Gari and *fufu* are processed cassava products with similar nutritional profile to the roots, but have slightly lower moisture contents due to processing. Protein contents of both *fufu* flour and *gari* showed some variations across varieties, which might influence their overall nutritional value. Proximate composition remained fairly stable across varieties, with carbohydrates being the dominant component.

Sensory evaluation scores for *fufu* and *gari* across varieties showed a decrease in hedonic scores (sensory liking) as storage time increased (from Day 0 to Day 14). This suggests a decline in *fufu* and *gari* quality during storage. Based on the scale (1-9), most initial scores (Day 0) fell between "liked slightly" (6) and "liked moderately" (7), indicating a generally favorable initial perception for all varieties. However, scores decreased towards "neither liked nor disliked" (5) by Day 14 for most sensory attributes. Variety NR87/184 maintained relatively high hedonic scores throughout storage compared to others. This suggests this variety might have a longer shelf life or better storage characteristics. TMS07/0539 had the most significant decline in hedonic scores across storage, with all attributes dropping below "neither liked nor disliked" (5) by Day 14. This

suggests a rapid decrease in consumer liking for *fufu* and *gari* processed from this variety.

5. RECOMMENDATIONS

Alternative storage methods can be explored to minimize moisture loss and extend shelf life while maintaining cassava root quality. Further work could be done to investigate the relationship between *fufu* and *gari* proximate composition and sensory characteristics preferred by consumers.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. LokkoY, Anderson J, Rudd S, Raji A, Horvath D, Mikel M, Dixon A. Characterization of an 18,166 EST dataset for cassava (*Manihot esculenta* Crantz) enriched for drought-responsive genes. *Plant cell reports*.2007;26:1605-1618.
2. OishimayaN. Top cassava producing countries in the world. *World Atlas*. In; 2017.
3. Rahmawati R, SukmaD, Ardie S, Sudarsono S. Postharvest physiological deterioration in cassava: Potential

- problems, possible inhibition, and resistant level identification. IOP Conference Series: Earth and Environmental Science; 2021,
4. ZainuddinIM, Fathoni A., Sudarmonowati E, Beeching JR, Gruissem W, Vanderschuren H. Cassava post-harvest physiological deterioration: From triggers to symptoms. *Postharvest Biology and Technology*.2018;142:115-123.
 5. FadeyibiA, Osunde Z, BelloK. Measures against damage of some perishable products on transit. *Advances in Agriculture, Sciences and Engineering Research*.2011;1(1):1-8.
 6. Uchechukwu-AguaAD, CalebOJ, Opara UL. Postharvest handling and storage of fresh cassava root and products: A review. *Food and Bioprocess Technology*.2015a;8:729-748.
 7. Bamidele OP, Fasogbon MB, Oladiran DA, Akande EO. Nutritional composition of fufu analog flour produced from Cassava root (*Manihot esculenta*) and Cocoyam (*Colocasia esculenta*) tuber. *Food science & nutrition*. 2015;3(6):597-603.
 8. Uchechukwu-Agua AD, CalebOJ, OparaUL. Postharvest handling and storage of fresh cassava root and products: A review. *Food and Bioprocess Technology*, 2015b;8(4):729-748.
 9. Bamidele O, FasogbonB, Oladiran DO, Akande E. Nutritional composition of fufu analog flour produced from Cassava root (*Manihot esculenta*) and Cocoyam (*Colocasia esculenta*) tuber. *Food Science & Nutrition*.2015;3. Available:<https://doi.org/10.1002/fsn3.250>
 10. AbassAB, Dzedzoave NT, AlenkheBE, James BD. Quality management manual for the production of gari. In: *International Institute of Tropical Agriculture*; 2013.
 11. Bechoff A, Tomlins K, FliedelG, Becerra Lopez-lavalle LA, Westby A, HersheyC, DufourD. Cassava traits and end-user preference: Relating traits to consumer liking, sensory perception, and genetics. *Critical Reviews in Food Science and Nutrition*.2018;58(4):547-567.
 12. Owolarafe O, Adetifa B, Oyekanmi A, Lemikan T, Samuel T. Assessment of fufu production technologies in Ogun State, Nigeria. *Arid Zone Journal of Engineering, Technology and Environment*. 2018;14(4):547-558.
 13. Ume S, Okoye F, Onwujiariri U, Achebe U. Analysis of Intensity of Adoption of Odourless Fufu Processing Technology by Pro vitamin A Cassava Variety Processors in Anambra State of Nigeria; An Implication to Health and Nutritional Food Security. *International Journal of Science and Healthcare Research*.2020;5(7):34-45.
 14. Oyeyinka SA, Ajayi OI, Gbadebo CT, Kayode RM, Karim OR, Adeloye AA. Physicochemical properties of gari prepared from frozen cassava roots. *LWT*.2019;99:594-599.
 15. SanchezT, Dufour D, Moreno JL, Pizarro M, Aragón I, Dominguez M, Ceballos H. Changes in extended shelf life of cassava roots during storage in ambient conditions. *Postharvest Biology and Technology*.2013;86:520-528.
 16. Tomlins K, Parmar A, Omohimi CI, Sanni LO, Adegoke AF, AdebowaleA-RA, Bennett B. Enhancing the shelf-life of fresh cassava roots: A field evaluation of simple storage bags. *Processes*.2021;9(4):577.
 17. UarrotaVG, NunesEdC, PeruchLAM, NeubertEdO, Coelho B, Moresco R, Dufour D. Toward better understanding of postharvest deterioration: Biochemical changes in stored cassava (*Manihot esculenta* Crantz) roots. *Food Science & Nutrition*. 2016;4(3):409-422.
 18. Akingbala JO, Oyewole OB, Uzo-Peters PI., Karim RO, Baccus-TaylorGS. Evaluating stored cassava quality in gari production. *Journal of Food, Agriculture & Environment*.2005;3(1):75-80.
 19. Idowu M, Akindele S. Effect of storage of cassava roots on the chemical composition and sensory qualities of gari and fufu. *Food chemistry*.1994;51(4):421-424.
 20. Venturini MT, SantosVdS, OliveiraEJd. Procedures for evaluating the tolerance of cassava genotypes to postharvest physiological deterioration. *Pesquisa Agropecuária Brasileira*, 2015;50:562-570.
 21. Aniedu C, Omodamiro R. Use of newly bred β -carotene cassava in production of value-added products: Implication for food security in Nigeria. *Global Journal of Science Frontier Research Agriculture and Veterinary Sciences*.2012;12(10):1-10.
 22. Horwitz W, Chichilo P, ReynoldsH. Official methods of analysis of the Association of Official Analytical Chemists. *Official methods of analysis of the Association of Official Analytical Chemists*; 1970.
 23. UdoroE, Kehinde A, OlasunkanmiS, Charles T. Studies on the physicochemical, functional and sensory properties of gari

- processed from dried cassava chips. Journal of Food Processing and Technology.2014;5(1):293.
24. IweM. Proximate, Physical And Sensory Properties Of Soy-Sweet Potato Flour Cookie. Global Journal of Pure and Applied Sciences.2002;8(2);187-192.
 25. Omosuli SV, IkujenlolaAV, AbisuwaAT. Quality assessment of stored fresh Cassava Roots and 'fufu'flour produced from stored roots. J Food Sci Nutr The. 2017;3(1):009-013.
 26. Bayata A. Review on nutritional value of cassava for use as a staple food. Sci J Anal Chem, 2019;7(4):83-91.
 27. Tivana L, Bvochora J, Mutukumira A, Owens J. A study of heap fermentation process of cassava roots in Nampula Province, Mozambique. Journal of Root Crops. 2007;33(2):119-128.
 28. Falade KO, Akingbala JO. Utilization of cassava for food. Food Reviews International.2010;27(1):51-83.
 29. Bradbury JH, Holloway WD. Chemistry of tropical root crops: significance for nutrition and agriculture in the Pacific; 1988.
 30. Charles AL, Sriroth K, Huang T-C. Proximate composition, mineral contents, hydrogen cyanide and phytic acid of 5 cassava genotypes. Food chemistry. 2005;92(4):615-620.
 31. Ukenye E, UkpabiU, ChijokeU, EgesiC, Njoku S. Physicochemical, nutritional and processing properties of promising newly bred white and yellow fleshed cassava genotypes in Nigeria. Pakistan Journal of Nutrition.2013;12(3):302.
 32. IkujenlolaA, OpawaleBO. Effects of processing on the yield and physico-chemical properties of cassava products. Advanced Materials Research; 2007.
 33. KokoCA, KouameBK., BlanchardYA, Amani GNg, Assidjo EN. Comparative study on physicochemical characteristics of cassava roots from three local cultivars in Côte d'Ivoire. European Scientific Journal. 2014;10(33).

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/118329>