



Assessment of the Knowledge, Attitude and Practice of Cervical Cancer Screening among Students at Ambrose Alli University in Southern Nigeria

Okpu U. Jeremiah ^{a*}, Uhomoibhi D. Osezimen ^b,
Ayemoro O. Temitope ^c, Okeke O.Daniel ^d,
Madu, W. Chimezie ^e, Ohaneneye M. Chinedu ^f,
Eze O.Samuel ^g, Aturamu A. Ayoyinka ^h,
Ugwoke O. Albert ⁱ, Ugwu E. Francis ^j, Okolo E. Ifeanyi ^k,
Idialu .O.Perpetual ^l, Dongo E. Naomi ^m,
Akhaine J. Precious ⁿ and Ezugwu O. Oliver ^o

^a Department of Obstetrics and Gynecology Irrua Specialist Teaching Hospital, Edo State, Nigeria.

^b Department of Cancer Biology, George Town University DC USA.

^c Department of Obstetrics and Gynecology, Central Hospital Edo State, Nigeria.

^d Department of Primary Health Care, Bradford Teaching Hospitals NHS Foundation Trust, United Kingdom.

^e General Practice, Bradford Teaching Hospitals NHS Foundation Trust, United Kingdom.

^f Department of Internal Medicine, University of Nigeria Teaching Hospital Ituku, Ozalla, Enugu State, Nigeria.

^g Department of Internal Medicine Federal Medical Center Owerri, Imo State, Nigeria.

^h Department of Internal Medicine, Ekiti State University Teaching Hospital, Ado Ekiti, Ekiti State

^l Department Pathology, University of Nigeria Teaching Hospital Ituku, Ozalla, Enugu State, Nigeria.

^j Department of Surgery, Federal Medical Centre, Owerri, Imo State, Nigeria.

^k Department of Internal Medicine, Delta State University Teaching Hospital, Nigeria.

^l Department of Internal Medicine, Irrua Specialist Teaching Hospital, Irrua Edo State, Nigeria.

^m Department of Public Health, Irrua Specialist Teaching Hospital, Irrua Edo State, Nigeria.

ⁿ Department of Obstetrics and Gynecology, Central Hospital Edo State, Nigeria.

^o Department Surgery, University of Nigeria Teaching Hospital Ituku, Ozalla, Enugu State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors OUJ, UDO and DEN conceived and designed the study. Authors AOT, OOD, MWC and OMC contributed to data acquisition and sorting. Data analysis and interpretation was carried out by authors AJP, EOS, AAA and UOA. First draft of the manuscript was written by authors OUJ, OEI, IOP, DEN and EOO. All authors critically reviewed the manuscript for important intellectual content. All Authors read and approved the final manuscript.

*Corresponding author: E-mail: okpujeremiah@gmail.com;

Cite as: Jeremiah, Okpu U., Uhomoibhi D. Osezimen, Ayemoro O. Temitope, Okeke O.Daniel, Madu, W. Chimezie, Ohaneneye M. Chinedu, Eze O.Samuel, Aturamu A. Ayoyinka, Ugwoke O. Albert, Ugwu E. Francis, Okolo E. Ifeanyi, Idialu .O.Perpetual, Dongo E. Naomi, Akhaine J. Precious, and Ezugwu O. Oliver. 2024. "Assessment of the Knowledge, Attitude and Practice of Cervical Cancer Screening Among Students at Ambrose Alli University in Southern Nigeria". Asian Research Journal of Gynaecology and Obstetrics 7 (1):307-25. <https://journalarjgo.com/index.php/ARJGO/article/view/239>.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/125077>

Original Research Article

Received: 25/08/2024
Accepted: 27/10/2024
Published: 04/11/2024

ABSTRACT

Background: Cervical cancer is a malignant condition affecting the cervix, with an average onset age of 45 years. It is the second most common cancer among women globally as in 2012, over 550,000 new cases of cervical cancer were diagnosed worldwide; of these, a large majority, about 85% occurred in less developed regions yet it is highly preventable if the premalignant stage is discovered through early routine screening.

Objectives: The objective of this study was to determine the knowledge, attitude and practice of cervical cancer screening among Ambrose Alli University student, Ekpoma, Edo state.

Methodology: Descriptive cross-sectional study was conducted among 326 female students of Ambrose Alli University using semi-structured interviewer administered questionnaire and multistage sampling technique was employed which helped to ensure flexibility of the study. Data was analyzed using IBM SPSS version 23.0 software for descriptive statistics.

Results: All respondents were female and it was observed that, 24.8% of respondents had good knowledge, 28.2% had fair knowledge, while 46.9% of respondents had poor knowledge. Knowledge increased with increasing age with highest knowledge (63.6%) among 29 -31 years age group. Knowledge also increased as observed in those in the higher university class. The attitude of the participants toward cervical cancer screening was poor. It was seen that 18.7% had positive attitude toward cervical cancer screening. Half of the respondents who had good knowledge also had good attitude toward cervical cancer screening. The practice of screening exercise by respondents was also found to be poor as 1.5% of the respondents had ever done Pap smear at least once in the past.

Conclusion: From this study, the knowledge, attitude and practice of cervical cancer screening among students of Ambrose Alli University were low which can have a negative impact in the general public on the long run as these girls can come down with cervical cancer and its complications.

Keywords: Cervical cancer; pap smear; knowledge; attitude; practice.

ABBREVIATIONS

FMOH	: Federal Ministry of Health
HPV	: Human Papilloma Virus
Pap Smear	: Papanicolaou Smear
WHA	: World Health Assembly
WHO	: World Health Organization

1. INTRODUCTION

1.1 Background

“Cervical cancer is a cancerous preventable state that affects the uterus especially the cervix,

with an average onset age of 45 years. It often has a pre-cancerous stage, which typically develops in younger women under the age of 40” [1]. “The major cause of cervical cancer is infection with one or more high-risk strains of the human papillomavirus (HPV), a common sexually transmitted virus which indicates that cervical cancer can be preventable” [2]. “The availability of effective screening options such as Pap smear, human papilloma virus-deoxyribonucleic acid testing and visual inspection with application of acetic acid and iodine makes it quite easy to comprehend” [3]. “Cervical cancer is associated with the following risk factors; early age at first

sexual intercourse between 16-18 years, multiple male sexual partners, male sexual partners who have had multiple partners, early age at first birth, multi-parity, smoking, long-term use of oral contraceptive pills, immunosuppressed states, HIV, poverty, family history of cervical cancer, lack of regular screening" [1].

"Cancer of the cervix is a preventable disease when detected and treated at early stage and a vital aspect of its prevention is the detection of the pre malignant form by cervical screening, and if detected in time, recovery is achievable" [1]. As a result of the long transition time from a premalignant lesion to frank cancer of the cervix, it gives enough time for early detection and achieve nearly complete cure even in secondary health care centers [1]. However, in developing countries it has been noticed that there is difficulty in implementing cytology-based screening programs [4]. "It has been noted that the incidence of cervical cancer over the past decades has decreased in developed countries, as it was reported that over 270,000 women died of cervical cancer worldwide; almost 9 out of every 10 of these, or 231,000 women in total, lived and died in low- to middle income countries" [5]. To this effect, it is emphasized that increase in awareness and more effective screening and prevention strategies employed in these countries [6]. Furthermore, the human papilloma virus vaccine has contributed to a decline in the incidence rate of cervical cancer [7]. "Although screening is a well-known and cost-effective strategy for reducing the burden of cervical cancer globally, its implementation remains significantly low, especially in developing countries" [8]. "It has been reported that challenges such as limited healthcare service, language and communication barriers, cultural and social barriers, fear and anxiety, lack of regular screening program but the major barrier to access is that most cervical cancer screening services (provided by governmental and nongovernmental agencies) in Nigeria had been sporadic and poorly coordinated" [9].

"Despite efforts by governmental and non-governmental organizations to enhance access to cervical cancer screening services in Nigeria, the uptake has been dismal. Various studies have identified factors influencing the practice of cervical screening tests globally. These factors include a woman's age, marital status, number of children, risk perception, financial constraints, and knowing someone with cervical cancer" [10]. Furthermore, "the cost of screening, which can be as high as twenty thousand naira (20-30

USD) may contribute to the low participation rate. In a country like Nigeria, where a significant portion of the population lives below the poverty line and healthcare is largely funded through out-of-pocket expenses, this cost can be prohibitive for many individuals" [11].

Globally, cervical cancer is the fourth most common cancer in women, with around 660 000 new cases in 2022. In the same year, about 94% of the 350 000 deaths caused by cervical cancer occurred in low- and middle-income countries. The highest rates of cervical cancer incidence and mortality are in sub-Saharan Africa (SSA), Central America and South-East Asia [12].

Cervical cancer is the second most common cancer both among women and in the general population in Nigeria. Of the 14,000 women who develop the disease annually, over half of them will die. In Nigeria, the Cervical Cancer Age-standardized Incidence Rate in 2018 was 27.2 per 100,000 women while the Cervical Cancer Age-standardized Mortality Rate was 20 per 100,000 women [13].

It has been reported that approximately 50% of invasive cervical malignancies are diagnosed in patients who have never been screened and that 10% of the remaining cervical cancer patients have not had a Pap smear in the 5 years before diagnosis [14]. A previous study from Zaria noted that only 270 patients were screened as part of routine screening in 5 years [15]. "The consequences of the increasing prevalence of cervical cancer include increasing morbidity and mortality of women due to an otherwise preventable disease. Nigeria has not had a great deal of success in implementing effective cervical cancer screening till date. In addition to presenting a significant burden in terms of morbidity and mortality, cervical cancer also increases economic risk, which imposes very high direct costs on health systems, communities, households and lost the productivity of patients, premature death, and disability" [16]. "Loosing a mother is devastating for a family, but even when the disease is not fatal, the consequences are severe. Surgical and radiological treatment of cervical cancer often leads to physical, psychological, economic and sexual issues as well as infertility. These debilitating outcomes warrant policies to reduce the prevalence of cervical cancer" [17].

The study was carried out among female students of 17 years and above in Ambrose Alli

University to determine their level of understanding of cervical cancer screening. It will be an effective means in creating and disseminating information about the importance of the screening programs available. The emphasis on cervical cancer screening at this point in time is imperative because of the continuous increase in cervical cancer cases in this environment as opposed to that in the developed world which is largely attributed to the widespread practice of cervical cancer screening among their women. The study intends to achieve a reduction in the incidence of cervical cancer in this environment by increasing women's knowledge and hence practice of cervical cancer screening.

The researchers aimed to determine the knowledge of cervical cancer screening among female students of Ambrose Alli University. The researcher also investigated the attitude of female students of Ambrose Alli University towards cervical cancer screening and to assess the practice of cervical cancer screening among participants.

2. METHODOLOGY

2.1 Study Area

The study was done in Ambrose Alli University, Ekpoma, Edo state. It is accredited and recognized by the National University Commission. It has several faculties and departments that are accredited and render educational services to its students and prospective students. The population currently covers an estimate of about 27,000 regular undergraduate students [18].

2.2 Study Design

A descriptive cross-sectional study design was used for the study.

2.3 Study Population

The study population targeted was the female students of Ambrose Alli University Ekpoma, Edo state, Nigeria.

2.3.1 Inclusion criteria

The study included only Female undergraduate students of Ambrose Alli University who consented to the study.

2.3.2 Exclusion criteria

Under graduate female from the Ambrose Alli University who were unwilling to participate in the study from 17years and above.

2.4 Study Duration

The study was done over a timeline of 6months from March to September 2021.

2.5 Sample Size Determination

The sample size was calculated using the Cochran formula for simple proportion [19]:

$$n = \frac{z^2 pq}{d^2}$$

Where:

n = minimum sample size

z = standard normal deviate (reliability coefficient at 95% confidence interval; standard value of 1.96)

d = degree of accuracy or margin of error at 5% (standard value is 0.05)

p = prevalence i.e. proportion of population with characteristics of interest = 73.9% (0.739)
Prevalence of knowledge of cervical cancer screening of 73.9% from a study done in Ibadan [20], was taken.

q = 1 – p

The sample size was thus estimated at:

$$n = \frac{1.96^2 \times 0.739 \times (1 - 0.739)}{(0.05)^2}$$

$$= \frac{3.8416 \times 0.739 \times 0.261}{0.0025}$$
$$= 296.$$

10% attrition was added for non-responses 10% of 296 = 29.6

Final sample size = 296 + 29.6 = 325.6 = 326.

2.6 Sampling Technique

Multistage sampling method was employed in the selection of the participants for the study using Interviewer-administered questionnaire.

First stage: This was done using simple random sampling method- balloting. The names of the various 14 faculties across all campuses of

Ambrose Alli University, Ekpoma were written on pieces of papers, folded and placed in an enclosed container, shaken together and then four faculties were selected. The faculties selected were faculty of clinical science, education, basic medical science and faculty of life science.

Second stage: This involved another simple random sampling were by the names of 31 departments from the selected four faculties were written on pieces of papers, folded and placed in four enclosed container based on the respective faculties. Each container was shaken and then one department was selected from each container. The selected departments were medicine and surgery, health education, anatomy and microbiology department.

Third stage: Involved a Stratified sampling, in which there was proportional allocation of students from selected departments.

2.7 Medicine and Surgery

Class	Population
100	350
200	325
300a	250
300b	300
400	175
500a	80
500b	125
Total	1605

2.8 Health Education

Year of study	Population
100 level	395
200 level	387
300 level	375
400 level	290
Total	1447

2.9 Microbiology

Year of study	Population
100 level	400
200 level	323
300 level	280
400 level	190
Total	1193

2.10 Anatomy

Year of study	Population
100 level	348
200 level	345
300 level	338
400 level	324
Total	1355

Sum Total = Medicine and Surgery (100-500 level) + Health Education (100-400Level) + Anatomy (100-400 level) + Microbiology (100-400 Level)

Sum Total = 1605 + 1447 + 1193 + 1355 = 5600
 The study population was 5600. The sample size of 326 was calculated from this as follows:
 Medicine and Surgery = $(1605 / 5600) \times 326 = 93$
 Health Education = $(1447 / 5600) \times 326 = 84$
 Anatomy = $(1193 / 5600) \times 326 = 69$
 Microbiology = $(1355 / 5600) \times 326 = 80$

Students in each department were randomly selected and given the questionnaire.

2.11 Tools for Data Collection

Data was collected using semi structured questionnaires (both closed and open ended) designed based on the specific objectives of the study. The questions were aimed to gather information regarding respondent's knowledge on screening for cervical carcinoma, their attitude and practice of cervical cancer screening. The questionnaire was also designed to obtain relevant socio-demographic characteristics of the respondents. The questionnaire was adapted from a previous study on knowledge, attitude and practice of cervical cancer screening amongst market women of Yola metropolis carried out in Adamawa State [21]. See attached in appendix.

2.12 Data Analysis

Descriptive statistics was conducted and data was entered into and analyzed with IBM SPSS software version 23.0. Proportions were derived for categorical variables while means and standard deviations were derived for continuous variables. The dependent variables were the socio-demographic factors while the outcome variables were knowledge and practice of cervical cancer screening. Knowledge, attitude and practice of cervical cancer screening were assessed by scoring the responses to the questions in section two, three and four

respectively in the questionnaire. Section two had a total of 15 questions beginning from question 7 and ending at question 21. A score of one was awarded to every correct answer except for question 8 which had no score because it was used to access the source of information of the respondents in regards to the knowledge of Pap smear. The highest obtainable score in this section was 14. Participants who scored below 6 had poor knowledge; participants who scored between 6-10 had fair knowledge while participants who scored above 10 had good knowledge of cervical cancer screening.

Section three which accessed the attitude of respondents toward cervical cancer screening had a total of 12 questions, beginning from question 22 and ending at question 33. Each question was awarded two marks and one mark respectively for every correct answer of “strongly agree” and “agree” or “strongly disagree” and “disagree”. The maximum obtainable score in section three was 24. Respondents who scored below 13 have negative attitude while those that scored 13 and above, have positive attitude toward cervical cancer screening.

Section four had a total of 6 questions, beginning from question 34 and ending at question 39. Questions 34 and 35 were awarded 2 marks each for each correct answer. For question 35 which accessed the number of times participants went for screening, had a maximum score of 3. A score of 1, 2 and 3 respectively was awarded for once, twice and multiple times while questions 36 to 38 had no score as they were used to access the influence on participants’ decision to go for pap smear, findings of the pap smear and action of the participants if pap smear was abnormal. The maximum obtainable score in this section is 7. Respondents who scored below 3 have poor practice while respondents who scored 3 and above, have good practice of cervical cancer screening.

3. RESULTS

Table 1 shows the demographic characteristics. Age of participants ranged from 17-31 with mean age 23.7±3.3. Most of the participants (35%) fall in the range of 23-25 years. Majority of them were single (74.8%). The dominant ethnic group was Esan ethnic group (38.7%), while majority (85.3%) of the respondents were Christians. The ethnicity was dominated by Esan ethnic group (38.7%), majority (85.3%) of the respondents were Christians while the highest number of

respondents were from year 3 (27.3%) and 4 (27.6).

Table 1. Socio-demographic characteristics of respondents

Variable	Frequency (n=326)	Percent
Age Group (in years)		
17 – 19	24	7.4
20 – 22	91	27.9
23 – 25	114	35
26 – 28	86	26.4
29 – 31	11	3.3
Mean ± S.D	23.7 ± 3.3	
Marital Status		
Single	244	74.8
Married	82	25.2
Ethnic Group		
Esan	126	38.7
Bini	90	27.6
Etsako	47	14.4
Yoruba	38	11.7
Igbo	20	6.1
Owan	5	1.5
Level		
100	28	8.6
200	87	26.7
300	89	27.3
400	90	27.6
500	32	9.8
Religion		
Christianity	278	85.3
Islam	48	14.7
Department		
Medicine	94	28.8
Anatomy	84	25.7
Health	68	20.9
Education		
English	80	24.5
Education		

Table 2. Respondents’ source of information about pap smear

Source of Information	Frequency (n=102)	Percent (%)
School	49	48.0
Health Worker	29	28.4
Social Media	16	15.7
Mass Media	8	7.8

Out of the 326 respondents, 102 (31.3%) have heard of Pap smear. Table 2 above shows the source of information of the respondents. School had the highest percentage (48.0%) while mass media (7.8%) was the list source of information.

Table 3. Knowledge of Ambrose Alli University students on cervical cancer and cervical cancer screening

Questions	Responses			
	Yes (%)	No (%)	I don't know	Total (%)
Heard of cancer of cervix	299 (91.7)	27 (8.3)	0 (0.0)	326 (100)
Heard of pap smear	102 (31.3)	224 (68.7)	0 (0.0)	326 (100)
Pap smear is the most helpful way to detect pre-cancer and cancer of the cervix	85 (26.1)	90 (27.6)	151 (46.3)	326 (100)
Women should have pap smear at least every 3years	136 (41.7)	48 (14.7)	142 (43.6)	326 (100)
The purpose of pap smear is to detect abnormal cells in the cervix	179 (54.9)	32 (9.8)	115 (35.3)	326 (100)
Pap smear is able to detect all types of female genital cancers	48 (14.7)	130 (39.9)	148 (45.4)	326 (100)
Pap smear is non-invasive and relatively inexpensive	71 (21.8)	120 (36.8)	135 (41.4)	326 (100)
Women should have pap smear since onset of sexual activity	151 (46.3)	27 (8.3)	148 (45.4)	326 (100)
Pap smear should be performed in both menstrual and non-menstrual periods.	58 (17.8)	120 (36.8)	148 (45.4)	326 (100)
A woman should not have sex 24hours before having pap smear	123 (37.7)	48 (14.7)	155 (47.5)	326 (100)
Pap smear should be discontinued after menopause	59 (18.1)	130 (39.9)	137 (42.0)	326 (100)
A patient with a normal Pap smear does not need pap smear in the future	43 (13.2)	162 (49.7)	121 (37.1)	326 (100)
There is no need to have pap smear if it is not administered by a doctor	196 (60.1)	58 (17.8)	72 (22.1)	326 (100)




Fig. 1. Pie chart showing the distribution of general knowledge of cervical cancer and cancer screening of participants

Table 4. Knowledge and socio-demographic characteristics of respondents

Socio-Demographic Characteristics.	Knowledge of respondents			Chi-Square	P-Value
	Good (%)	Fair (%)	Poor (%)		
Age Group (years)					
17 – 19	4 (16.7)	2 (8.3)	18 (75.0)		
20 – 22	11 (12.1)	20 (22.0)	60 (65.9)		
23 – 25	31 (27.4)	39 (34.5)	43 (38.1)		
26 – 28	27 (31.4)	32 (37.2)	27 (31.4)		
29 – 31	9(63.6)	2 (18.2)	2 (18.2)	154.495	<0.0001*
Marital status					
Single	58 (23.7)	61 (25.0)	125 (51.3)		
Married	22 (26.8)	32 (40.2)	28 (33.0)	8.282	0.016
Religion					
Christian	70 (25.2)	77 (27.7)	131 (47.1)		
Muslim	11 (22.9)	15 (31.3)	22 (45.8)	0.221	0.895
Level					
100	4 (14.3)	1 (3.6)	23 (82.1)		
200	5 (5.7)	38 (43.7)	44 (50.6)		
300	12 (13.5)	17 (19.1)	60 (67.4)		
400	37 (41.1)	32 (35.6)	21 (23.3)		
500	22 (68.8)	7 (21.8)	3 (9.4)	117.910	<0.0001*
Department					
Medicine	5)	7)	8)		
Anatomy)	2)	0)		
Health education	2)	5)	3)		
Microbiology)	2)	3)	0	1*

*=Significant ($p < 0.05$)

Table 5. Attitude of respondents towards cervical cancer screening

Questions	Frequency (n=326)			
	Strongly agree (%)	Agree (%)	Disagree (%)	Strongly disagree (%)
Would feel embarrassed if a male doctor performed the Pap smear	120 (36.8)	80 (24.5)	37 (11.3)	89 (27.3)
Would go for screening even if single	98 (28.5)	7 (2.1)	8 (2.5)	218 (66.9)
Being married will prevent you from going for screening	258 (79.1)	21 (6.4)	12 (3.7)	35 (10.7)
Thinks pap smear screening is too expensive	106 (32.5)	158 (48.5)	17 (5.2)	45 (13.8)
Thinks it is painful to have pap smear	63 (19.3)	185 (56.7)	10 (3.1)	68 (20.9)
Thinks having a pap smear is unpleasant and embarrassing.	112 (34.4)	139 (42.9)	25 (7.7)	50 (15.3)
Thinks it is difficult to make out time to go for screening	199 (61.0)	36 (11.0)	30 (9.2)	61 (18.7)
Thinks it is difficult to get to the pap smear clinic	201 (61.7)	51 (15.6)	21 (6.4)	53 (16.3)
Thinks Pap smear is unnecessary if there are no signs and symptoms.	137 (42.0)	65 (19.9)	33 (10.1)	91 (27.9)
Thinks being busy is a barrier to Pap smear	151 (46.3)	55 (16.9)	46 (14.1)	74 (22.7)
Thinks it is unnecessary to go for only Pap smear screen	125 (38.3)	108 (33.1)	26 (8.0)	67 (20.6)
Afraid that something wrong will be detected if gone for pap smear screen.	35 (10.7)	17 (5.2)	32 (9.8)	242 (74.2)




Fig. 2. General attitude of respondents toward cervical cancer screening

Table 6. Response of Ambrose Alli University Students Regarding Practice of Cervical Cancer Screening

Questions	Frequency (n=326)	Percent (%)
Have had at least one screening for cervical cancer		
Yes	5	1.5
No	321	98.5
n=5		
Number of times screened for cervical cancer		
Once	5	100.0
Twice	0	0
Multiple	0	0
Influence on decision to go for pap smear		
Friends and family	3	60
Doctors' advice	2	40
Pap smear finding		
Normal	5	100
Would encourage people to go for screening.		
Yes	65	19.9
No	261	80.1

The knowledge of the respondents on cervical cancer and cervical cancer screening as assessed in Table 3, is shown in Fig. 1. 24.8% of respondents had good knowledge, 28.2% had fair knowledge, while 153(46.9%) of respondents had poor knowledge.

There was statistically significant ($P < 0.05$) relationship between the age of respondents and their knowledge of Pap smear. Similarly, a statistically significant association was found between the participants' ethnic group, level and knowledge of Pap smear as well as their

department and knowledge of Pap smear. However, there was no statistically significant association between respondents' religion and knowledge of Pap smear.

From Table 6 above, only 5(1.5) participant have done pap smear, of which 100% of them only had pap smear done once. Friends and family accounted for the highest influence on their decision to go for pap smear. 19.9% of respondents were willing to encourage others to go for screening.




Fig. 3. General practice of respondents toward cervical cancer screening

Table 7. General knowledge and general attitude of respondents

General knowledge of pap smear	General attitude of respondents		Test statistics	P-Value
	Positive (%)	Negative (%)		
Good	44 (53.7)	38 (43.6)	96.432	<0.0001*
Fair	15 (16.3)	77 (83.7)		
poor	2 (1.3)	150(98.7)		

*=Significant ($p < 0.05$)

There is statistically significant relationship between general knowledge of pap smear and general attitude of respondents toward cervical cancer screening.

4. DISCUSSION

This study was conducted to assess the knowledge, attitude and practice of cervical cancer screening among female students at Ambrose Alli University.

This research showed that majority of the participants has heard of cervical cancer. This is higher than the finding in Ethiopia (54.4%) [22] and Cameroon (28%) [23], however it was similar when compared to another study done among female medical students of Crimea State Medical University Ukraine where 80% of the

participants had heard about cervical cancer [24] However, it was observed that despite the fact that most of the respondent have heard of cervical cancer, half of them had poor knowledge on cervical cancer screening. This result was also similar to the study done in Lagos [25], however it was in contrast to another study done in Saudi Arabia [26] as majority of the participants had good knowledge on cervical cancer screening.

Knowledge of cervical cancer screening increased with increasing age as most of the participant between 29 to 31 years had good knowledge of cervical cancer screening while most of the participants aged 17 to 19 had poor knowledge. Similarly, Monica Idestrom's study of Denmark showed that knowledge and concerns regarding Pap smear test were dependent on

age, where younger women showed lesser degree of awareness and practice as compared to the older participants [27]. There was also a significant relationship between knowledge of cervical cancer and age of participant. Since some of the respondents were medical students, this result was expected.

Knowledge of cervical cancer screening also increased with increasing university level as students in 400 and 500 level had higher knowledge when compared to the other levels. The better knowledge of these students can be explained by the fact that these students, especially medical students and those in final year, must have been taught about cervical cancer as a part of their curriculum recently and therefore were more informed about the disease. This was in keeping with similar research done in Al-Ahsa Saudi Arabia where most of the student's had good knowledge of cervical cancer screening [28]. There was also statistical relationship between the knowledge of cervical cancer screening and university level of participants.

In this study, only a few of the study participants had a positive attitude towards cervical cancer screening. This is much lower than a similar study conducted in Hawassa University among female students [29]. 36.8% and 24.5% of the participants strongly agree and agree respectively that administration of Pap smear by a male doctor will be uncomfortable and embarrassing to them and it is in keeping with a research done South Africa [30], while 79.1% and 6.4% strongly agree and agree that being married will prevent them from going for screening. It is also important to state that 106 and 158 respondents strongly agree and agree that Pap smear is expensive. This finding was similar to a research done in Haiti [31] and this may be a huge deterrent in the attitude as well as practice of cervical cancer screening amongst participants. Majority of the respondent are confident that nothing wrong will be detected if they go for Pap smear while 42.0% and 19.9% of respondent strongly agrees and agrees that going for Pap smear is unnecessary if there are no signs and symptoms. This therefore confirms that the knowledge of Pap smear amongst the respondents is low as its purpose is for the prevention of cervical cancer, so individuals need not wait before the onset of signs and symptoms before going for screening.

One third of the respondents think it's unnecessary to go to the hospital just for the

purpose of Pap smear. This therefore implies that the hospital is a good means of encouraging women to go for Pap smear as most visits to the hospital by women are for other reasons and not Pap smear.

There was statistical significance between the level of knowledge of cervical cancer screening and the attitude of the respondents. Over half of the participants with good knowledge of cervical cancer screening had a positive attitude toward cervical cancer screening. However, despite having good knowledge and positive attitude toward screening, the practice of cervical cancer screening among them was low. This could be due to the fact that majority of the respondents were confident that nothing wrong will be detected if they go for Pap smear.

Although practice of screening is reported to be low in Nigeria, studies from rural areas and slums have reported lower practice of cervical cancer screening. Only five (1.5%) respondents in this study had done a cervical cancer screening test at some time. Among women interviewed in two slums, in Lagos, also none had been screened or was aware of a screening test for cervical cancer [32]. In a study at rural Okada a community in Edo state, Southern Nigeria, none had been screened for cervical cancer [33]. A low practice of cervical cancer screening was however, also observed in a study in Olusosun, a commercial and residential area of Lagos where only 5% of the female respondents had undertaken a pap smear [34]. Similarly, in a study in Onitsha, a metropolitan city in Anambra, Southeast Nigeria, only 1.8% of respondents had done a cervical screening test [35].

A survey in Britain reported 91% of women have had a cervical cancer screening test at least once [36] Practice of cervical cancer screening varies globally, being higher in developed countries compared with less developed. Studies carried out in some other developing countries also showed low practice of cervical cancer screening. In a Kenyan study, practice was 6% [37] and 0.8% in a community based study in Elmina, Ghana [38].

Out of the 5 respondents who have been screened for cervical cancers, 3 of them went for screening because of the advice from family and friends while the remaining went because a doctor recommended or advised them to for screening. This further suggests that doctors and

health care workers should advise patients who are eligible for the commencement of screening to go for screening.

The low practice could be due to poor availability of screening services within Esan West Local government and poor knowledge of people about the disease and its screening. Moreover, cervical cancer screening services are mostly available in tertiary health institutions with fairly high cost in most cases.

5. CONCLUSION

In conclusion, knowledge, and attitude regarding cervical cancer screening among female student- participants of Ambrose Alli University recorded a poor outcome. Cervical cancer screening is also low as only 5 (1.5%) of the respondent had ever gone for screening. It was noted that majority of the respondent are confident that nothing wrong will be detected if they go for Pap smear while 42.0% and 19.9% of respondent strongly agrees and agrees that going for Pap smear is unnecessary if there are no signs and symptoms. This therefore confirms that the knowledge of Pap smear amongst the respondents is low as its purpose is for the prevention of cervical cancer, so individuals need not wait before the onset of signs and symptoms before going for screening.

6. RECOMMENDATIONS

6.1 To the Government

1. Public service announcements promoting cervical cancer screening with low cost and effective method also avail free service for screening and its benefits should be carried out by the government, non-government organizations and schools.
2. Policies that encourage doctors to recommend the procedure to their female patients who meet the necessary requirements for commencing cervical cancer screening should be instituted.
3. Policies that encourage the orientation of the new university intakes about cervical cancer and cervical cancer screening should also be instituted.
4. Specific recommendation for screening for cervical cancer screening for all HIV infected women of reproductive age enrolled in care with VIA and officer appropriate channel for treatment and care

for all women screened positive for cervical cancer.

6.2 To the Public Health Physician/and Health Workers

1. Encourage or as appropriate mandate that all women coming for antenatal services are provided with the options of a Pap smear test.
2. Embark on awareness and health sensitization creation on educating the populace on the condition of cervical cancer, with the emphasis on the fact that it is preventable.

6.3 To the University and the General Population

The general population should take the issue of cervical cancer screening seriously as preventing the disease is much better than its management. The more enlightened once who have undergone screening, should advice and encourage others to go the screening as well. Also, lots of research activities should be encouraged and conducted by the university community to ensure further addition to knowledge.

7. LIMITATIONS OF THE STUDY

5. The study cannot be generalized because it was done in restricted the setting it was conducted.
6. There was a financial constraint on the part of researchers as it impeded on the coverage of more participants.
7. The study was a cross-sectional study which implied that it was done within a short time frame and presented a snap shot of the possible outcome. However, a longitudinal study would have covered a longtime frame and presented a holistic view on the subject matter.
8. Researchers encountered participants who did not fill most of questionnaires and some answers were perceived to be false hence, data had to scrutinized and cleanse before analysis.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image

generators have been used during writing or editing of manuscripts. This guarantees the fact that the originality of the study was maintained.

CONSENT AND ETHICAL APPROVAL

Individual's informed consent was obtained from the respondents after detailed explanation of the objectives, scope, and benefits of the study had been made known to them.

The permission of the various head of departments in the selected department was granted before the research was commenced. Ethical clearance was sorted from the Hospital Research Ethics Committee affiliated with the university.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Ellenson L, Pirog E. The female genital tract: Robins and Cotran pathologic basis of disease. Elsevier. 2015;9:991-1009.
2. WHO. Comprehensive Cervical Cancer Control; A Guide to Essential Practice. 2016;16.
3. WHO, United Nations, The World Bank, IARC Globocan. Cervical cancer crisis card; 2013. Available: <http://www.who.int/hpvcentre/statistics/en/>. Accessed on: 8 Feb 2021.
4. Ale M, Vassilakos p. Cervical Cancer In Developing Countries. Available: [gfmer.ch/ccdc/cervicalcancer.htm](http://www.gfmer.ch/ccdc/cervicalcancer.htm). Accessed 9 Feb, 2021.
5. Adegoke O, Kulasmgan S, Vimig B. Cervical cancer trends in the United States: A 35-year population-based analysis. *Journal of Women's Health*. 2012;21:1031-1037.
6. Tome L, Bray F, Siegel R, Ferley J, Lortet-Tieulent J, Jemal A. Global cancer statistics. 2012: *Cancer Journal for Clinicians*. 2015;65:87-108.
7. McClung N, Gargano J, Bennett N. Trends in human papilloma virus vaccine type 16 and 18 in cervical precancers 2008-2014: *Cancer Epidemiol Biomarkers*. 2019;28:602-609.
8. Jeronmon J, Bansil P, Lim J. A multi country evaluation of HPV testing, visual inspection with acetic acid, and papanicolaou testing for the detection of cervical cancer: *International Journal of Gynecological Cancer*. 2014;24:2014.
9. Olowoolere S, Fagbemi A, and Ogunlaja O. Determinants of cervical cancer screening uptake among women In Ilorin, North Central Nigeria: A Community-Based Study: *Journal of Cancer Epidemiology*; 2016. Available: doi.org/10.1155/2016/6469240 Accessed on: 9 Feb, 20201.
10. Butho N, Amita B, Jeremy K, Patricia B, Paulina J. Factors associated with the uptake of cervical cancer screening among women in Portland Jamaica, North American *Journal of Medical Sciences*. 2015;7:104–113.
11. Idowu A, Olowookere S, Fagbemi A, Ogunlaja O. Determinants of Cervical Cancer Screening Uptake among Women in Ilorin, North Central Nigeria: A community-based study, *Journal of Cancer Epidemiology*. Available: <https://doi.org/10.1155/2016/6469240>. Access On: 4 Feb, 2021.
12. WHO. Comprehensive cervical cancer control; A guide to essential practice. 2022;2:27-39.
13. WHO. Costing and national strategic plan on prevention and control of cervical cancer: Nigeria. 2017-2021;2020:3
14. Subramaniam A, Fauci J, Schneider K, Whitworth J, Erickson B, Kim K, Huh W. Invasive cervical cancer and screening: what are the rates of unscreened and underscreened women in the Modern Era. 2011;15(2):110-113.
15. Oguntaye A, Samaila M, Prevention of cervical intraepithelial neoplasia Zaria. *Ann. Afr. Med*. 2010;(9):194 –195
16. Souza J, Hunt B, Asirwa F, Adebamowo C, Lopes G. Cancer care outcome disparity in high, middle and low income countries. *Journal Clinical Oncology*. 2016;34(1):6- 13.
17. Yadav K, Cree I, Field A, Vielh P, Mehrotra R. Importance of cytopathologic diagnosis in early cancer diagnosis in resource- constrained countries. *JCO Glob Oncol*. 2022 F;8:e2100337. DOI: 10.1200/GO.21.00337. Erratum in: *JCO Glob Oncol*. 2022;8:e2200086 DOI: 10.1200/GO.22.00086 PMID: 35213215; PMCID: PMC8887942.
18. FIO, Angela O, Easter A. Perceived Effects of Computer-Based Examination (CBE) on Essay Writing Skills of

- Undergraduates in Ambrose Alli University, Ekpoma, Edo State. *Randwick International of Education and Linguistics Science Journal*. 2021;2(1):59-68. Available:<https://doi.org/10.47175/rielsj.v2i1.200>
19. Cochran formula on statistics and probability. Available:www.enwikipedia.org. Accessed On:Feb 21, 2021.
 20. Ayinde OA, Omigbodun AO, Ilesanmi AO. Awareness of cervical cancer, papanicolaou's smear and its utilisation among female undergraduates in Ibadan. *African Journal of Reproductive Health / La Revue Africaine De La Santé Reproductive*. 2017;8:68-80
 21. James VJ. knowledge, attitude and practice concerning cervical cancer screening among market women in yola-metropolis, adamawa state. *Afr J Reproductive Health*: 2012;7(2):68-80
 22. Tilahun T, Tulu T, Dechasa W. Knowledge, attitude and practice of cervical cancer screening and associated factors amongst female students at Wollega University, western Ethiopia. 2019;12:518
 23. Ahmedin J. Global cancer statistics. *CA Cancer J Clin*. 2011;61:69–90.
 24. Okyere, J., Duodu, P.A., Aduse-Poku, L. *et al*. Cervical cancer screening prevalence and its correlates in Cameroon: secondary data analysis of the 2018 demographic and health surveys. *BMC Public Health*. 2021;21:1071.
 25. Available:<https://doi.org/10.1186/s12889-021-11024-z>
 26. Olubodun T, Odukoya OO, Balogun MR. Knowledge, attitude and practice of cervical cancer prevention, among women residing in an urban slum in lagos, South West, Nigeria. *Pan Afr Med J*. 2019;30:130
 27. Humariya Heena, Sajid Durrani, Isamme AlFayyad, Muhammad Riaz, Rabeena Tabasim, Gazi Parvez, Amani Abu-Shaheen, "Knowledge, Attitudes, and Practices towards Cervical Cancer and Screening amongst Female Healthcare Professionals: A Cross-Sectional Study", *Journal of Oncology*. 2019;2019;9 Article ID 5423130. Available:<https://doi.org/10.1155/2019/5423130>
 28. Tapera R, Manyala E, Erick P, Maswabi TM, Tumoyagae T, Letsholo B, Mbongwe B. Knowledge and attitudes towards cervical cancer screening amongst University of Botswana Female Students. *Asian Pac J Cancer Prev*. 2017;18(9):2445-2450. DOI: 10.22034/APJCP.2017.18.9.2445. PMID: 28952274; PMCID: PMC5720649.
 29. Al-Darwish A, Al-Naim AF, Al-Mulhim KS. Knowledge about cervical cancer early warning signs and symptoms, risk factors and vaccination among students at a medical school in Al-Ahsa, Kingdom of Saudi Arabia," *Asian Pacific Journal of Cancer Prevention*. 2014;15(6):2529–2532.
 30. Tsegaye S. Knowledge, attitude, practice of cervical cancer screening and its associated factors among female students in Hawassa University College of medicine and health science Hawassa Ethiopia. Available:<http://www.hu.edu.et/hu/images/pdf/proceedings/>. Accessed on:3 Mar 2017.
 31. Byrd TL, Chavez R, Wilson KM. Barriers and facilitators of cervical cancer screening among Hispanic women. *Ethn Dis*. 2007;17:129-34.
 32. Brown DR, Wilson RM, Boothe MA, Harris CE. Cervical cancer screening among ethnically diverse black women: Knowledge, attitudes, beliefs, and practices. *J Natl Med Assoc*. 2011; 103:719-28.
 33. Balogun MR, Odukoya OO, Oyediran MA, Ujomi PI. Cervical cancer awareness and preventive practices: A challenge for female urban slum dwellers in Lagos, Nigeria. *Afr J Reprod Health*. 2012;16(1):75-82.
 34. Igwilo AI, Igwilo UU, Hassan F, Idanwekhai M, Igbinomwanhia O, Popoola AO. The knowledge, attitude and practice of the prevention of cancer of the cervix in Okada Community. *Asian Journal of Medical Sciences*. 2012;4(3):95-98.
 35. Wright KO, Aiyedehin O, Akinyinka MR, Ilozumba O. Cervical cancer: community perception and preventive practices in an urban neighborhood of Lagos (Nigeria). *ISRN Preventive Medicine*. 2014;2014:950534.
 36. Nwozor CM, Oragudosi AL. Awareness and uptake of cervical cancer screening among women in Onitsha, South-East, Nigeria. *Greener Journal of Medical Sciences*. 2013;3(8):283-288.

37. Moser K, Patnick J, Beral V. Inequalities in reported use of breast and cervical screening in Great Britain: analysis of cross sectional survey data. *BMJ*. 2009;338:b2025.
38. Sudenga SL, Rositch AF, Otieno WA, Smith JS. Brief report: knowledge, attitudes, practices and perceived risk of cervical cancer among Kenyan women. *Int J Gynecol Cancer*. 2013 Jun;23(5):895-899.
39. Ebu NI, Mupepi SC, Siakwa MP, Sampsele CM. Knowledge, practice and barriers toward cervical cancer screening in Elmina, Southern Ghana. *Int J Wom Health*. 2015;7:31–39.

APPENDIX

KNOWLEDGE, ATTITUDE AND PRACTICE OF CERVICAL CANCER SCREENING AMONG AMBROSE ALLI UNIVERSITY STUDENTS

SECTION ONE: Socio-demographic characteristics of respondents.

Tick as appropriate

Date: _____

1. Age at last birthday: _____
2. Marital status:

Single	[]
Married	[]
Divorced	[]
Separated	[]
Widowed	[]
3. Ethnic group: _____
4. Religion:

Christian	[]
Muslim	[]
ATR	[]
5. Level :

100	[]
200	[]
300	[]
400	[]
500	[]
600	[]
6. Departement: _____

Section Two: Knowledge of cervical cancer screening.

1. Have you ever heard about Pap smear?
 - a. Yes []
 - b. No []
2. If yes, what is your source of information?
 - a. School []
 - b. Health worker []
 - c. Mass media []
 - d. Social media []
 - e. Others (Please specify): _____

Questions	Yes	No	Don't know
9. Have you ever heard of cancer of the cervix?			
10. Women should have Pap Smears at least every 3 years			
11. Pap smear is the most helpful way to detect pre-cancer and cancer of the cervix			
12. The purpose of the Pap smear is to detect abnormal cells in the cervix			
13. Pap smear is able to detect all types of female genital cancer			

Questions	Yes	No	Don't know
14. Pap smear is a non-invasive and relatively inexpensive method			
15. Women should have Pap smear since the onset of sexual activity			
16. Pap smears can be performed at both menstrual and non-menstrual period			
17. A woman should not have sex 24 hours before having Pap smear			
18. Pap smear should be discontinued after menopause			
19. Having a normal Pap smear, means that you don't need Pap smear in the future			
20. There is no need to have a Pap smear if it is not administered by a doctor			
21. Have you heard about HPV vaccine?			

Section Three: Attitude towards cervical cancer screening

s/n		Strongly agree	Agree	Disagree	Strongly disagree
22	If male doctor performed the test, would you feel embarrassed?				
23	You will go for Pap smear even when single				
24	Being married will discourage you from going for Pap smear				
25	Going for Pap smear screening is too expensive				
26	Is it painful to have a Pap smear?				
27	Having a Pap smear is unpleasant/embarrassing				
28	It is difficult to take time off to go for Pap smear				
29	It is difficult to get to the Pap smear clinic				
30	Pap smear is unnecessary if there are no signs and symptoms				
31	Being busy is a barrier to Pap smear				
32	Is it unnecessary to go only for Pap smear test				
33	I am afraid that something wrong will be detected if I go for Pap smear				

Section Four: Practice of cervical cancer screening

s/n		Yes	No
34.	Have you ever done a pap smear?		
35.	If yes, how many times have you been screened for cervical cancer? Once Twice Multiple times		

36. What influenced your decision to go for Pap smear? _____

37. If you had a Pap smear done before, what was the finding?

Normal [], abnormal [], I don't know [].

38. If the Pap smear was abnormal, what next did you do? _____

39. Have you ever encouraged anybody to go for cervical cancer screening? Yes [], No []

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125077>