



Occlusion of the Central Retinal Artery Complicating a Thrombosed Abdominal Aortic Aneurysm: An Exceptional Complication

Belfaiza Soukaina^{a++*}, Mouzari Yassine^{a#}, Reda Karim^{a#} and Oubaz Abdelbar^{a#}

^a *Military Hospital Mohammed V., University Mohammed V., Rabat, Morocco.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95676>

Case Report

Received 23 October 2022
Accepted 29 December 2022
Published 29 December 2022

ABSTRACT

We report the case a patient who presented a sudden decrease in visual acuity of the right eye secondary to an occlusion of the central retinal artery. In addition, the patient complained of intense abdominal pain for which a CT scan was performed and it revealed a thrombosed aneurysm of the abdominal aorta so the diagnosis retained is occlusion of the central retinal artery complicating a thrombosed aneurysm of the abdominal aorta .Occlusion of the central retinal artery (CRAO) is a very rare vascular pathology. It is often secondary to cardiovascular pathology but it can also be a complication of a distant vascular anomaly, hence the interest of an exhaustive etiological diagnosis and the elimination of vascular emergencies such as thrombosed aneurysms of an artery whatever its location such as abdominal or thoracic aorta , especially in the presence of warning signs. The scientific interest of our case is that the occlusion of the central artery of the retina is rare and exceptionally can complicate a thrombosed aneurysm of the abdominal aorta which is a vital emergency.

Keywords: Occlusion of the central retinal artery; thrombosed aneurysm; aortic aneurysm.

⁺⁺ *Intern in Ophthalmology;*

[#] *Professor in Ophthalmology;*

^{*}*Corresponding author: E-mail: soukainabelfaiza@gmail.com;*

1. INTRODUCTION

Occlusion of the central retinal artery (CRAO) is a very rare vascular pathology [1]. It is often secondary to cardiovascular pathology. The most frequent etiology is atheroma of the carotid artery [2], but it can also be a complication of a distant vascular anomaly such as thrombosed abdominal aortic aneurysms, sending emboli that can lead to the occlusion of the ocular vessels [3], hence the interest of a broad diagnostic work-up. We report the case of a patient with an occlusion of the central retinal artery complicating a thrombosed abdominal aortic aneurysm.

2. CASE REPORT

We report the case of a 50-year-old patient, hypertensive, who presented to the emergency department with sudden decreased vision in the left eye. The best-corrected visual acuity was limited to luminous perception in the left eye and 10/10 in the right eye. Slit lamp examination found a relative afferent pupillary deficit in the left eye. The fundus of the left eye found a diffuse retinal edema with a cherry-red aspect of the macula evoking an occlusion of the central retinal artery (Fig. 1). A cardiovascular check-up was normal (heart echo and echodoppler of the neck vessels). An abdominal CT scan was requested by the gastrologist to explore the abdominal pain, which revealed a thrombosed abdominal aortic aneurysm (Fig. 2). The diagnosis of occlusion of the central artery of the retina complicating a thrombosed abdominal aortic aneurysm was retained. The thrombosed abdominal aortic aneurysm was responsible for emboli leading to the occlusion of the central artery of the retina. The patient was given on antiplatelet medication and was operated on by vascular surgery for the thrombosed abdominal aortic aneurysms with the placement of a stent graft.

3. DISCUSSION

Central retinal artery occlusion (CRAO) is a rare vascular pathology, with an estimated frequency of 1 in 10,000 (4). CRAO most often affects subjects in their sixties. The accident is bilateral in only 1 to 2% of cases. The patients describe a sudden decrease in visual acuity. Examination of the eye reveals a white, painless eye. The visual acuity is collapsed, most often limited to a luminous perception. The pupil is in mydriasis areflective [4]. At the fundus, in the first moments, the retina can still have a normal

aspect, but there is a diffuse narrowing of the arterial caliber. It is only in the following hours that a whitish retinal ischemic edema appears, translating the ischemic suffering of the internal layers of the retina. The foveola, which is only vascularized by the choroid, keeps its normal coloring and appears redder in contrast with the rest of the ischemic retina ("cherry-red spot of the macula") [4].

The fluorescein angiography is not essential for diagnosis and should not delay the initiation of treatment. It shows an extreme delay in perfusion of the branches of the central retinal artery and a prolongation of the retinal arteriovenous filling time(4). The causes can be classified in several major classes according to the mechanism involved: embolisms; thrombosis or coagulation disorders [5]. The most frequent emboligenic pathologies are: carotid atheroma, carotid maneuvers or a postoperative context (carotid endarterectomy); emboligenic cardiopathies.much more rarely, lipidic embolisms consecutive to a bone fracture or tumor embolisms (myxoma of the atrium).The second mechanism is thrombosis ,it can be secondary to Horton's disease and systemic diseases, such as systemic lupus erythematosus, Wegener's disease, Takayashu's disease, Kawasaki's disease and Churg-Strauss syndrome. Lastly, they can be linked to coagulation disorders (resistance to activated protein C, deficiency of protein C, protein S or antithrombin); antiphospholipid syndrome; hyperhomocysteinemia) [5]. The etiologies mentioned above are the most frequent, hence the interest of an exhaustive etiological diagnosis and the elimination of vascular emergencies such as thrombosed aneurysms of an artery whatever its location such as abdominal or thoracic aorta, especially in the presence of warning signs [6,7]. The thrombosed aortic aneurysm is responsible for emboli leading to the occlusion of vessels as Central retinal artery [8]. The thrombosed abdominal aortic aneurysm was responsible for emboli leading to the occlusion of the central artery of the retina .The spontaneous evolution is almost always unfavorable in the absence of a rapid repermeabilization, the papilla becomes pale, atrophic and the retinal arteries become sparse, filiform, and the internal layers of the retina atrophy [9]. The treatment of OACR remains disappointing. The objective of all the proposed treatments is to obtain a arterial repermeabilization as soon as possible before the appearance of definitive ischemic retinal lesions: hypotonizing treatment; vasodilator



Fig. 1. The fundus of the left eye found a diffuse retinal edema with a cherry-red aspect at the macula evoking an occlusion of the central retinal artery



Fig. 2. An abdominal CT scan revealed a thrombosed abdominal aortic aneurysm (Arrow)

treatment and anticoagulant treatment [9]. All these treatments are not effective and only the fibrinolytic treatment by intravenous or intra-arterial catheterization of the ophthalmic artery which is really effective [10], but it must be initiated very early and the etiological treatment is necessary for preventing the Central retinal artery occlusion (CRAO) of the other eye ant to save the life of the patient because the etiology can be an emergence pathology vascular as in our case.

4. CONCLUSION

Central retinal artery occlusion (CRAO) is one of the rare true emergencies in ophthalmology. Indeed, not only is visual function threatened in

the very short term, but it can also be a sign of an underlying systemic pathology. This is why the etiological assessment must be exhaustive and eliminate a vascular emergency such as an aneurysm or stenosis of the abdominal aorta, especially in front of warning signs as in our case.

CONSENT

The patient has given their informed consent for the case report to be published.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Brown GC. Real arterial obstructive disease. *Retina*. 1994;2:1361-77.
2. Miyazawa A, Inoue M, Yamada S, Hirota K, Sano RY, Nishiyama K, Hirakata A. Higher incidence of carotid stenosis in patients with central retinal artery occlusion. *Acta Ophthalmologica*. 2011; 89(4):e378-9.
3. Speelman L, Schurink GW, Bosboom EM, Buth J, Breeuwer M, van de Vosse FN, Jacobs MH. The mechanical role of thrombus on the growth rate of an abdominal aortic aneurysm. *Journal of Vascular Surgery*. 2010 Jan 1;51(1):19-26.
4. Duker JS, Sivalingam A, Brown GC, Reber R. A prospective study of acute central retinal artery obstruction: the incidence of secondary ocular neovascularization. *Archives of ophthalmology*. 1991 Mar 1;109(3):339-42.
5. Greven CM, Slusher MM, Weaver RG. Retinal arterial occlusions in young adults. *American Journal of Ophthalmology*. 1995 Dec 1;120(6):776-83.
6. Moulakakis, Maras, Bountouris, Pomoni, Georgakis, Andrikopoulos. "Silent" thrombosis of an abdominal aortic aneurysm not producing acute limb ischemia. *Vasa*. 2010 Aug 1;39(3):265-7.
7. Martelli E, Ippoliti A, Ventoruzzo G, De Vivo G, Marchetti AA, Pistolesse GR. Popliteal artery aneurysm. *Int Angiol*. 2004;23:54-65.
8. Watson TJ, Childers WK, Haga L, Calaitges J. Acute Bilateral Lower Extremity Paralysis Secondary to Acute Thrombosis of an Infrarenal Abdominal Aortic Aneurysm. *AORTA*. 2017 Jun;5(03): 91-5.
9. Lim JY, Lee JY, Chung HW, Yoon YH, Kim JG. Treatment of branch retinal artery occlusion with transluminal Nd: YAG laser embolysis. *Korean Journal of Ophthalmology*. 2009 Dec 1;23(4):315-7.
10. Schumacher M, Schmidt D, Jurklics B, Gall C, Wanke I, Schmoor C, Maier-Lenz H, Solymosi L, Brueckmann H, Neubauer AS, Wolf A. Central retinal artery occlusion: local intra-arterial fibrinolysis versus conservative treatment, a multicenter randomized trial. *Ophthalmology*. 2010 Jul 1;117(7):1367-75.

© 2022 Soukaina et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/95676>