

Recovery Degree in Pediatric Patients with Antecedent Severe Traumatic Brain Injury in Mexico

Sujey Araceli Zurita Rivera ^a, Jose Alfredo Vidal Vidal ^{b*},
Froylan Eduardo Hernandez Lara Gonzalez ^c, Patricia Zagoya Martinez ^d,
Jose Felix Urbina Hernandez ^e, Francisco Tinoco Lemus ^f,
Rosario Ponce Dávila ^g, Atalia Saraí Ruíz Austria ^h,
Emma Juanita Carbajal Reyes ⁱ, Ivonne Juarez Reyes ^j,
Carolina Contreras Camacho ^k, Manuel Alejandro Resendiz López ^k
and Angel Ramon Saucedo Medina ^l

^a Department of Pediatric Patient Medicine in Critical Condition, Mexican Institute of Social Security, Hospital General de Zona No. 1, Colima, Mexico.

^b Department of Critical Medicine, Hospital Para el Niño Poblano, Puebla, Mexico.

^c Teaching and Research Coordinator, Hospital Para el Niño Poblano, Puebla, Mexico.

^d Department of Hematology, Mexican Institute of Social Security, National Medical Center Gral, Manuel Avila Camacho, Puebla, Mexico.

^e Department of Head of the Critical Medicine, Hospital Para el Niño Poblano, Puebla, Mexico.

^f Department of Anesthesiology, Regional University Hospital, Colima, Mexico.

^g Department of Pediatric Patient Medicine in Critical Condition, Mexican Institute of Social Security, National Medical Center, Adolfo Ruiz Cortinez, Veracruz, Mexico.

^h Department of Pediatric Patient Medicine in Critical Condition, Mexican Institute of Social Security, Hospital General de Zona No. 1, Queretaro, Mexico.

ⁱ Department of Critical Medicine, Hospital Para el Niño Poblano, Universidad Nacional Autonoma de Mexico (UNAM), Puebla, Mexico.

^j Infectology Area, Hospital Para el Niño Poblano, Puebla, Mexico.

^k Department of Pediatric Oncology, Mexican Institute of Social Security, Hospital General de Zona No. 1, Colima, Mexico.

^l Department of Pediatric, Mexican Institute of Social Security, Hospital General de Sub Zona No. 4, Colima, Mexico.

Authors' contributions

This work was carried out in collaboration among all authors. Authors SAZR, JAVV and FEHLG collected the data and were involved in writing the manuscript. Authors SAZR, JAVV, FEHLG, PZM, JFUH, FTL, RPD, ASRA and EJCR were involved in writing and editing the manuscript. Authors IJR, CC, MARL, ARSM and FEHLG wrote the discussion and participated in literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJPR/2021/v7i430230

Editor(s):

(1) Dr. Nicolas Padilla-Raygoza, Institute of Public Health, Mexico.

(2) Dr. Emmanouil Magiorkinis, Athens University Medical School, Greece.

Reviewers:

(1) Shiv Bagga, Mount Sinai Hospital, U.S.A.

(2) Ariel Varela Hernandez, University of Talca Regional Hospital, Chile.

(3) Oscar Rojo Del Moral, México.

Complete Peer review History, details of the editor(s), Reviewers and additional Reviewers are available here:

<https://www.sdiarticle5.com/review-history/81094>

[≡] Associate Professor;

*Corresponding author: Email: vidal_vidal18@hotmail.com;

ABSTRACT

Aims: To determine the recovery degree in pediatric patients with antecedent severe traumatic brain injury.

Study Design: Prospective, analytical and relational.

Place and Duration of Study: Pediatrics Intensive Care Area of the "Hospital para el Niño Poblano" from March 2014 to February 2015.

Methodology: Patients aged 2 to 192 months, both sexes, with antecedent severe traumatic brain injury. The degree of recovery was determined with the Glasgow Outcome Score (GOS) at hospital discharge and one year after. Descriptive and inferential statistics were used data analysis.

Results: The study consisted of 23 patients, of which 69.56% were men. The mean age was 73.17±50.33 months. The predominant age group was middle childhood (34.78%). The main mechanism of injury was run over by a vehicle (39.13%). The group of toddler remained hospitalized for 17.50±8.58 days. Moreover, GOS at hospital discharge ($p=0.391$) and at one year ($p=0.789$) was not associated with sex. Additionally, an association of GOS at hospital discharge was found with GOS at one year of care ($p=0.003$), with greater improvement being observed, in those cases with less initial damage brain.

Conclusion: The degree of recovery at one year after hospital discharge in pediatric patients with antecedent severe traumatic brain injury is associated with the degree of recovery at hospital discharge. Middle childhood, being the group at greatest risk. The complications can reduce the recovery of the patient. Rehabilitation therapy provided by the public health services and the family is of vital importance.

Keywords: Critical care; Glasgow Outcome Scale (GOS); head injury; neurotrauma.

1. INTRODUCTION

Traumatic brain injury is defined, as any physical injury or functional impairment of the cranial content secondary to a sudden exchange of mechanical energy [1] or as physical injuries produced on brain tissue that temporarily or permanently alter brain function [2]. The categorization is established from the Glasgow Coma Scale as mild (13-15), moderate (9-12) and severe (3-8) [3]. In recent years, traumatic brain injury has become a major public health problem [4] as it represents, the leading cause of death and disability worldwide among all trauma-related injuries [5].

In the United States, there are 500-800 estimated cases of traumatic brain injury per 100,000 people [6]. In Mexico, mortality related to traumatic brain injury is 38.8 per 100,000 people, especially in population, aged 15-45 years. The most frequent cause, is traffic accidents (75%) [7]. Traumatic brain injury in pediatric patients, is common in Emergency Department, which represents a high risk of

presenting sequels [8]. Despite this, in the general population, it is the leading cause of mortality and disability in children older than 12 months in high-income countries, estimating that about 10% of the children under 18 years of age will suffer some type of head trauma during this stage, with a mortality rate twice as high in children under 12 months [9].

Traumatic brain injury can cause cell death, neurotoxicity mediated by alteration in neurotransmission, cerebral edema, vasospasm and compromised angiogenesis, causing secondary lesions with potential irreversibility and neurological disability [10]. Glasgow Outcome Scale (GOS) is recommended, as an instrument to measure the degree of recovery from traumatic brain injury. It is simple to apply and has reliability, validity and stability in its results [11], categorizing patients in death, neurovegetative status, severe disability, moderate disability and good recovery [12]. The objective of this paper, is to determine the recovery degree in pediatric patients with antecedent severe traumatic brain injury.

2. METHODOLOGY

Prospective, analytical and relational study in Pediatrics Intensive Care Area of the “Hospital para el Niño Poblano” during the period March 2014 to February 2015 with patients, aged 2 to 192 months, both sexes, with severe traumatic brain antecedent injury, regarding Glasgow Coma Scale. Patients with previous neurological pathology, were excluded. Patients were classified in: infancy (28 days-12 months), toddler (13-24 months), early childhood (25-71 months), middle childhood (72-132 months) and early adolescence (133-198 months) [13].

The variables studied were: age, sex, hospital stay and injury mechanisms (run over by vehicle, fall at the same level, fall at different levels, car accident, other). The degree of recovery was obtained by Glasgow Outcome Score (death (1); neurovegetative state (2); severely disabled (3); moderately disabled (4); and good recovery (5); at hospital discharge and one year after [14]. All patients received rehabilitation for one year by the institution (once a moth) and supplemented by their families (Every day of the week). Rehabilitation was oriented to physical, neurological and sensoperception therapy. Improvement with rehabilitation after one year was evaluated using the Glasgow Outcome Score. Descriptive and inferential (Mann-Whitney U test, Kruskal-Wallis test, Chi square, V Cramer, Wilcoxon test) statistics were used data

analysis, with a confidence interval 95%, using the statistical program SPSS Ver. 25.

3. RESULTS

The study was carried out with 23 patients, of which 16 (69.56%) were males. The median age did not present, a statistical difference between the sexes ($p = 0.109$), being higher in males (84.94 ± 55.39 months; median: 82.5 months) than in females (46.29 ± 20.12 months; median: 48 months) (Table 1). The predominant age group, were middle childhood (34.78%) with a mean age of 91.13 ± 22.16 months, followed by early childhood (21.74%) (Table 2).

The principal injury mechanisms, were run over by vehicle (39.13%), fall at the same level (13.39%), fall at different levels (13.04%) and car accident (8.34%). The length of hospital stay respecting gender did not display significant differences ($p = 0.867$), with a median of 12 days (Table 3).

The toddler group stayed longer (17.50 ± 8.58 days; median: 17.50 days), followed by middle childhood (12.13 ± 4.26 days; median: 12.5 days) ($p = 0.464$) (Table 4). Glasgow Outcome Scale (GOS) at hospital discharge, was not associated with sex ($p = 0.391$). Males in this stage showed lower values than females, with 56.25% they found in GOS 2 and GOS 3 (Table 5).

Table 1. Age in months with respect to the sex of the patients

Sex	P value	N	Median	Mean	Standard deviation	Minimum value	Maximum value
Males	0.109	16	82.5	84.94	55.39	2	168
Females		7	48	46.29	20.12	24	80
Total		23	12	73.17	50.33	2	168

Table 2. Classification of patients respect age stage

Classification	N	Median	Mean	Standard deviation	Minimum value	Maximum value
Infancy	2	6	6.00	5.66	2	10
Toddler	4	19	18.75	6.08	13	24
Early childhood	5	55	51.00	9.35	36	59
Middle childhood	8	86.5	91.13	22.16	71	132
Early adolescence	4	150	153.00	11.49	144	168
Total	23	71	73.17	50.33	2	168

Table 3. Hospital stay in days respect to sex

Sex	P value	N	Median	Mean	Standard deviation	Minimum value	Maximum value
Males	0.867	16	12	12.00	4.00	5	20
Females		7	9	13.57	8.60	5	28
Total		23	12	12.48	5.62	5	28

Table 4. Hospital stay in days respect to age stage

Classification	P value	N	Median	Mean	Standard deviation	Minimum value	Maximum value
Infancy	0.464	2	10.5	10.50	2.12	9	12
Toddler		4	17.5	17.50	8.58	7	28
Early childhood		5	8	10.00	6.32	5	21
Middle childhood		8	12.5	12.13	4.26	5	20
Early adolescence		4	12	12.25	3.77	8	17
Total		23	12	12.48	5.62	5	28

Table 5. Glasgow Outcome Scale (GOS) in hospital discharge and one year after, respect to sex

Sex	GOS hospital discharge						GOS one year after hospital discharge					
	1	2	3	4	5	Total	1	2	3	4	5	Total
Males	-	2	7	7	-	16	-	-	1	6	9	16
Females	-	-	2	5	-	7	-	-	1	2	4	7
Total	-	2	9	12	-	23	-	-	2	8	13	23

Table 6. Glasgow Outcome Scale (GOS) in hospital discharge and one year after, respect to age stage

Classification	GOS hospital discharge						GOS one year after hospital discharge					
	1	2	3	4	5	Total	1	2	3	4	5	Total
Infancy	-	-	2	-	-	2	-	-	-	1	1	2
Toddler	-	-	2	2	-	4	-	-	1	1	2	4
Early childhood	-	-	2	3	-	5	-	-	-	2	3	5
Middle childhood	-	1	1	6	-	8	-	-	-	2	6	8
Early adolescence	-	1	2	1	-	4	-	-	1	2	1	4
Total	-	2	9	12	-	23	-	-	2	8	13	23

GOS one year after hospital discharge, was not associated with gender ($p=0.789$). An association of GOS at discharge was found with GOS, one year after hospital discharge ($p = 0.003$). The most frequent hospital discharge was GOS 4 (52.17%), followed by GOS 3 (39.13%) and GOS 1 (8.7%). One year after hospital discharge, those that predominated were, GOS 5 (56.52%) and GOS 4 (34.78%). Furthermore, it observed an significant increase in the score ($p=0.000$) of the patients under study (Table 6).

4. DISCUSSION

The number of males, included in this study was higher than the number of females, coinciding with other investigations carried out in Mexico and other countries [6,7]. Regarding the average age of the patients studied, it was similar to reported by another investigations [15,16]. However, it should be emphasized that the level of medical care of each institution, affected this variable.

The predominant age groups were, middle childhood and early childhood, similar data to those recorded in other investigations. [16,17] This is due to the fact that, the minor begins increasing the risk of suffering accidents without an appropriate care.

The foregoing is related to being run over by vehicle, as the main mechanism of injury found in this research and other authors ratified [7,18,19]. The longest hospital stay was, in the toddler group, which is similar to the data reported, in another investigation carried out in Mexico [7]. Nevertheless, it differs from other investigations, probably due to the availability of resources and the complications that each patient presents [20].

The Glasgow Outcome Scale (GOS) recorded, at hospital discharge the male presented, the lowest values (GOS 2 and GOS 3) and they are consistent with other studies [16,17,21,22] it found an association of sex patients with the mechanisms of injury, the level of severity and the length of hospital stay. Besides, GOS one year after hospital discharge was similar to the reported by other studies [21,22] that indicated a pathophysiological response of the brain to trauma which was higher in children and young people.

5. LIMITATIONS

The results generated in this study, they are still insufficient to establish a strong management criteria, regarding the recovery of pediatric patients with antecedents of severe traumatic brain injury, mainly due to the lack of resources allocated to the development of research, medical follow-up and rehabilitation. The experience resulting, in this study should be considered, as an opportunity to improve the quality of medical care and the educational and training processes, within the institution.

6. CONCLUSION

The degree of recovery at one year after hospital discharge, in pediatric patients with antecedent severe traumatic brain injury is associated with the degree of recovery at hospital discharge. Middle childhood was the group at greatest risk. Rehabilitation therapy provided by the public health services and the family are of great importance. Nonetheless, due to the prevailing social and economic characteristics in Mexico and access to health services it is difficult to achieve this and above all related to

rehabilitation. Therefore, it is important to carry out educational actions about accident prevention, in pediatric patients will reduce this type of injuries and their possible consequences, in the medium and long term. Therefore, it is recommended continue with research related to the subject, in order to propose management alternatives.

CONSENT

All authors declare that written informed consent was obtained, from the parents of patients for publication of this paper.

ETHICAL APPROVAL

The research work was examined and approved by the hospital research and ethics committee.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Keating CE, Cullen DK. Mechanosensation in traumatic brain injury. *Neurobiology of Disease*. 2021;148:105210. DOI: 10.1016/j.nbd.2020.105210
2. Schepici G, Silvestro S, Bramanti P, Mazzon E. Traumatic brain injury and stem cells: an overview of clinical trials, the current treatments and future therapeutic approaches. *Medicina*. 2020; 56(3):137. DOI:10.3390/medicina56030137
3. Jain S, Iverson LM. Glasgow Coma Scale. In: *StatPearls*. StatPearls Publishing, Treasure Island (FL); 2021. PMID: 30020670.
4. Jiang JY, Gao GY, Feng JF, Mao Q, Chen LG, Yang XF, et al. Traumatic brain injury in China. *The Lancet Neurology*. 2019;18(3):286-295. DOI: 10.1016/S1474-4422(18)30469-1
5. Mollayeva T, Mollayeva S, Colantonio A. Traumatic brain injury: sex, gender and intersecting vulnerabilities. *Nature Reviews Neurology*. 2018;14(12):711-722. DOI: 10.1038/s41582-018-0091-y
6. Dewan MC, Rattani A, Gupta S, Baticulon RE, Hung Y, Punchak M, Agrawal A, Adeleye AO, Shrivastava MG, Rubiano AM, Rosenfeld JV, Park KB. Estimating the global incidence of traumatic brain injury.

- Journal of Neurosurgery JNS. 2019;130(4): 1080-1097.
DOI: 10.3171/2017.10.JNS17352
7. Muñana RJE., Ramírez EA. Glasgow Coma Scale: Origin, analysis, and appropriate use. *Enfermería universitaria*. 2014;11(1):24-35.
DOI: 10.1016/S1665-7063(14)72661-2
 8. Pavlovic D, Pekic S, Stojanovic M, Popovic V. Traumatic brain injury: neuropathological, neurocognitive and neurobehavioral sequelae. *Pituitary*. 2019;22:270–282.
DOI: 10.1007/s11102-019-00957-9
 9. Crompton EM, Lubomirova I, Cotlarciuc I, Han TS, Sharma SD, Sharma P. Meta-analysis of therapeutic hypothermia for traumatic brain injury in adult and pediatric patients. *Critical Care Medicine*. 2017; 45(4):575-583.
DOI: 10.1097/CCM.0000000000002205
 10. Ng SY, Lee AYW. Traumatic brain injuries: pathophysiology and potential therapeutic targets. *Frontiers in Cellular Neuroscience*. 2019;13:528.
DOI: 10.3389/fncel.2019.00528
 11. Rossato E, Verzini E, Scandola M, Ferrari F, Bonadiman S. Role of LCF scale as an outcome prognostic index in patients with traumatic brain injury. *Neurological Sciences*. 2021; 42(7):2747-2752.
DOI: 10.1007/s10072-020-04852-1
 12. Giner J, Galán LM, Teruel SY, Espallargas MG, López CP, Guerrero AI, Frade JR. Traumatic brain injury in the new millennium: new population and new management. *Neurología (English Edition)*; 2021.
DOI: 10.1016/j.nrleng.2019.03.024
 13. Williams K, Thomson D, Seto I, Contopoulos-Ioannidis DG, Ioannidis JP, Curtis S, Constantin E, Batmanabane G, Hartling L, Klassen T. Standard 6: age groups for pediatric trials. *Pediatrics*. 2012;129S(3):S153–S160
DOI:10.1542/peds.2012-00551
 14. Kreitzer NP, Hart K, Lindsell CJ, Manley GT, Dikmen SS, Ratcliff JJ, et al. A Comparison of Satisfaction with Life and the Glasgow Outcome Scale Extended after Traumatic Brain Injury: An Analysis of the TRACK–TBI Pilot Study. *The Journal of Head Trauma Rehabilitation*. 2019; 34(3):E10.
DOI: 10.1097/HTR.0000000000000457
 15. Kochanek PM, Tasker RC, Carney N, Totten AM, Adelson PD, Selden NR, et al. Guidelines for the management of pediatric severe traumatic brain injury: update of the brain trauma foundation guidelines, executive summary. *Neurosurgery*. 2019;84(6):1169-1178.
DOI: 10.1093/neuros/nyz051
 16. Asarnow RF, Newman N, Weiss RE, Su E. Association of attention-deficit/hyperactivity disorder diagnoses with pediatric traumatic brain injury: a meta-analysis. *JAMA Pediatrics*. 2021;175(10):1009-1016.
DOI: 10.1001/jamapediatrics.2021.2033
 17. Treble-Barna A, Schultz H, Minich N, Taylor HG, Yeates KO, Stancin T, Wade SL. Long-term classroom functioning and its association with neuropsychological and academic performance following traumatic brain injury during early childhood. *Neuropsychology*. 2017;31(5):486.
DOI: 10.1037/neu0000325
 18. Nakhjavan-Shahraki B, Yousefifard M, Hajighanbari MJ, Oraii A, Safari S, Hosseini M. Pediatric Emergency Care Applied Research Network (PECARN) prediction rules in identifying high risk children with mild traumatic brain injury. *European journal of trauma and emergency surgery*. 2017;43(6): 755-762.
DOI: 10.1007/s00068-017-0811-9
 19. Chen C, Peng J, Sribnick EA, Zhu M, Xiang H. Trend of age-adjusted rates of pediatric traumatic brain injury in US emergency departments from 2006 to 2013. *International Journal Of Environmental Research and Public Health*. 2018;15(6):1171.
DOI: 10.3390/ijerph15061171
 20. Lumba-Brown A, Yeates KO, Sarmiento K, Breiding MJ, Haegerich TM, Gioia GA, et al. Centers for Disease Control and Prevention guideline on the diagnosis and management of mild traumatic brain injury among children. *JAMA pediatrics*.2018;172(11):e182853-e182853.
DOI: 10.1001/jamapediatrics.2018.2853
 21. Krishnamoorthy V, Temkin N, Barber J, Foreman B, Komisarow J, Korley FK. et al. Association of early multiple organ dysfunction with clinical and functional outcomes over the year following traumatic brain injury: a transforming research and clinical knowledge in traumatic brain injury study. *Critical care medicine*. 2021;49(10):1769-1778.
DOI: 10.1097/CCM.0000000000005055

22. Slovis JC, Gupta N, Li NY, Kernie SG, Miles DK. Assessment of recovery following pediatric traumatic brain injury. *Pediatric Critical Care Medicine*. 2018; 19(4):353-360. DOI: 10.1097/PCC.0000000000001490

© 2021 Rivera et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/81094>