



# **Musculo-skeletal Symptoms and Its Severity among Radiographers and Sonographers in Southern Nigeria**

**Benjamin Effiong Udoh<sup>1\*</sup>, Ulu Okoro Ulu<sup>2</sup>, Joseph Chukwuemeka Eze<sup>3</sup>  
and Hyacienth Uche Chiegwu<sup>3</sup>**

<sup>1</sup>Department of Radiography and Radiological Science, Faculty of Allied Medical Sciences, University of Calabar, Calabar, Cross River State, Nigeria.

<sup>2</sup>Department of Radiology, Jos University Teaching Hospital, Jos, Plateau State, Nigeria.

<sup>3</sup>Department of Radiography and Radiological Science, Faculty of Health Sciences and Technology, Nnamdi Azikiwe University, Nnewi, Anambra State, Nigeria.

## **Authors' contributions**

*This work was carried out in collaboration among all authors. Author BEU designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors UOU and JCE managed the analyses of the study. Author HUC managed the literature searches. All authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/JAMMR/2019/v29i1230152

### Editor(s):

(1) Dr. Ashish Anand, Department of Orthopaedic Surgery, GV Montgomery Veteran Affairs Medical Center, Jackson, MS, USA.

(2) Dr. Panagiotis Korovessis MD, PhD, Chief Orthopaedic Surgeon, Orthopaedic Department, General Hospital "Agios Andreas" Patras, Greece.

### Reviewers:

(1) Loveness Anila Nkhata, University of Zambia, Zambia.

(2) Pedro Gómez Piqueras, University of Castilla-La Mancha, Spain.

(3) Ashraf Ramadan Hafez, Deraya University, Egypt.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/34720>

**Original Research Article**

**Received 07 May 2017**

**Accepted 26 July 2017**

**Published 24 June 2019**

## **ABSTRACT**

**Background:** A lot of activities have been found to contribute to musculoskeletal symptoms (MSS) among different professionals such as lifting of heavy objects, prolonged sitting or standing, long schedule of duties equipment manipulation and others. The aims of this study were to assess the prevalence and severity as well as risk factors of musculoskeletal symptoms among practicing radiographers and sonographers in the southern part of Nigeria.

\*Corresponding author: E-mail: [ben123trial@yahoo.com](mailto:ben123trial@yahoo.com);

**Methods:** This study involved distribution of questionnaire to radiographers and sonographers within the southern part of Nigeria. A total of 48 radiographers and 45 sonographers were involved in this study. Information collected from the participants included categories of work performed, schedule/ duration of duty, types of equipment used, work organizational factors, prevalence and severity of musculoskeletal symptoms to mention a few.

**Results:** Out of the 48 Radiographers and 45 sonographers that participated in the study, 76.3% complained of at least one episode of symptom. Low back pain was the most frequent complaint (52.6%), followed by neck pain, shoulder pain and hand/wrist pain. Reports showed that some of the respondents were prevented from performing their duties as a result of the symptoms experienced. The major risk factors were prolonged standing/sitting during prolonged duration of duty, lifting of patients and heavy equipment, overstretching of the neck following heavy work schedules.

**Conclusion:** There is a high prevalence of musculoskeletal symptoms among radiographers and sonographers in Southern Nigeria. Work schedule/duration of duty and lifting of heavy loads are important factors to be controlled in order to reduce the prevalence and severity of musculoskeletal symptoms among radiographers and sonographers in Southern Nigeria.

*Keywords: Muscles; pain; symptoms; waste; neck; sonographers; radiographer.*

## 1. INTRODUCTION

Musculoskeletal symptoms (MSS) also called repetitive motion disorders, are injuries of the muscles, tendons, ligaments, joints, cartilages, nerves and spinal disks [1]. They are degenerative diseases and inflammatory conditions that cause pain and impair normal activities [2]. They can affect many different parts of the body including the upper and lower back, neck, shoulders and extremities [3]. Common symptoms of musculoskeletal pain are localized or wide spread and can be worsened with movement, stiffness of the entire body, fatigue, sleep disturbance, twitching muscle and the sensation of burning in muscles [4,5].

Low back pain is the most common type of musculoskeletal pain and is caused by overuse, poor posture or prolonged immobilization, and wrong movements used during long working hours [1,4]. Some researchers [6] investigated the prevalence of back pain among radiographers and the various work exposures in form of method utilized to perform specific tasks. The results obtained showed that individual strategy is a source of variability for the work performed which influences the presence of back pain and other forms of disorder.

A lot of activities have been found to contribute to MSS among different health care professionals. These include highly repetitive movements, poor posture, forceful hand application, hand-arm vibration, and direct mechanical pressure on body tissues, cold work environments, work organization, and workers' perception of work

organization [7-12]. In addition, psychosocial factors such as work demands, job control, job content, and social support have been implicated for increased report of work-related musculoskeletal disorders (WRMDs) [13]. Faulty ergonomics workstation has been noted to increase the prevalence of musculoskeletal disorders [14]. Keyserling [15] stated that weight of objects lifted, horizontal reaching distance, awkward posture and frequency of exertions play a significant role in the development of low back pain. Other studies have shown that ergonomic stressors such as repetitive motion, intense force, awkward posture, and duration of work activity may be related to the development of upper extremity discomfort, and musculoskeletal disorder such as tendinitis, and carpal tunnel syndrome [16-19]. It is thought that through the various tasks required of some health professionals such as Radiographers and sonographers, they may be exposed to the factors mentioned above, and therefore are at risk of developing work-related musculoskeletal disorders (MSDs) [20].

MSDs are an increasing healthcare issue globally, being the second leading cause of disability [5]. For example in the US, there were more than 16 million strains and sprain treated in 2004, and the total cost for treating MSDs is estimated to be more than \$125 billion per year [21]. In 2006, approximately 14.3% of the Canadian population was living with a disability, with nearly half due to MSDs [22].

Radiographers and sonographers are trained healthcare professionals responsible for

performing diagnostic and therapeutic imaging procedures and providing essential service in clinical healthcare system. These tasks are achieved by manual handling of patients, equipments and materials at different postures and work schedules and as a result, they are prone to occupational musculoskeletal symptoms and disorders because of the heavy patients, equipment and materials handled [23,24].

Although a great variety of tasks are performed by radiographers and sonographers in clinical settings, but hitherto very few works have been carried out to assess the prevalence of musculoskeletal symptoms among radiographers and sonographers in southern part of Nigeria. The aims of this study were to assess the prevalence and severity as well as risk factors of musculoskeletal symptoms among practicing radiographers and sonographers in the southern part of Nigeria.

## 2. METHODS

This was a non-experimental descriptive study that involved collecting information from practicing radiographers and sonographers in different hospitals located within different states in the southern part of Nigeria. These hospitals include; Nnamdi Azikiwe University Teaching Hospital (NAUTH) Nnewi, New Hope Diagnostic Centre Onitsha, both in Anambra State, National Orthopaedic Hospital, Enugu, University of Nigeria Teaching Hospital (UNTH) Ituku-Ozala, Enugu State University of Science and Technology Teaching Hospital, Parklane, Enugu State, Federal Teaching Hospital Abakali, Ebonyi State, Assurance Medical Diagnostic Centre, Calabar, Cross River State and many other hospitals and diagnostic centres within the region of study. Ethical approval for the study was obtained from the Human Research and Ethics Committee of Assurance Medical Diagnostic Center in Calabar.

Data for this study was collected using questionnaires which were previously given to experienced experts in the field of study to validate question wordings, question contents, response alternatives and other forms of validations to the questionnaire. Questionnaires were issued to 53 radiographers and 50 sonographers practicing in the named hospitals. Out of these numbers, 48 and 45 questionnaires were returned and accepted from radiographers and sonographers respectively. The questionnaires contained questions that provide

demographic data and work characteristics, work description, postures maintained for prolonged periods, frequency of equipment use and areas of the body where the participants (radiographers and sonographers) experienced pain/symptom and other necessary information. The information on pain experienced was restricted to four major areas of the body that is, the neck, shoulders, low back and hand/wrist.

Respondents were asked to state the possible cause of the symptom(s) experienced. Exposures to the risk factors of MSSs were measured by the time spent per work day moving patients, standing, sitting or using computers and the postures taken to perform the normal daily task.

Data was analyzed using the statistical package for social sciences (SPSS) version 16.0 for windows. Descriptive statistics were used to obtain the mean, standard deviation and frequency of the variables studied while the chi square was used to assess the significance of the demographic factors and work characteristics on the development of musculoskeletal symptoms. Correlation equations were used to examine the relationships between potential risk factors and MSSs. Statistical level of significance was  $P \leq 0.05$ .

## 3. RESULTS

Out of the 48 radiographers and 45 sonographers that participated in the study, 76.3% complained of at least one episode of symptom. Low back pain was the most frequent complaint (52.6%) among the radiographers, while neck/ shoulder pains was the most frequent complaint among the sonographers (40.4%) followed by hand/wrist pains. Results showed that 66.8% of the respondents were prevented from performing their duty as a result of the symptoms experienced. The major risk factors were prolonged standing during prolonged duration of duty, lifting of patients and heavy equipment, overstretching of the neck/ shoulder following heavy work schedules and cold work environment. Some demographic and anthropometric variables such as gender, age and body mass index had significant association with occurrence of symptoms.

## 4. DISCUSSION

Excessive forces can trigger different pathophysiological processes depending on the

**Table 1. Correlation (r, p) between job content/activity and occurrence of symptoms**

<b>Activity</b>	<b>Low back pain</b>	<b>Shoulder pain</b>	<b>Neck pain</b>	<b>Hand/wrist pain</b>
Excessive standing/walking while on duty.	0.83	0.14	0.02	0.03
Lifting of heavy load/patients	0.00	0.07	0.23	0.12
Excessive stretching of the neck	0.88	0.25	0.01	0.53
Prolonged stretching of the shoulder	0.00	0.04	0.32	0.03
Excessive gripping of object with the hand	0.01	0.17	0.81	0.07
	0.32	0.06	0.00	0.26
	0.03	0.85	0.03	0.26
	0.14	0.00	0.14	0.04
	0.01	0.34	0.02	0.74
	0.32	0.04	0.23	0.01

**Table 2. Severity of symptoms**

<b>Severity</b>	<b>Low back pain</b>	<b>Shoulder pain</b>	<b>Neck pain</b>	<b>hand/wrist pain</b>
Hospitalized due to symptoms	0%	0%	0%	0%
Changed duty temporally due to symptoms	9.0%	5.2%	17.4%	12.1%
Excused duty due to symptoms	60.7%	80.1%	70.7%	7.3%
Managed to work with symptoms	30.3%	14.7%	3.5%	80.6%

tissues affected. It is the magnitude, duration and frequency of loading as well as the time for recovery which determine the physiological effect. Different symptoms in different body sites will depend on how far the physiological processes have advanced.

More than seventy six percent of the respondents complained of at least one episode of symptoms of musculoskeletal pains which occurred as a result of performance of their professional task. Low back (waist) pain was the most frequent complaint among the participants, followed by neck/shoulder pains and hand/wrist pains. To date, low back pain remains the most prevalent and most common work related injury among health workers [25]. Other studies have reported low back pain as most common musculoskeletal pain among radiographers and other health workers [26-29].

The origin of most musculoskeletal disorders (MSDs) lies in a mismatch between the external load and the capacity of the human body to resist biomechanical and physiological strain [30]. The waist, being the center or fulcrum of the body is surrounded and suspended by muscles and ligaments, which receive most of the impacts of these mechanical forces. The muscles produce the forces needed for different activities at work. The mechanical output of the human body is best in the neutral posture of the joints. The local forces acting on the tissues increase at the

extreme postures and extreme limits of the joint movements. The capacity depends on individual factors such as body build and size, gender, age, and general health, and it weakens with age due to MSDs. In addition to the foregoing, gravity acts continuously on parts of the body and muscle force is needed to maintain certain body postures. Studies have shown that Mechanical forces are the most important factors placing stress onto the musculoskeletal system [30].

Several different physiological responses can account for the muscle pain related to muscular work [31]. By nature, muscle contraction means increased pressures within the muscles resulting in obstruction of blood flow in the vessels. Lack of blood flow (ischemia) is a potential cause of pain due to static contractions. With high muscular exertions, internal rupture of muscle cells is possible. In addition, the accumulation of Ca<sup>++</sup> ions can cause cell damage [32].

Our results also show that some demographic and anthropometric variables such as gender, age and body mass index had significant association with occurrence of symptoms (p< 0.05). Females were twice more likely to suffer from MSS than their male counterpart irrespective of their age. This could be explained by the fact that the mean maximal muscle strength of women is about 2/3 of that of men, independent of body size [33,34]. It therefore means that males have a higher capacity of the

body to resist biomechanical and physiological strain than their female counterpart. The results show that 68.2% of the MSS occurred in people of 40 years and above. Age was therefore a factor for the occurrence of MSS. Studies have shown that muscle strength grows during adolescence but starts a gradual decrease before the age of 30 years. First this decrease is small but the decline increases with years and is about 8–16% per decade after approximately 50 years of age [33,34].

Factors which predisposed the respondents to the risks of MSSs from the study (as suggested by the respondents) include lifting of patients and equipments, prolonged standing/walking during prolong duty schedule, wrong postures and excessive work load. This finding is in consonant with that of Kumar et al. [35] who conducted a research on the biomechanical loads placed on x-ray technologists during a selected number of tasks such as pushing a mobile unit, lifting a patient from wheel chair to mention a few. Their studies concluded that the occurrence of musculoskeletal symptoms increases linearly with the tasks of x-ray technologists. On the other hand, Vahdati et al. [36], in their study, stated that no significant relationship was found between incidence of lower back pain and lifting of heavy loads. The environment and task encountered by individual radiographer/sonographer vary depending on the clinical specialty. Tasks in some clinical specialties may be associated with increased risk of developing a certain musculoskeletal symptoms than others. Sonographers, in our study, had increased risk of developing neck and shoulder pain than low back pain while radiographers had increased risk of developing low back pain than neck pain. This is so because for the sonographers, most of the mechanical forces are exerted on the neck and shoulder as the muscles of these region are always stretched during the scanning process. The radiographers usually take some uncomfortable postures and repetitive movements like bending to lift the patient from wheel chair and positioning of the patients on the x-ray couch during radiographic examinations and therefore more prone to risk of low back pain. Bongers et al. [37] found that there was an association between monotonous work, high workload, time pressure, poor work content, stress symptoms at work and the development of neck and shoulder pain. It therefore means that MSSs would be reduced with modification of work characteristics, work content, equipment design and work environment.

## 5. CONCLUSION

Low back pain, neck/shoulder pains were the most common musculoskeletal symptoms experienced by radiographers and sonographers in Southern Nigeria. Although none of the respondent who experienced the symptoms was hospitalized, most of them were prevented from performing their normal duties due to the symptom(s) experienced. Adjusting working schedules (shifts), and modification of equipment design can reduce work-stress and consequently musculoskeletal symptoms.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

As per international standard or university standard, written approval of Ethics committee has been collected and preserved by the authors.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Karahan A, Kav S, Abbasoglu A, Dogan N Low back pain: Prevalence and associated risk factors among hospital staff. *J. Adv Nurs.* 2009;65(3):516-24.
2. Cote JN, Ngomo S, Stock S, Messing K, Vezina N, Antle D, Delisle A, St-Vincent M. Quebec research on work-related musculoskeletal disorders. Deeper understanding for better prevention. *Industrial Relations.* 2013;68(4):643-660.
3. Kuorinka I, Jenssen B, Kilbom A, Vinterberg H, Biering-Sørensen I, Anderson G, Jørgensen K. Standardized nordic questionnaires for the analysis of musculoskeletal symptoms. *Applied Ergonomics.* 1987;18(3):233-237.
4. [www.Clevelandclinic.org](http://www.Clevelandclinic.org). Musculoskeletal pain. (Accessed 13<sup>th</sup> January 2017)
5. Barbe MF, Gallagher S, Massicotte VS, Tyteu M, Popoff SN, Bar-Giuespie AE. The interaction of force and repetition on musculoskeletal and neural tissue responses and sensorimotor behaviour in

- a rat model of work-related musculoskeletal disorders. *BMC Musculoskeletal Disorders*. 2013;14(1):1-51.
6. Wright DL, Witt PL. Initial study of back pain among radiographers. *Radiologic Technology*. 1993;64(5):283-289.
  7. Habibi E, Pourabdian S, Atabaki AK, Hoseini M. Evaluation of work-related psychosocial and ergonomics factors in relation to low back discomfort in emergency unit Nurses. *Int J Prev Med* 2012;3:564-8.
  8. Rambabu T, Suneetha K. Prevalence of work related musculoskeletal disorders among physicians, surgeons and dentists: A comparative study. *Ann Med Health Sci Res*. 2014;4:578-82.
  9. Chyuan JA, Du C, Yeh W, Li C. Musculoskeletal disorders in hotel restaurant workers. *Occup Med (Lond)* 2004;54:55-7.
  10. Attar SM. Frequency and risk factors of musculoskeletal pain in nurses at a tertiary centre in Jeddah, Saudi Arabia: A cross sectional study. *BMC Res Notes*. 2014;7:61-7.
  11. Alghadir A, Zafar H, Iqbal ZA. Work-related musculoskeletal disorders among dental professionals in Saudi Arabia. *J Phys Ther Sci*. 2015;27:1107-12.
  12. Lorusso A, Vimercati L, L'Abbate N. Musculoskeletal complaints among Italian X-ray technology students: A cross-sectional questionnaire survey. *BMC Res Notes*. 2010;3:114.
  13. Oyewole OO, Adeniyi EA, Ajayi BF, Olajitan AA, Oritogun KS. Work-related musculoskeletal disorders and ergonomic stressors among direct and non direct contact health care workers from a Nigerian tertiary health facility. *Journal of Orthopedics and Allied Sciences*. 2016;15(1):7-13.
  14. Thibodeau PL, Melamut SJ. Ergonomics in the electronic library. *Bull Med Libr Assoc*. 1995;83:322-9.
  15. Keyserling WM. Workplace risk factors and occupational musculoskeletal disorders: A review of biomechanical and psychophysical research on risk factors associated with low-back pain. *AIHAJ*. 2000b;61:39-50.
  16. Silverstein BA, Fine LJ, Armstrong TJ. Occupational factors and carpal tunnel syndrome. *American Journal of Industrial Medicine*. 1987;11:343-358.
  17. Smith AC, Wolf JG, Xie GY, Smith MD. Musculoskeletal pain in cardiac ultrasonographers: Results of a random survey. *Journal of the American Society of Echocardiography*. 1997;12(4):363-66.
  18. Sjogaard G, Sogaard K. Muscle injury in repetitive motion disorder. *Clinical Orthopedics and Related Research*. 1998; 351:21-31.
  19. Keyserling WM. Workplace risk factors and occupational musculoskeletal disorders: A review of biomechanical and psychophysical research on risk factors associated with upper extremity disorders, *AIHAJ*. 2000a;61:231-243.
  20. Lamar SI. Investigation of factors associated with prevalence and severity of musculoskeletal symptoms among workers in clinical specialties of Radiologic Technologic: An Ergonomic and Epidemiological Approach; 2014. Available:<http://www.Lib.ncsu.edu/resolver>
  21. Gallagher S, Heberger JR. Examining the interaction of force and repetition on musculoskeletal disorder risk: A systemic literature review. *Human Factors*. 2013; 55(1):108-124.
  22. Goodridge D, Lawson J, Marciniuk D, Rennie D. A population-based profile of adult Canadians living with participation and activity limitations. *Canadian Medical Association Journal*. 2011;183(13):1017-1024.
  23. Smedley J, Inskip H, Trevelyan F, Buckle P, Cooper C, Coggon D. Risks factors for incident neck and shoulder pain in hospital nurses. *Occupational and Environmental Medicine*. 2003;60(11):564-69.
  24. Lagerstnom M, Wenemark M, Hagberg Hjeem EW. Occupational and individual factors related to musculoskeletal symptoms in five body regions among Swedish nursing personnel. *International Archives of Occupational and Environmental Health*. 1995;68(1):27-35.
  25. www. Global year against musculoskeletal pain; 2009-2010. (Assessed 28<sup>th</sup> January 2017)
  26. Ugwu AC, Egwu OA, Ochie K, Ewunonu EO, Ovumba KN, Njoku CO. Incidence of occupational stress among medical radiographers: a population based zonal survey. *Nigerian Journal of Physiological Sciences*. 2007;22(1-2):123-127.
  27. Abdulmujeeb AB. Prevalence and factors associated with low back pain among healthcare workers in Kibuli Muslim

- Hospital Kampala, Uganda. *Epidemiology (Sunnyvale)*. 2017;7:287.  
DOI:10.4172/2161-1165.1000287
28. Wong TS, Teo N, Kyaw MO. Prevalence and risk factors associated with low back pain among health care providers in a district hospital. *Malaysian Orthopaedic Journal*. 2010;4(2):23-28.
  29. Ramin M, Narges SSH, Sara A, Mina YM. Prevalence of low back pain in health care workers and comparison with other occupational categories in Iran: A Systematic Review. *Iran J Med Sci*. 2016;41(6):467-478.
  30. Esa-Pekka T, Klaus K. Pathophysiological mechanisms of musculoskeletal disorders. Available: <https://oshwiki.eu/index.php> (Accessed 25<sup>th</sup> January 2017)
  31. Visser B, van Dieen JH. Pathophysiology of upper extremity muscle disorders. *J Electromyogr Kinesiol*. 2006;16:1-16.
  32. Stoll T, Huber E, Seifert B, Michel BA, Stucki G. Maximal isometric muscle strength: Normative values and gender-specific relation to age. *Clin Rheumatol*. 2000;19:105-13.
  33. Danneskiold-Samsøe B, Bartels EM, Bulow PM, Lund H, Stockmarr A, Holm CC, Wtjen I, Appleyard M, Bliddal H. Isokinetic and isometric muscle strength in a healthy population with special reference to age and gender', *Acta Physiol Oxf*. 2009;197(673):1-68.
  34. Kumar S, Moro L, Yogesh NA. Biomechanical analysis of loads on x-ray technologists: A field study. *Ergonomics* 2003;46(5):502-517.
  35. Vahdati SS, Khiavi RS, Ghafouri RR, Adimi I. Evaluation of prevalence of low back pain among residents of Tabriz University of Medical Sciences in relation with their position in work. *Turk J Emerg Med*. 2014;14(3):125-129.
  36. Bongers PM, Dewineer CR, Kompier MAJ. Psychosocial factors at work and musculoskeletal disease. *Scandinavian Journal of Work Environment and Health*. 1993;19:297-312.
  37. Bongers PM, de Winter CR, Kompier MA, Hildebrandt VH. Psychosocial factors at work and musculoskeletal disease. *Scandinavian Journal of Work, Environment & Health*. 1993;297-312.

© 2019 Udoh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle3.com/review-history/34720>