



# Effect of Sowing Date and Variety on Pigeonpea Production in Nepal

Anil Pokhrel <sup>a\*</sup>, Sarita Manandhar <sup>b</sup>  
and Sangharsh Raj Dangi <sup>c</sup>

<sup>a</sup> Ginger Research Program, Kapurkot, Salyan, Nepal.

<sup>b</sup> Nepal Agricultural Research Council, Khumaltar, Nepal.

<sup>c</sup> Grain Legumes Research Program, Khajura, Banke, Nepal.

## Authors' contributions

This work was carried out in collaboration among all authors. Authors AP and SM arranged the research concept, design, data analysis and writing the manuscript. Author SRD had contributed recording and entry a data of the experiment. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJRCS/2023/v8i3177

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98753>

Original Research Article

Received: 07/02/2023

Accepted: 09/04/2023

Published: 10/04/2023

## ABSTRACT

**Aims:** An unsuitable sowing time and selection of inappropriate variety were the main yield limiting factors in pigeonpea (*Cajanus cajan* L. Millsp.) cultivation. Therefore, a field experiment was conducted to study the effect of sowing dates on growth and yield of pigeonpea varieties.

**Study Design:** The experiment was laid out in split plot design.

**Place and Duration of Study:** This experiment was conducted at the Grain Legumes Research Program (GLRP), Khajura, Banke, Nepal in two consecutive years, 2020/21 and 2021/22.

**Methodology:** The experiment consisted five sowing dates treatments viz., 2<sup>nd</sup> July, 17<sup>th</sup> July, 1<sup>st</sup> August, 16<sup>th</sup> August and 31<sup>st</sup> August as the main-factor and two promising varieties viz., ICPL-88039 and MA-6 as the sub-factor, with three replications.

**Results:** Both the sowing time and variety significantly influenced all the tested growth and yield parameters of pigeonpea at one percent significance level. Early sowing (2<sup>nd</sup> July) recorded longer days to flowering and maturity and taller plants that affected the yield. Pigeonpea produced the

\*Corresponding author: E-mail: [anilp.narc@gmail.com](mailto:anilp.narc@gmail.com), [ngrp.narc@gmail.com](mailto:ngrp.narc@gmail.com);

highest ( $P \leq 0.01$ ) seed yield (1482 kg/ha), while seeded on 2<sup>nd</sup> July. Moreover, the pigeonpea sown on 2<sup>nd</sup> to 17<sup>th</sup> July produced 36% more seed yield than sown on 1<sup>st</sup> to 31<sup>st</sup> August. Similarly, a promising variety of pigeonpea MA-6 (1028 kg/ha) produced significantly higher ( $P \leq 0.01$ ) seed yield compared to the variety ICPL-88039 (888 kg/ha).

**Conclusion:** The study identified the 2<sup>nd</sup> July as the most appropriate time of sowing and the MA-6 as a high yielding variety of pigeonpea for the study area in Mid-Western Terai Region of Nepal.

*Keywords: Pigeonpea; sowing time; variety; yield.*

## 1. INTRODUCTION

Pigeonpea (*Cajanus cajan* L. Millsp.) is a hardy and widely adopted drought tolerant crop of the South Asia, and a most important crop in the context of climate change. It is grown in an area about 6.4 million ha in the world with a production of 5.5 million t, but the South Asia contributes an about 83 and 86% of the total world area and its production, respectively [1]. Cultivation of pigeonpea in a tropical and sub-tropical regions is not only an economical benefits for harvesting the grains, but has tremendous important for improving marginal soils by shedding its large amount of its leaves and fixing atmospheric nitrogen by nodulation [2].

Pigeonpea is the most important legume crop in Nepal and its' cultivation is gaining popularity from Terai to Hilly Regions. Pigeonpea is grown in about 16591 ha with the production and productivity of 16649 t and 1.0 t/ha, respectively [3]. It is mostly grown in upland or marginal lands under the rainfed condition in both Terai and Hilly regions of the country. The data over the 15 years indicates the increasing trend in its cultivated area, but the productivity seemed very low and stagnant at less than 1.0 t/ha [4]. There are several constraints like drought, unavailability of suitable variety, inappropriate time of sowing, poor plant population, etc. for the lower productivity of pigeonpea [5,6]. The study by [7-9] also reported the severe yield reduction of pigeonpea due to the inappropriate sowing time and variety. An appropriate sowing time and selection of suitable varieties are the most important factors for improving the productivity of any crops, mostly in legume crops [10].

Consequently, it is important to study the effect of sowing times and varieties on the growth and yield parameters of pigeonpea in Nepal. Therefore, the study was conducted with the objective for improving the pigeonpea yield by identifying the suitable time of sowing with best suited variety.

## 2. MATERIALS AND METHODS

A field experiment was conducted at agricultural farm of Nepal Agricultural Research Council, Grain Legumes Research Program (GLRP), Khajura, Banke, Nepal situated at 28° 06' 45" N latitude, 81° 35' 58" E longitude and 182 meters above Mean Sea Level in two consecutive years, 2020/21 (Year I) and 2021/22 (Year II). It was laid out in a split-plot design involving five sowing dates viz., D<sub>1</sub>: 2<sup>nd</sup> July, D<sub>2</sub>: 17<sup>th</sup> July, D<sub>3</sub>: 1<sup>st</sup> August, D<sub>4</sub>: 16<sup>th</sup> August and D<sub>5</sub>: 31<sup>st</sup> August as the main-factor, and two promising varieties viz., V<sub>1</sub>: ICPL-88039 and V<sub>2</sub>: MA-6 as the sub-factor with ten treatments combinations in three replications. The size of the experiment plot was 9 m<sup>2</sup>, where the crop was sown at a row spacing of 75 cm and 25 cm plant to plant spacing. Fertilizers were applied @ 20:40:20 kg N:P<sub>2</sub>O<sub>5</sub>:K<sub>2</sub>O per ha at the time of sowing and other cultural practices were applied as per needed. Harvesting was done manually at physiological maturity stage.

The soil characteristics of experimental plots was sandy loam with neutral in pH consisting of a low amount of organic matter (1.76%) and nitrogen (0.08%) but a high level of phosphorous (138.7 kg/ha) and potassium (283.5 kg/ha). All the data on yield, growth and development parameters were analyzed statistically at probability level of  $P \leq .05$  by using statistical software SPSS version 16.0.

## 3. RESULTS AND DISCUSSION

### 3.1 Effect of Sowing Date

The effect of sowing dates and varieties on growth, development and yield of pigeonpea are presented in Tables 2 and 3. A combined result indicated a significant effect at  $P \leq 0.01$  in flowering days, maturity days, plant height and seed yield due to the effect of sowing dates (Table 3). The similar results on the all recorded growth and yield parameters were shown in both the experimental years (Table 2).

**Table 1. The physical and chemical properties of soil of the study area**

Particular	Organic matter (%)	Nitrogen (%)	Phosphorous (kg/ha)	Potassium (kg/ha)
Soil sample I	1.76	0.08	138.7	283.5

The pooled analysis showed significantly longer days to flowering in pigeonpea under early sown condition *i.e.*, 2<sup>nd</sup> July (188 days) followed by 17<sup>th</sup> July (183 days). Similarly, in both the years, pigeonpea took longer duration to flower when seeded on 2<sup>nd</sup> July. It required 166 days (31<sup>st</sup> August) to 185 days (2<sup>nd</sup> July) in the Year I and 160 days (31<sup>st</sup> August) to 192 days (2<sup>nd</sup> July) in the Year II, for 50% flowering. Further the maturity days ranged from 213 days (31<sup>st</sup> August) to 247 days (2<sup>nd</sup> July) in the Year I, while it ranged from 213 (31<sup>st</sup> August) to 260 days (2<sup>nd</sup> July) in the Year II. Similarly, a longer day to mature was recorded when seeded on 2<sup>nd</sup> July to 1<sup>st</sup> August (254 days) under pooled analysis (Table 3).

The pigeonpea sown on 2<sup>nd</sup> July to 1<sup>st</sup> August (251 to 243 cm) had significantly taller plants than sown on 16<sup>th</sup> August to 31<sup>st</sup> August (150 to 143 cm), based on the combined analysis (Table 3). During the Year II, the plant height ranged from 141 cm (16<sup>th</sup> August) to 252 cm (2<sup>nd</sup> July), while in Year II it ranged from 138 cm (31<sup>st</sup> August) to 250 cm (2<sup>nd</sup> July) (Table 2).

A combined result of the Year I and Year II recorded a significant higher seed yield at  $P \leq 0.01$  when sown on 2<sup>nd</sup> July (1482 kg/ha) followed by 17<sup>th</sup> July sown conditions (1279 kg/ha), whereas pigeonpea sowing on 31<sup>st</sup> August produced the lowest seed yield (411 kg/ha) followed by 16<sup>th</sup> August (593 kg/ha). In both years of the experiment, a significantly higher seed yield of pigeonpea was observed with 2<sup>nd</sup> July to 17<sup>th</sup> July sowing conditions. The seed yield of pigeonpea ranged from 390 kg/ha (31<sup>st</sup> August) to 1575 kg/ha (2<sup>nd</sup> July) in Year I, while from 432 kg/ha (31<sup>st</sup> August) to 1399 kg/ha (17<sup>th</sup> July) in Year II.

In the study, earlier flowering (15 to 25 days) and maturity (34 to 40 days), shorter plant height (90 to 108 cm) and three-fold lower yield of pigeonpea was recorded under late sown conditions. This might be due to the compression of growing periods that hinder the proper growth and development of the crops. However, the earlier sown condition provides the ample time and favorable climatic conditions, especially light, temperature, *etc.* during growth, development

and maturity stages for proper crop growth and development. Similar observation was reported by Fukugawa et al. [11] where heavy vegetative growth was followed by more flowers in early sown crop. The late sowing pigeonpea affects seed germination due to higher soil moisture and affects yield due to lower temperature coupled with low rainfall during reproductive and maturity stages [12,13]. The shorter plant and longer days for flowering and maturity and low yield of pigeonpea in late sown as compared to early sown condition were also observed by different authors [5,6,9], where they reported 37%, 397% and 7% more yield under earlier sown condition than later sown condition, respectively. The results of the study are conformity with the findings of [7,13,14], where they found early sowing of pigeonpea was more productive with regard to growth and yield parameters.

### 3.2 Effect of Variety

The tested promising varieties of pigeonpea had significant effect at  $P \leq 0.01$  on its flowering and maturity time, plant height and seed yield, however in Year I plant height did not differ significantly. The variety of pigeonpea *i.e.*, MA-6 possessed the longer duration to flower (193 days) and maturity (248 days), taller plant (214 cm) and higher seed yield (1028 t/ha), as compared to ICPL-8803. The days to flower, days to maturity, plant height and seed yield were found to be 152 days, 229 days, 197 cm and 888 kg/ha, respectively in ICPL-88039. In both Year I and Year II, the days to flowering, days to maturity, plant height and seed yield in both varieties were noted at 150–194 days and 154–193 days, 220–249 days and 239–248 days, 200–211 cm and 195–216 cm, 866–1024 kg/ha and 911–1031 kg/ha, respectively.

The pigeonpea variety MA-6 required more number of days *i.e.*, 41 days and 19 days for flowering and maturity as compared to ICPL-88039. The long duration variety (MA-6) was 17 cm taller and had 16% more seed yield compared with early maturing variety (ICPL-88039). The differential flowering time, maturing time, plant height and seed yield of the varieties was due to the disparity in their genetic makeup and characters. A study by Kithan et al. [7] also

reported the different flowering and maturing days, plant height and seed yield due to the effect of varietal characteristics in pigeonpea production, where they noted higher yield of variety UPAS 120 (969 kg/ha), among the three tested varieties. The yield variation in pigeonpea


due to varietal difference was also noted by Kuri, Chawhan and Rani et al. [5,9,14] in their study. A researcher [15] reported that seed yield of pigeonpea depends on source and sink relation which in turn depends on environmental factors.

**Table 2. Effects of sowing time and variety on growth attributes and yield of pigeonpea in two different years**

Treatment	Days to flowering		Days to maturity		Plant height, cm		Seed yield, kg ha <sup>-1</sup>	
	Year I	Year II	Year I	Year II	Year I	Year II	Year I	Year II
<b>Sowing time</b>								
2 <sup>nd</sup> July	185	192	247	260	252	250	1575	1389
17 <sup>th</sup> July	181	186	246	262	238	242	1158	1399
1 <sup>st</sup> August	168	169	250	258	248	238	963	1090
16 <sup>th</sup> August	160	163	216	225	141	160	641	545
31 <sup>st</sup> August	166	160	213	213	149	138	390	432
SEm (±)	1.30	3.36	7.01	1.80	10.48	2.32	61.84	61.00
LSD (0.05)	3.85	10.00	20.84	5.34	31.13	6.89	183.73	181.23
<b>Variety</b>								
ICPL-88039	150	154	220	239	200	195	866	911
MA-6	194	193	249	248	211	216	1024	1031
SEm (±)	0.82	2.13	4.44	1.14	6.63	1.47	39.11	38.58
LSD (0.05)	2.44	6.32	13.18	3.38	19.69	4.36	116.20	114.62
<b>F-test</b>								
Sowing time	**	**	**	**	**	**	**	**
Variety	**	**	**	**	ns	**	**	*
Sowing time x Variety	**	**	ns	ns	ns	**	ns	ns
CV%	1.8	4.7	7.3	1.8	12.5	2.8	16.6	15.4

**Table 3. Effects of sowing time and variety on growth attributes and yield of pigeonpea (Pooled data of two years)**

Treatment	Days to flowering	Days to maturity	Plant height, cm	Seed yield, kg ha <sup>-1</sup>
<b>Sowing time</b>				
2 <sup>nd</sup> July	188	254	251	1482
17 <sup>th</sup> July	183	254	240	1279
1 <sup>st</sup> August	168	254	243	1026
16 <sup>th</sup> August	162	220	150	593
31 <sup>st</sup> August	163	213	143	411
SEm (±)	1.80	3.79	5.70	42.94
LSD (0.05)	5.16	10.86	16.32	122.93
<b>Variety</b>				
ICPL-88039	152	229	197	888
MA-6	193	248	214	1028
SEm (±)	1.14	2.40	3.61	27.16
LSD (0.05)	3.27	6.87	10.32	77.75
<b>F-test</b>				
Year	ns	**	ns	ns
Sowing time	**	**	**	**
Variety	**	**	**	**
Sowing time x Variety	**	ns	ns	ns
CV%	3.6	5.5	9.6	15.5



**Fig. 1. Interaction effects of sowing time and variety on seed yield of pigeonpea (F-test = ns; LSD (0.05) = 173.9)**

### 3.3 Interaction Effect

The interaction effect of sowing time and variety showed non-significant difference on the seed yield of pigeonpea (Fig. 1). However, the results of the experiments indicated that the highest yield of pigeonpea was found from variety MA-6 with the sowing time of 2<sup>nd</sup> July (1578 kg/ha), while recorded lowest from variety ICPL-88039 (371 kg/ha) seeded on 31<sup>st</sup> August. Both the tested variety performed well under early sowing condition and poor under late sown condition. Similar observation, a non-significant interaction effect of variety and sowing date on pigeonpea was also reported by Ram et al. [16].

### 4. CONCLUSION

Selection of appropriate variety and sowing time is the most important factor for pigeonpea production. A high yielding variety sown at the right time could help to utilize all the production inputs efficiently and optimize yield. Based on the above findings it can be concluded that the suitable time for pigeonpea sowing is the month of July, after which yield reduced radically. The pigeonpea sown on July produced about 50% more yield as compared to sowing on August in the study area. Similarly, a long duration variety performed better compared to a short duration variety for the pigeonpea cultivation in Mi-West Region of Nepal. However, more research can be carried out for more detailed study of the findings.

### ACKNOWLEDGEMENTS

The authors gratefully acknowledge the necessary support received from the climate-smart technology development in grain legumes project of Nepal Agricultural Research Council (NARC), Nepal.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. FAOSTAT. Crop and livestock products. 2023; Accessed 12 March 2023. Available: <https://www.fao.org/faostat/en/#data/qcl>.
2. Rao K and Dart PJ. Nodulation, nitrogen fixation and nitrogen uptake in pigeonpea (*Cajanus cajan* L. Millsp) of different maturity groups. Plant and Soil. 1987;99: 255-266.
3. MoALD. Statistical Information on Nepalese Agriculture (2020/21). Government of Nepal, Ministry of Agriculture and Livestock Development. Planning and Development Coordination Division, Statistics and Analysis Section, Singhdurbar, Kathmandu, Nepal; 2022. Accessed 16 March 2023. Available: <https://moald.gov.np/wp-content/uploads/2022/07/statistical-information-on-nepalese-agriculture-2077-78.pdf>

4. Pokhrel A, Aryal L and Poudel PP. A review on research work of Grain Legumes Research Program, NARC, NARC Publication serial number:00686-500/2017/18; 2018.
5. Kuri S, Shivaramu HS, Thimmegowda MN, Yogananda SB, Prakash SS and Murukannappa. Effect of Row Spacing, Varieties and Sowing Dates on Growth and Yield of Pigeonpea. International Journal of Current Microbiology and Applied Sciences. 2018;7(8):1123-1128. DOI: <https://doi.org/10.20546/ijcmas.2018.708.127>.
6. Kittur CN and Guggari AK. Effect of sowing time and planting geometry on the growth and yield of pigeonpea in Northern Dry Zone (Zone 3) of Karnataka. Journal of Farm Science. 2017;334-337.
7. Kithan L, Sharma MB and Longchar A. Effect of Planting Dates on the Performance of Promising Pigeonpea Genotypes under NEHZ. International Journal of Economic Plants. 2020;7:6-8.
8. Chopade BJ, Mehendale SK, Desai VS, Narangalkar AL and Sapkal SD. Effect of different dates of sowing on pod borer complex of pigeon pea, (*Cajanus cajan* (L.) Mill Sp.). Journal of Entomology and Zoology Studies. 2020;8(5):1048-1051.
9. Chawhan, RG, Chahande RV and Deshmukh HS. Effect of sowing date on seed quality of pigeonpea [*Cajanus cajan* (L.) Mill sp.]. The Pharma Innovation Journal. 2019;8(7):784-789.
10. Pokhrel A and Dangi SR. Increasing the productivity of rajma through proper sowing date and plant geometry. Journal of Agriculture and Natural Resources. 2022;5(1):12-18. DOI:<https://doi.org/10.3126/janr.v5i1.50348>
11. Fukugawa Y and Zhenga SUH. Growth response to sowing dates of pigeonpea in northern kyushu of Japan. Japanese Journal of Crop Science. 1999;68:33-38.
12. Dhanoji MM and Patil JR. Effect of Date of Sowing on Potential Source and Sink Realization in Pigeon Pea. Environment and Ecology. 2011;29(3):1003-1005.
13. Kumar N, Gopinath KA, Srivastva AK and Mahajan V. Performance of pigeonpea (*Cajanuscajan* (L.) Millsp.) at different sowing dates in the mid-hills of Indian Himalaya. Archives of Agronomy and Soil Science. 2008;54 (5):507–514.
14. Rani BP and Reddy RD. Performance of pigeonpea in sole and intercropping system in vertisols of Krishna - Godavari zone in Andhra Pradesh. Indian Journal of Agricultural Research. 2010;44(3): 225–228.
15. Singh I. Flowering and podding behavior in determinate and indeterminate pigeonpea genotypes. Indian Journal of Agricultural Research. 2000;34(1):67-70.
16. Ram H, Singh G, Sekhon HS and Khanna V. Effect of sowing time on the performance of pigeonpea genotypes. Journal of Food Legumes. 2011;24(3): 207-210.

© 2023 Pokhrel et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/98753>